The New Prime theorem（28）

$$P_1 = P \pm m \quad \text{and} \quad P_1 = 2P \pm n$$

Chun-Xuan Jiang

P.O.Box3924, Beijing100854, P.R. China. Jiangchunxuan@vip.sohu.com

Abstract: Using Jiang function we prove that $P_1 = P \pm m$ and $P_1 = 2P \pm n$ have infinitely many prime solutions.

Keywords: prime; theorem; function; number; new

Theorem 1. Let m be an even number.

$$P_1 = P + m \tag{1}$$

has infinitely many prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P - 1 - \chi(P)], \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$q + m \equiv 0 \pmod{P}, q = 1, \cdots, P - 1. \tag{3}$$

If $P \mid m$ then $\chi(P) = 0$. $\chi(P) = 1$ othewise.

Substituting it into (2) we have.

$$J_2(\omega) = \prod_{3 \le P} (P - 2) \prod_{P \mid m} \frac{P - 1}{P - 2} \neq 0 \tag{4}$$

We prove that (1) has infinitely many primes solutions.

We have asymptotic formula [1,2]

$$\pi_2(N, 2) = \left| \{ P \le N : P_1 = prime \} \right| \sim \frac{J_2(\omega)\omega}{\phi^2(\omega)} \frac{N}{\log^2 N} = 2 \prod_{3 \le P} \left(1 - \frac{1}{(P-1)^2} \right) \prod_{P \mid m} \frac{P - 1}{P - 2} \frac{N}{\log^2 N}. \tag{5}$$

Where $\phi(\omega) = \prod_P (P - 1)$.

In the same way we are able to prove that $P_1 = P - m$ has infinitely many prime solutions.

Theorem 2. Let n be an odd number.

$$P_1 = 2P + n$$

has infinitely many prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P - 1 - \chi(P)], \tag{6}$$

where $\chi(P)$ is the number of solutions of congruence

$$2q + n \equiv 0 \pmod{P} \tag{7}$$

$$q = 1, \cdots, P - 1.$$

If $P \mid n$ then $\chi(P) = 0$. $\chi(P) = 1$ otherwise.

Substituting it into (6) we have

$$J_2(\omega) = \prod_{3 \le P}(P-2)\prod_{P|n}\frac{P-1}{P-2} \ne 0$$

（8）

We prove that (6) has infinitely many prime solutions.

We have asymptotic formula [1,2]

$$\pi_2(N,2) = \left|\{P \le N : P_1 = prime\}\right| \sim \frac{J_2(\omega)\omega}{\phi^2(\omega)}\frac{N}{\log^2 N}$$

（9）

Remark. The prime number theory is basically to count the Jiang function $J_{n+1}(\omega)$ and Jiang prime k-tuple singular series $\sigma(J) = \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} = \prod_P\left(1 - \frac{1+\chi(P)}{P}\right)\left(1 - \frac{1}{P}\right)^{-k}$ [1,2], which can count the number of prime number. The prime distribution is not random. But Hardy prime k-tuple singular series $\sigma(H) = \prod_P\left(1 - \frac{\nu(P)}{P}\right)\left(1 - \frac{1}{P}\right)^{-k}$ is false [3-8], which cannot count the number of prime numbers.

Szemerdi's theorem does not directly to the primes, because it can not count the number of primes. It is unusable. Cramr's random model can not prove prime problems. It is incorrect. The probability of $1/\log N$ of being prime is false. Assuming that the events "P is prime", "$P+2$ is prime" and "$P+4$ is prime" are independent, we conclude that P, $P+2$, $P+4$ are simultaneously prime with probability about $1/\log^3 N$. There are about $N/\log^3 N$ primes less than N. Letting $N \to \infty$ we obtain the prime conjecture, which is false. The tool of additive prime number theory is basically the Hardy-Littlewood prime tuple conjecture, but can not prove and count any prime problems[6].

Mathematicians have tried in vain to discover some order in the sequence of prime numbers but we have every reason to believe that there are some mysteries which the human mind will never penetrate. Leonhard Euler

It will be another million years, at least, before we understand the primes. Paul ErdÖs

References
1. Chun-Xuan Jiang, Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture. Inter. Acad. Press, 2002, MR2004c:11001, (http://www.i-b-r.org/docs/jiang.pdf) (http://www.wbabin.net/math/xuan13.pdf)(http://vixra.org/numth/).
2. Chun-Xuan Jiang, Jiang's function $J_{n+1}(\omega)$ in prime distribution.(http://www. wbabin.net/math /xuan2. pdf.) (http://wbabin.net/xuan.htm#chun-xuan) (http://vixra.org/numth/).
3. Chun-Xuan Jiang, The Hardy-Littlewood prime k-tuple conjectnre is false.(http://wbabin.net/xuan.htm# chun-xuan)(http://vixra.org/numth/).
4. G. H. Hardy and J. E. Littlewood, Some problems of "Partitio Numerorum", III: On the expression of a number as a sum of primes. Acta Math., 44(1923)1-70.
5. W. Narkiewicz, The development of prime number theory. From Euclid to Hardy and Littlewood. Springer-Verlag, New York, NY. 2000, 333-353. 这是当代素数理论水平.
6. B. Green and T. Tao, Linear equations in primes. To appear, Ann. Math.
7. D. Goldston, J. Pintz and C. Y. Yildirim, Primes in tuples I. Ann. Math., 170(2009) 819-862.
8. T. Tao. Recent progress in additive prime number theory, preprint. 2009. http://terrytao.files.wordpress. com/2009/08/prime-number-theory 1.pdf.
9. Vinoo Cameron. Prime Number 19, The Vedic Zero And The Fall Of Western Mathematics By Theorem. *Nat Sci* 2013;11(2):51-52. (ISSN: 1545-0740). http://www.sciencepub.net/nature/ns1102/009_15631ns1102_51_52.pdf.
10. Vinoo Cameron, Theo Den otter. PRIME NUMBER COORDINATES AND CALCULUS. *Rep Opinion* 2012;4(10):16-17. (ISSN: 1553-9873). http://www.sciencepub.net/report/report0410/004_10859report0410_16_17.pdf.
11. Vinoo Cameron, Theo Den otter. PRIME NUMBER COORDINATES AND CALCULUS. *J Am Sci* 2012;8(10):9-10. (ISSN: 1545-1003). http://www.jofamericanscience.org/journals/am-sci/am0810/002_10859bam0810_9_10.pdf.
12. Chun-Xuan Jiang. Automorphic Functions And Fermat's Last Theorem (1). *Rep Opinion* 2012;4(8):1-6. (ISSN: 1553-9873). http://www.sciencepub.net/report/report0408/001_10009report0408_1_6.pdf.
13. Chun-Xuan Jiang. The Hardy-Littlewood prime k-tuple conjecture is false. *Rep Opinion* 2012;4(8):35-38. (ISSN: 1553-9873). http://www.sciencepub.net/report/report0408/008_10016report0408_35_38.pdf.
14. Chun-Xuan Jiang. A New Universe Model. *Academ Arena* 2012;4(7):12-13 (ISSN 1553-992X). http://sciencepub.net/academia/aa0407/003_10067aa0407_12_13.pdf.

5/1/2015