The New Prime theorem (42)

On the
$$P, jP^7 + k - j(j = 1, \dots, k-1)$$

Chun-Xuan Jiang

P. O. Box 3924, Beijing 100854, P. R. China jiangchunxuan@vip.sohu.com

Abstract: Using Jiang function we prove that $jP^7 + k - j$ contain infinitely many prime solutions.

[Chun-Xuan Jiang. The New Prime theorem (42) On the $P, jP^7 + k - j(j = 1, \dots, k-1)$. Academ Arena 2015;7(1s): 79-80]. (ISSN 1553-992X). http://www.sciencepub.net/academia. 42

Keywords: prime; theorem; function; number; new

Theorem . Let k be a given prime.

$$P, jP^7 + k - j(j = 1, \dots, k - 1)$$
 (1)

We have Jiang function [1,2]

$$J_{2}(\omega) = \prod_{P} [P - 1 - \chi(P)] \tag{2}$$

where $\omega = \prod_{P} P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^7 + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P - 1$$
(3)

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

we prove that (1) contain infinitely many prime solutions.

We have asymptotic formula [1,2]

$$\pi_{k}(N,2) = \left| \left\{ P \le N : jP^{7} + k - j = prime \right\} \right| \sim \frac{J_{2}(\omega)\omega^{k-1}}{(7)^{k-1}\phi^{k}(\omega)} \frac{N}{\log^{k}N}$$
Where
$$\phi(\omega) = \prod_{P} (P-1)$$
(5)

Remark. The prime number theory is basically to count the Jiang function $J_{n+1}(\omega)$ and Jiang prime k-tuple

$$\sigma(J) = \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} = \prod_P \left(1 - \frac{1 + \chi(P)}{P}\right) \left(1 - \frac{1}{P}\right)^{-k}$$
 [1,2], which can count the number of prime

numbers. The prime distribution is not random. But Hardy-Littlewood prime k -tuple singular series $\sigma(H) = \prod_{P} \left(1 - \frac{\nu(P)}{P} \right) (1 - \frac{1}{P})^{-k}$ is false [3-8], which cannot count the number of prime numbers[3].

References

- Chun-Xuan Jiang, Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and conjecture. Inter. Acad. Press, 2002, MR2004c:11001, (http://www.i-b-r.org/docs/jiang.pdf) (http://www.wbabin.net/math/xuan13.pdf) (http://vixra.org/numth/).
- Chun-Xuan Jiang, Jiang's function $J_{n+1}(\omega)$ in prime distribution.(http://www.wbabin.net/math/xuan2.pdf.) (http://wbabin.net/xuan.htm#chun-xuan) (http://vixra.org/numth/).
- Chun-Xuan Jiang, The Hardy-Littlewood prime k -tuple conjecture is false.(http://wbabin.net/xuan.htm# 3.

- chun-xuan)(http://vixra.org/numth/).
- 4. G. H. Hardy and J. E. Littlewood, Some problems of "Partitio Numerorum", III: On the expression of a number as a sum of primes. Acta Math., 44(1923)1-70.
- W. Narkiewicz, The development of prime number theory. From Euclid to Hardy and Littlewood. Springer-Verlag, New York, NY. 2000, 333-353.
- 6. B. Green and T. Tao, Linear equations in primes. To appear, Ann. Math.
- 7. D. Goldston, J. Pintz and C. Y. Yildirim, Primes in tuples I. Ann. Math., 170(2009) 819-862.
- 8. T. Tao. Recent progress in additive prime number theory, preprint. 2009. http://terrytao.files.wordpress.com/2009/08/prime-number-theory1.pdf.
- 9. Vinoo Cameron. Prime Number 19, The Vedic Zero And The Fall Of Western Mathematics By Theorem. *Nat Sci* 2013;11(2):51-52. (ISSN: 1545-0740). http://www.sciencepub.net/nature/ns1102/009 15631ns1102 51 52.pdf.
- 10. Vinoo Cameron, Theo Den otter. PRIME NUMBER COORDINATES AND CALCULUS. *Rep Opinion* 2012;4(10):16-17. (ISSN: 1553-9873). http://www.sciencepub.net/report/report/410/004 10859report0410 16 17.pdf.
- 11. Vinoo Cameron, Theo Den otter. PRIME NUMBER COORDINATES AND CALCULUS. J Am Sci 2012;8(10):9-10. (ISSN: 1545-1003). http://www.jofamericanscience.org/journals/am-sci/am0810/002 10859bam0810 9 10.pdf.
- 12. Chun-Xuan Jiang. Automorphic Functions And Fermat's Last Theorem (1). *Rep Opinion* 2012;4(8):1-6. (ISSN: 1553-9873). http://www.sciencepub.net/report/408/001/10009report0408/1/6.pdf.
- 13. Chun-Xuan Jiang. Jiang's function $J_{n+1}(\omega)$ in prime distribution. Rep Opinion 2012;4(8):28-34. (ISSN: 1553-9873). http://www.sciencepub.net/report/report/0408/007 10015report0408 28 34.pdf.
- 14. Chun-Xuan Jiang. The Hardy-Littlewood prime *k*-tuple conjecture is false. *Rep Opinion* 2012;4(8):35-38. (ISSN: 1553-9873), http://www.sciencepub.net/report/report/d08/008 10016report0408 35 38.pdf.
- Chun-Xuan Jiang. A New Universe Model. Academ Arena 2012;4(7):12-13 (ISSN 1553-992X). http://sciencepub.net/academia/aa0407/003 10067aa0407 12 13.pdf.

Szemerédi's theorem does not directly to the primes, because it cannot count the number of primes. Cramér's random model cannot prove any prime problems. The probability of $1/\log N$ of being prime is false. Assuming that the events "P is prime", "P+2 is prime" and "P+4 is prime" are independent, we conclude that P, P+2, P+4 are simultaneously prime with probability about $1/\log^3 N$. There are about $N/\log^3 N$ primes less than N. Letting $N \to \infty$ we obtain the prime conjecture, which is false. The tool of additive prime number theory is basically the Hardy-Littlewood prime tuples conjecture, but cannot prove and count any prime problems[6].

Mathematicians have tried in vain to discover some order in the sequence of prime numbers but we have every reason to believe that there are some mysteries which the human mind will never penetrate.

Leonhard Euler (1707-1783)

It will be another million years, at least, before we understand the primes.

Paul Erdos (1913-1996)

Of course, the primes are a deterministic set of integers, not a random one, so the predictions given by random models are not rigorous (Terence Tao, Structure and randomness in the prime numbers, preprint). 陶哲轩认为素数不是随机的。

Erdos and Turán (1936) contributed to probabilistic number theory, where the primes are treated as if they were random, which generates Szemerédi's theorem (1975) and Green-Tao theorem (2004). But they cannot actually prove and count any simplest prime examples: twin primes and Goldbach's conjecture. They don't know what prime theory means, only conjectures.

5/1/2015