

The New Prime theorems (1191)— (1240)

Jiang, Chun-Xuan (蒋春暄)

Institute for Basic Research, Palm Harbor, FL34682-1577, USA

And: P. O. Box 3924, Beijing 100854, China (蒋春暄, 北京 3924 信箱, 100854)

jiangchunxuan@sohu.com, cxjiang@mail.bcf.net.cn, jcxuan@sina.com, Jiangchunxuan@vip.sohu.com,
jcxxxx@163.com

Abstract: Using Jiang function we are able to prove almost all prime problems in prime distribution. This is the Book proof. No great mathematicians study prime problems and prove Riemann hypothesis in AIM, CLAYMA, IAS, THES, MPIM, MSRI. In this paper using Jiang function $J_2(\omega)$ we prove that the new prime theorems (1191)-(1240) contain infinitely many prime solutions and no prime solutions. From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$. This is the Book theorem.

[Jiang, Chun-Xuan (蒋春暄). **The New Prime theorems (1191)— (1240)**. *Academ Arena* 2016;8(1s): 1136-1203]. (ISSN 1553-992X). <http://www.sciencepub.net/academia>. 20. doi:[10.7537/marsaai0801s1620](https://doi.org/10.7537/marsaai0801s1620).

Keywords: new; prime theorem; Jiang Chunxuan; mathematics; science; number; function

It will be another million years at least, before we understand the primes.

Paul Erdos (1913-1996)

The New Prime Theorem (1191)

$$p, jp^{2302} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2302} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2302} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2302} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2302} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2302} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2302)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1192)

$$p, jp^{2304} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2304} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2304} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2304} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2304} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2304} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2304)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \tag{6}$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$

Example 1. Let $k = 3, 5, 7, 13, 17, 19, 37, 73, 97, 193, 577, 769, 1153$

From (2) and (3) we have

$$J_2(\omega) = 0 \tag{7}$$

We prove that for $k = 3, 5, 7, 13, 17, 19, 37, 73, 97, 193, 577, 769, 1153$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 7, 13, 17, 19, 37, 73, 97, 193, 577, 769, 1153$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{8}$$

We prove that for $k \neq 3, 5, 7, 13, 17, 19, 37, 73, 97, 193, 577, 769, 1153$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1193)

$$p, jp^{2306} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang
 jiangchunxuan@vip.sohu.com
 Abstract

Using Jiang function we prove that $jp^{2306} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2306} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2306} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2306} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2306} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2306)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1194)

$$p, jp^{2308} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2308} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2308} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2308} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2308} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2308} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2308)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 2309$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 2309$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 2309$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 2309$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1195)

$$p, jp^{2310} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2310} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2310} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2310} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2310} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2310} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2310)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \tag{6}$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$

Example 1. Let $k = 3, 7, 11, 23, 31, 71, 211, 331, 2311$

From (2) and (3) we have

$$J_2(\omega) = 0 \tag{7}$$

We prove that for $k = 3, 7, 11, 23, 31, 71, 211, 331, 2311$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 7, 11, 23, 31, 71, 211, 331, 2311$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{8}$$

We prove that for $k \neq 3, 7, 11, 23, 31, 71, 211, 331, 2311$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1196)

$$p, jp^{2312} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang
 jiangchunxuan@vip.sohu.com
 Abstract

Using Jiang function we prove that $jp^{2312} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2312} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2312} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2312} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2312} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2312)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 137$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 137$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 137$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 137$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1197)

$$p, jp^{2314} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2314} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2314} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2314} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2314} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2314} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2314)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 179$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 179$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 179$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 179$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1198)

$$p, jp^{2316} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2316} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2316} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2316} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2316} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2316} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2316)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 7, 13, 773$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 7, 13, 773$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 7, 13, 773$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 7, 13, 773$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1199)

$$p, jp^{2318} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2318} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2318} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2318} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2318} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2318} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2318)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1200)

$$p, jp^{2320} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2320} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2320} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2320} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2320} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2320} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2320)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \tag{6}$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$

Example 1. Let $k = 3, 5, 11, 17, 41, 59, 233$

From (2) and (3) we have

$$J_2(\omega) = 0 \tag{7}$$

We prove that for $k = 3, 5, 11, 17, 41, 59, 233$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 11, 17, 41, 59, 233$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{8}$$

We prove that for $k \neq 3, 5, 11, 17, 41, 59, 233$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1201)

$$p, jp^{2322} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang
 jiangchunxuan@vip.sohu.com
 Abstract

Using Jiang function we prove that $jp^{2322} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2322} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2322} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2322} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2322} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2322)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 7, 19, 173$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 7, 19, 173$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 7, 19, 173$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 7, 19, 173$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1202)

$$p, jp^{2324} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2324} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2324} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2324} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2324} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2324} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2324)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 29, 1163$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 29, 1163$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 29, 1163$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 29, 1163$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1203)

$$p, jp^{2326} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2326} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2326} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2326} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2326} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2326} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2326)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1204)

$$p, jp^{2328} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2328} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2328} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2328} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2328} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2328} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2328)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 7, 13, 389$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 7, 13, 389$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 7, 13, 389$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 7, 13, 389$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1205)

$$p, jp^{2330} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2330} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2330} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2330} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2330} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2330} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2330)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 11, 467$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 11, 467$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 11, 467$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 11, 467$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1206)

$$p, jp^{2332} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2332} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2332} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2332} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2332} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2332} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2332)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 23, 107, 2333$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 23, 107, 2333$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 23, 107, 2333$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 23, 107, 2333$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1207)

$$p, jp^{2334} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2334} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2334} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2334} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2334} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2334} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2334)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 7$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 7$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 7$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 7$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1208)

$$p, jp^{2336} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2336} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2336} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2336} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2336} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2336} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2336)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 17, 293$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 17, 293$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 17, 293$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 17, 293$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1209)

$$p, jp^{2338} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2338} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2338} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2338} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2338} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2338} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2338)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 2339$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 2339$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 2339$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 2339$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1210)

$$p, jp^{2340} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2340} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2340} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2340} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2340} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2340} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2340)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 7, 11, 19, 31, 53, 61, 79, 131, 181, 1171, 2341$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 7, 11, 19, 31, 53, 61, 79, 131, 181, 1171, 2341$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 7, 11, 19, 31, 53, 61, 79, 131, 181, 1171, 2341$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 7, 11, 19, 31, 53, 61, 79, 131, 181, 1171, 2341$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1211)

$$p, jp^{2342} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2342} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2342} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2342} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2342} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2342} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2342)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1212)

$$p, jp^{2344} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2344} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2344} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2344} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2344} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2344} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2344)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 587$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 587$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 587$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 587$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1213)

$$p, jp^{2346} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2346} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2346} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2346} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2346} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2346} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2346)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \tag{6}$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$

Example 1. Let $k = 3, 7, 47, 103, 139, 2347$

From (2) and (3) we have

$$J_2(\omega) = 0 \tag{7}$$

We prove that for $k = 3, 7, 47, 103, 139, 2347$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 7, 47, 103, 139, 2347$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{8}$$

We prove that for $k \neq 3, 7, 47, 103, 139, 2347$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1214)

$$p, jp^{2348} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang
 jiangchunxuan@vip.sohu.com
 Abstract

Using Jiang function we prove that $jp^{2348} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2348} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2348} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2348} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2348} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2348)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \tag{6}$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5$

From (2) and (3) we have

$$J_2(\omega) = 0 \tag{7}$$

We prove that for $k = 3, 5$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{8}$$

We prove that for $k \neq 3, 5$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1215)

$$p, jp^{2350} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang
 jiangchunxuan@vip.sohu.com
 Abstract

Using Jiang function we prove that $jp^{2350} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2350} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2350} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2350} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2350} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2350)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \tag{6}$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$

Example 1. Let $k = 3, 11, 2351$

From (2) and (3) we have

$$J_2(\omega) = 0 \tag{7}$$

We prove that for $k = 3, 11, 2351$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 11, 2351$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{8}$$

We prove that for $k \neq 3, 11, 2351$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1216)

$$p, jp^{2352} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang
 jiangchunxuan@vip.sohu.com
 Abstract

Using Jiang function we prove that $jp^{2352} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2352} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2352} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2352} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2352} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2352)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \tag{6}$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$

Example 1. Let $k = 3, 5, 7, 13, 29, 43, 197, 337$

From (2) and (3) we have

$$J_2(\omega) = 0 \tag{7}$$

We prove that for $k = 3, 5, 7, 13, 29, 43, 197, 337$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 7, 13, 29, 43, 197, 337$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{8}$$

We prove that for $k \neq 3, 5, 7, 13, 29, 43, 197, 337$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1217)

$$p, jp^{2354} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang
 jiangchunxuan@vip.sohu.com
 Abstract

Using Jiang function we prove that $jp^{2354} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2354} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2354} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2354} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2354} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2354)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 23$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 23$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 23$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 23$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1218)

$$p, jp^{2356} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2356} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2356} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2356} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2356} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2356} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2356)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 2357$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 2357$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 2357$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 2357$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1219)

$$p, jp^{2358} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2358} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2358} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2358} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2358} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2358} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2358)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 7, 19, 263, 787$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 7, 19, 263, 787$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 7, 19, 263, 787$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 7, 19, 263, 787$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1220)

$$p, jp^{2360} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2360} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2360} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2360} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2360} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2360} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2360)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \tag{6}$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$

Example 1. Let $k = 3, 5, 11, 41$

From (2) and (3) we have

$$J_2(\omega) = 0 \tag{7}$$

We prove that for $k = 3, 5, 11, 41$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 11, 41$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{8}$$

We prove that for $k \neq 3, 5, 11, 41$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1221)

$$p, jp^{2362} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang
 jiangchunxuan@vip.sohu.com
 Abstract

Using Jiang function we prove that $jp^{2362} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2362} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2362} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2362} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2362} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2362)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1222)

$$p, jp^{2364} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2364} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2364} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2364} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2364} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2364} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2364)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 7$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 7$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 7$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 7$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1223)

$$p, jp^{2366} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2366} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2366} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2366} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2366} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2366} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2366)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1224)

$$p, jp^{2368} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2368} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2368} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2368} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2368} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2368} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2368)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 17, 149, 563$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 17, 149, 563$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 17, 149, 563$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 17, 149, 563$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1225)

$$p, jp^{2370} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2370} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2370} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2370} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2370} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2370} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2370)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 7, 11, 31, 2371$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 7, 11, 31, 2371$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 7, 11, 31, 2371$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 7, 11, 31, 2371$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1226)

$$p, jp^{2372} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2372} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2372} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2372} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2372} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2372} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2372)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 1187$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 1187$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 1187$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 1187$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1227)

$$p, jp^{2374} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2374} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2374} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2374} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2374} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2374} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2374)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1228)

$$p, jp^{2376} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2376} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2376} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2376} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2376} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2376} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2376)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$

Example 1. Let $k = 3, 5, 7, 13, 19, 23, 37, 67, 73, 89, 109, 199, 397, 2377$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 7, 13, 19, 23, 37, 67, 73, 89, 109, 199, 397, 2377$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 7, 13, 19, 23, 37, 67, 73, 89, 109, 199, 397, 2377$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 7, 13, 19, 23, 37, 67, 73, 89, 109, 199, 397, 2377$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1229)

$$p, jp^{2378} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2378} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2378} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2378} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2378} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2378} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2378)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 59, 83$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 59, 83$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 59, 83$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 59, 83$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1230)

$$p, jp^{2380} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2380} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2380} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2380} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2380} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2380} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2380)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 11, 29, 71, 239, 2381$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 11, 29, 71, 239, 2381$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 11, 29, 71, 239, 2381$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 11, 29, 71, 239, 2381$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1231)

$$p, jp^{2382} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2382} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2382} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2382} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2382} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2382} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2382)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 7, 2383$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 7, 2383$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 7, 2383$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 7, 2383$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1232)

$$p, jp^{2384} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2384} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2384} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2384} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2384} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2384} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2384)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 17, 1193$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 17, 1193$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 17, 1193$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 17, 1193$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1233)

$$p, jp^{2386} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2386} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2386} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2386} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2386} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2386} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2386)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \tag{6}$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$

Example 1. Let $k = 3$

From (2) and (3) we have

$$J_2(\omega) = 0 \tag{7}$$

We prove that for $k = 3$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{8}$$

We prove that for $k \neq 3$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1234)

$$p, jp^{2388} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang
jiangchunxuan@vip.sohu.com
Abstract

Using Jiang function we prove that $jp^{2388} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2388} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2388} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2388} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2388} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2388)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \tag{6}$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$

Example 1. Let $k = 3, 5, 7, 13, 2389$

From (2) and (3) we have

$$J_2(\omega) = 0 \tag{7}$$

We prove that for $k = 3, 5, 7, 13, 2389$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 7, 13, 2389$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{8}$$

We prove that for $k \neq 3, 5, 7, 13, 2389$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1235)

$$p, jp^{2390} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang
 jiangchunxuan@vip.sohu.com
 Abstract

Using Jiang function we prove that $jp^{2390} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2390} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2390} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2390} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2390} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2390)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 11, 479$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 11, 479$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 11, 479$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 11, 479$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1236)

$$p, jp^{2392} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2392} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2392} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2392} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2392} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2392} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2392)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \tag{6}$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$

Example 1. Let $k = 3, 5, 47, 53, 599, 2393$

From (2) and (3) we have

$$J_2(\omega) = 0 \tag{7}$$

We prove that for $k = 3, 5, 47, 53, 599, 2393$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 47, 53, 599, 2393$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{8}$$

We prove that for $k \neq 3, 5, 47, 53, 599, 2393$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1237)

$$p, jp^{2394} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang
 jiangchunxuan@vip.sohu.com
 Abstract

Using Jiang function we prove that $jp^{2394} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2394} + k - j (j = 1, \dots, k-1) \tag{1}$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \tag{2}$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2394} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \tag{3}$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \tag{4}$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2394} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \tag{5}$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2394} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2394)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 7, 19, 43, 127$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 7, 19, 43, 127$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 7, 19, 43, 127$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 7, 19, 43, 127$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1238)

$$p, jp^{2396} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2396} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2396} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2396} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2396} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2396} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2396)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1239)

$$p, jp^{2398} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2398} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2398} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2398} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2398} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2398} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2398)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 23, 2399$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 23, 2399$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 23, 2399$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 23, 2399$

(1) contain infinitely many prime solutions.

The New Prime Theorem (1240)

$$p, jp^{2400} + k - j (j = 1, 2, \dots, k-1)$$

Chun-Xuan Jiang

jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jp^{2400} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$p, jp^{2400} + k - j (j = 1, \dots, k-1) \quad (1)$$

Contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_{P>2} [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} (jq^{2400} + k - j) \equiv 0 \pmod{p}, q = 1, \dots, p-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions that is for any k there are infinitely many primes P such that each of $jp^{2400} + k - j$ is a prime.

Using Fermat's little theorem from (3) we have $\chi(P) = P-1$. Substituting it into (2) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jp^{2400} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(2400)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N}, \quad (6)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

From (6) we are able to find the smallest solution $\pi_k(N_0, 2) \geq 1$.

Example 1. Let $k = 3, 5, 7, 11, 13, 17, 31, 41, 61, 97, 101, 151, 241, 401, 601$

From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 7, 11, 13, 17, 31, 41, 61, 97, 101, 151, 241, 401, 601$

(1) contain no prime solutions. 1 is not a prime.

Example 2. Let $k \neq 3, 5, 7, 11, 13, 17, 31, 41, 61, 97, 101, 151, 241, 401, 601$

From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 7, 11, 13, 17, 31, 41, 61, 97, 101, 151, 241, 401, 601$

(1) contain infinitely many prime solutions.

Remark. The prime number theory is basically to count the Jiang function $J_{n+1}(\omega)$

$$\sigma(J) = \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} = \prod_P \left(1 - \frac{1 + \chi(P)}{P} \right) \left(1 - \frac{1}{P} \right)^{-k} \quad [1,2],$$

and Jiang prime k -tuple singular series

which can count the number of prime numbers. The prime distribution is not random. But Hardy-Littlewood prime

k -tuple singular series $\sigma(H) = \prod_P \left(1 - \frac{\nu(P)}{P} \right) \left(1 - \frac{1}{P} \right)^{-k}$ is false [3-17], which cannot count the number of prime numbers[3].

References

1. Chun-Xuan Jiang, Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture. Inter. Acad. Press, 2002, MR2004c:11001, (<http://www.i-b-r.org/docs/jiang.pdf>) (<http://www.wbabin.net/math/xuan13.pdf>) (<http://vixra.org/numth/>).
2. Chun-Xuan Jiang, Jiang's function $J_{n+1}(\omega)$ in prime distribution. (<http://www.wbabin.net/math/xuan2.pdf>) (<http://wbabin.net/xuan.htm#chun-xuan>) (<http://vixra.org/numth/>)
3. Chun-Xuan Jiang, The Hardy-Littlewood prime k -tuple conjecture is false. (<http://wbabin.net/xuan.htm#chun-xuan>) (<http://vixra.org/numth/>)
4. G. H. Hardy and J. E. Littlewood, Some problems of "Partitio Numerorum", III: On the expression of a number as a sum of primes. Acta Math., 44(1923)1-70.
5. W. Narkiewicz, The development of prime number theory. From Euclid to Hardy and Littlewood. Springer-Verlag, New York, NY. 2000, 333-353.
6. B. Green and T. Tao, Linear equations in primes. Ann. Math, 171(2010) 1753-1850.
7. D. Goldston, J. Pintz and C. Y. Yildirim, Primes in tuples I. Ann. Math., 170(2009) 819-862.
8. T. Tao. Recent progress in additive prime number theory, preprint. 2009. <http://terrytao.files.wordpress.com/2009/08/prime-number-theory1.pdf>
9. J. Bourgain, A. Gamburd, P. Sarnak, Affine linear sieve, expanders, and sum-product, Invent math, 179 (2010)559-644.
10. K. Soundararajan, The distribution of prime numbers, In: A. Granville and Z. Rudnik (eds), Equidistribution in number theory, an Introduction, 59-83, 2007 Springer.
11. B. Kra, The Green-Tao theorem on arithmetic progressions in the primes: an ergodic point of view, Bull. Amer. Math. Soc., 43(2006)3-23.

12. K. Soundararajan, Small gaps between prime numbers: The work of Goldston-Pintz-Yildirim, Bull. Amer. Math. Soc., 44(2007)1-18.
13. D. A. Goldston, S. W. Graham, J. Pintz and C. Y. Yildirim, Small gaps between products of two primes, Proc. London Math. Soc., 98(2009)741-774.
14. B. Green and T. Tao, The primes contain arbitrarily long arithmetic progressions, Ann. Math., 167(2008) 481-547.
15. D. A. Goldston, J. Pintz and C. Y. Yildirim, Primes in tuples II, Acta Math., 204(2010),1-47.
16. B. Green, Generalising the Hardy-Littlewood method for primes, International congress of mathematicians, Vol, II, 373-399, Eur. Math. Soc., Zurich, 2006.
17. T. Tao, The dichotomy between structure and randomness, arithmetic progressions, and the primes, International congress of mathematicians Vol. I, 581-608, Eur. Math. Soc., Zurich 2006.

Szemerédi’s theorem does not directly to the primes, because it cannot count the number of primes. Cramér’s random model cannot prove any prime problems. The probability of $1/\log N$ of being prime is false. Assuming that the events “ P is prime”, “ $P+2$ is prime” and “ $P+4$ is prime” are independent, we conclude that P , $P+2$, $P+4$ are simultaneously prime with probability about $1/\log^3 N$. There are about $N/\log^3 N$ primes less than N . Letting $N \rightarrow \infty$ we obtain the prime conjecture, which is false. The tool of additive prime number theory is basically the Hardy-Littlewood prime tuples conjecture, but cannot prove and count any prime problems[6].

Mathematicians have tried in vain to discover some order in the sequence of prime numbers but we have every reason to believe that there are some mysteries which the human mind will never penetrate.

Leonhard Euler(1707-1783)

It will be another million years, at least, before we understand the primes.

Paul Erdos(1913-1996)

Jiang’s function $J_{n+1}(\omega)$ in prime distribution

Chun-Xuan Jiang

P. O. Box 3924, Beijing 100854, P. R. China
 jiangchunxuan@vip.sohu.com

Dedicated to the 30-th anniversary of hadronic mechanics

Abstract: We define that prime equations

$$f_1(P_1, \dots, P_n), \dots, f_k(P_1, \dots, P_n) \tag{5}$$

are polynomials (with integer coefficients) irreducible over integers, where P_1, \dots, P_n are all prime. If Jiang’s function $J_{n+1}(\omega) = 0$ then (5) has finite prime solutions. If $J_{n+1}(\omega) \neq 0$ then there are infinitely many primes P_1, \dots, P_n such that f_1, \dots, f_k are primes. We obtain a unite prime formula in prime distribution

$$\begin{aligned} \pi_{k+1}(N, n+1) &= |\{P_1, \dots, P_n \leq N : f_1, \dots, f_k \text{ are } k \text{ primes}\}| \\ &= \prod_{i=1}^k (\deg f_i)^{-1} \times \frac{J_{n+1}(\omega)\omega^k}{n! \phi^{k+n}(\omega) \log^{k+n} N} (1 + o(1)). \end{aligned} \tag{8}$$

Jiang’s function is accurate sieve function. Using Jiang’s function we prove about 600 prime theorems [6]. Jiang’s function provides proofs of the prime theorems which are simple enough to understand and accurate enough to be useful.

Mathematicians have tried in vain to discover some order in the sequence of prime numbers but we have every

reason to believe that there are some mysteries which the human mind will never penetrate.

Leonhard Euler

It will be another million years, at least, before we understand the primes.

Paul Erdős

Suppose that Euler totient function

$$\phi(\omega) = \prod_{2 \leq P} (P-1) = \infty \quad \text{as } \omega \rightarrow \infty, \quad (1)$$

where $\omega = \prod_{2 \leq P} P$ is called primorial.

Suppose that $(\omega, h_i) = 1$, where $i = 1, \dots, \phi(\omega)$. We have prime equations

$$P_1 = \omega n + 1, \dots, P_{\phi(\omega)} = \omega n + h_{\phi(\omega)} \quad (2)$$

where $n = 0, 1, 2, \dots$.

(2) is called infinitely many prime equations (IMPE). Every equation has infinitely many prime solutions. We have

$$\pi_{h_i} = \sum_{\substack{P_i \leq N \\ P_i \equiv h_i \pmod{\omega}}} 1 = \frac{\pi(N)}{\phi(\omega)} (1 + o(1)). \quad (3)$$

where π_{h_i} denotes the number of primes $P_i \leq N$ in $P_i = \omega n + h_i$ $n = 0, 1, 2, \dots$, $\pi(N)$ the number of primes less than or equal to N .

We replace sets of prime numbers by IMPE. (2) is the fundamental tool for proving the prime theorems in prime distribution.

Let $\omega = 30$ and $\phi(30) = 8$. From (2) we have eight prime equations

$$\begin{aligned} P_1 = 30n + 1, \quad P_2 = 30n + 7, \quad P_3 = 30n + 11, \quad P_4 = 30n + 13, \quad P_5 = 30n + 17, \\ P_6 = 30n + 19, \quad P_7 = 30n + 23, \quad P_8 = 30n + 29, \quad n = 0, 1, 2, \dots \end{aligned} \quad (4)$$

Every equation has infinitely many prime solutions.

THEOREM. We define that prime equations

$$f_1(P_1, \dots, P_n), \dots, f_k(P_1, \dots, P_n) \quad (5)$$

are polynomials (with integer coefficients) irreducible over integers, where P_1, \dots, P_n are primes. If Jiang's function $J_{n+1}(\omega) = 0$ then (5) has finite prime solutions. If $J_{n+1}(\omega) \neq 0$ then there exist infinitely many primes P_1, \dots, P_n such that each f_k is a prime.

PROOF. Firstly, we have Jiang's function [1-11]

$$J_{n+1}(\omega) = \prod_{3 \leq P} [(P-1)^n - \chi(P)] \quad (6)$$

where $\chi(P)$ is called sieve constant and denotes the number of solutions for the following congruence

$$\prod_{i=1}^k f_i(q_1, \dots, q_n) \equiv 0 \pmod{P} \quad (7)$$

where $q_1 = 1, \dots, P-1, \dots, q_n = 1, \dots, P-1$.

$J_{n+1}(\omega)$ denotes the number of sets of P_1, \dots, P_n prime equations such that $f_1(P_1, \dots, P_n), \dots, f_k(P_1, \dots, P_n)$ are prime equations. If $J_{n+1}(\omega) = 0$ then (5) has finite prime solutions. If $J_{n+1}(\omega) \neq 0$ using $\chi(P)$ we sift out from (2) prime equations which can not be represented P_1, \dots, P_n , then

residual prime equations of (2) are P_1, \dots, P_n prime equations such that $f_1(P_1, \dots, P_n), \dots, f_k(P_1, \dots, P_n)$ are prime equations. Therefore we prove that there exist infinitely many primes P_1, \dots, P_n such that $f_1(P_1, \dots, P_n), \dots, f_k(P_1, \dots, P_n)$ are primes.

Secondly, we have the best asymptotic formula [2,3,4,6]

$$\begin{aligned} \pi_{k+1}(N, n+1) &= |\{P_1, \dots, P_n \leq N : f_1, \dots, f_k \text{ are } k \text{ primes}\}| \\ &= \prod_{i=1}^k (\deg f_i)^{-1} \times \frac{J_{n+1}(\omega)\omega^k}{n! \phi^{k+n}(\omega)} \frac{N^n}{\log^{k+n} N} (1+o(1)). \end{aligned} \tag{8}$$

(8) is called a unite prime formula in prime distribution. Let $n=1, k=0$, $J_2(\omega) = \phi(\omega)$. From (8) we have prime number theorem

$$\pi_1(N, 2) = |\{P_1 \leq N : P_1 \text{ is prime}\}| = \frac{N}{\log N} (1+o(1)). \tag{9}$$

Number theorists believe that there are infinitely many twin primes, but they do not have rigorous proof of this old conjecture by any method. All the prime theorems are conjectures except the prime number theorem, because they do not prove that prime equations have infinitely many prime solutions. We prove the following conjectures by this theorem.

Example 1. Twin primes $P, P+2$ (300BC).

From (6) and (7) we have Jiang's function

$$J_2(\omega) = \prod_{3 \leq P} (P-2) \neq 0$$

Since $J_2(\omega) \neq 0$ in (2) exist infinitely many P prime equations such that $P+2$ is a prime equation. Therefore we prove that there are infinitely many primes P such that $P+2$ is a prime.

Let $\omega = 30$ and $J_2(30) = 3$. From (4) we have three P prime equations

$$P_3 = 30n + 11, \quad P_5 = 30n + 17, \quad P_8 = 30n + 29$$

From (8) we have the best asymptotic formula

$$\begin{aligned} \pi_2(N, 2) &= |\{P \leq N : P+2 \text{ prime}\}| = \frac{J_2(\omega)\omega}{\phi^2(\omega)} \frac{N}{\log^2 N} (1+o(1)) \\ &= 2 \prod_{3 \leq P} \left(1 - \frac{1}{(P-1)^2}\right) \frac{N}{\log^2 N} (1+o(1)). \end{aligned}$$

In 1996 we proved twin primes conjecture [1]

Remark. $J_2(\omega)$ denotes the number of P prime equations, $\frac{\omega}{\phi^2(\omega)} \frac{N}{\log^2 N} (1+o(1))$ the number of solutions of primes for every P prime equation.

Example 2. Even Goldbach's conjecture $N = P_1 + P_2$. Every even number $N \geq 6$ is the sum of two primes.

From (6) and (7) we have Jiang's function

$$J_2(\omega) = \prod_{3 \leq P} (P-2) \prod_{P|N} \frac{P-1}{P-2} \neq 0$$

Since $J_2(\omega) \neq 0$ as $N \rightarrow \infty$ in (2) exist infinitely many P_1 prime equations such that $N - P_1$ is a prime equation. Therefore we prove that every even number $N \geq 6$ is the sum of two primes.

From (8) we have the best asymptotic formula

$$\begin{aligned}\pi_2(N, 2) &= \left| \{P_1 \leq N, N - P_1 \text{ prime}\} \right| = \frac{J_2(\omega)\omega}{\phi^2(\omega)} \frac{N}{\log^2 N} (1 + o(1)). \\ &= 2 \prod_{3 \leq P} \left(1 - \frac{1}{(P-1)^2} \right) \prod_{P|N} \frac{P-1}{P-2} \frac{N}{\log^2 N} (1 + o(1))\end{aligned}$$

In 1996 we proved even Goldbach's conjecture [1]

Example 3. Prime equations $P, P+2, P+6$.

From (6) and (7) we have Jiang's function

$$J_2(\omega) = \prod_{5 \leq P} (P-3) \neq 0$$

$J_2(\omega)$ denotes the number of P prime equations such that $P+2$ and $P+6$ are prime equations. Since $J_2(\omega) \neq 0$ in (2) exist infinitely many P prime equations such that $P+2$ and $P+6$ are prime equations. Therefore we prove that there are infinitely many primes P such that $P+2$ and $P+6$ are primes.

Let $\omega = 30$, $J_2(30) = 2$. From (4) we have two P prime equations

$$P_3 = 30n + 11, \quad P_5 = 30n + 17$$

From (8) we have the best asymptotic formula

$$\pi_3(N, 2) = \left| \{P \leq N : P+2, P+6 \text{ are primes}\} \right| = \frac{J_2(\omega)\omega^2}{\phi^3(\omega)} \frac{N}{\log^3 N} (1 + o(1)).$$

Example 4. Odd Goldbach's conjecture $N = P_1 + P_2 + P_3$. Every odd number $N \geq 9$ is the sum of three primes.

From (6) and (7) we have Jiang's function

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 3) \prod_{P|N} \left(1 - \frac{1}{P^2 - 3P + 3} \right) \neq 0$$

Since $J_3(\omega) \neq 0$ as $N \rightarrow \infty$ in (2) exist infinitely many pairs of P_1 and P_2 prime equations such that $N - P_1 - P_2$ is a prime equation. Therefore we prove that every odd number $N \geq 9$ is the sum of three primes.

From (8) we have the best asymptotic formula

$$\begin{aligned}\pi_2(N, 3) &= \left| \{P_1, P_2 \leq N : N - P_1 - P_2 \text{ prime}\} \right| = \frac{J_3(\omega)\omega}{2\phi^3(\omega)} \frac{N^2}{\log^3 N} (1 + o(1)) \\ &= \prod_{3 \leq P} \left(1 + \frac{1}{(P-1)^3} \right) \prod_{P|N} \left(1 - \frac{1}{P^3 - 3P + 3} \right) \frac{N^2}{\log^3 N} (1 + o(1))\end{aligned}$$

Example 5. Prime equation $P_3 = P_1 P_2 + 2$.

From (6) and (7) we have Jiang's function

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 2) \neq 0$$

$J_3(\omega)$ denotes the number of pairs of P_1 and P_2 prime equations such that P_3 is a prime equation. Since $J_3(\omega) \neq 0$ in (2) exist infinitely many pairs of P_1 and P_2 prime equations such that P_3 is a prime equation. Therefore we prove that there are infinitely many pairs of primes P_1 and P_2 such that P_3 is a prime.

From (8) we have the best asymptotic formula

$$\pi_2(N, 3) = |\{P_1, P_2 \leq N : P_1 P_2 + 2 \text{ prime}\}| = \frac{J_3(\omega)\omega}{4\phi^3(\omega)} \frac{N^2}{\log^3 N} (1 + o(1)).$$

Note. $\deg(P_1 P_2) = 2$.

Example 6 [12]. Prime equation $P_3 = P_1^3 + 2P_2^3$.

From (6) and (7) we have Jiang's function

$$J_3(\omega) = \prod_{3 \leq P} [(P-1)^2 - \chi(P)] \neq 0$$

where $\chi(P) = 3(P-1)$ if $2^{\frac{P-1}{3}} \equiv 1 \pmod{P}$; $\chi(P) = 0$ if $2^{\frac{P-1}{3}} \not\equiv 1 \pmod{P}$; $\chi(P) = P-1$ otherwise.

Since $J_3(\omega) \neq 0$ in (2) there are infinitely many pairs of P_1 and P_2 prime equations such that P_3 is a prime equation. Therefore we prove that there are infinitely many pairs of primes P_1 and P_2 such that P_3 is a prime.

From (8) we have the best asymptotic formula

$$\pi_2(N, 3) = |\{P_1, P_2 \leq N : P_1^3 + 2P_2^3 \text{ prime}\}| = \frac{J_3(\omega)\omega}{6\phi^3(\omega)} \frac{N^2}{\log^3 N} (1 + o(1)).$$

Example 7 [13]. Prime equation $P_3 = P_1^4 + (P_2 + 1)^2$.

From (6) and (7) we have Jiang's function

$$J_3(\omega) = \prod_{3 \leq P} [(P-1)^2 - \chi(P)] \neq 0$$

where $\chi(P) = 2(P-1)$ if $P \equiv 1 \pmod{4}$; $\chi(P) = 2(P-3)$ if $P \equiv 1 \pmod{8}$; $\chi(P) = 0$ otherwise.

Since $J_3(\omega) \neq 0$ in (2) there are infinitely many pairs of P_1 and P_2 prime equations such that P_3 is a prime equation. Therefore we prove that there are infinitely many pairs of primes P_1 and P_2 such that P_3 is a prime.

From (8) we have the best asymptotic formula

$$\pi_2(N, 3) = |\{P_1, P_2 \leq N : P_3 \text{ prime}\}| = \frac{J_3(\omega)\omega}{8\phi^3(\omega)} \frac{N^2}{\log^3 N} (1 + o(1)).$$

Example 8 [14-20]. Arithmetic progressions consisting only of primes. We define the arithmetic progressions of length k .

$$P_1, P_2 = P_1 + d, P_3 = P_1 + 2d, \dots, P_k = P_1 + (k-1)d, (P_1, d) = 1 \tag{10}$$

From (8) we have the best asymptotic formula

$$\begin{aligned} \pi_2(N, 2) &= |\{P_1 \leq N : P_1, P_1 + d, \dots, P_1 + (k-1)d \text{ are primes}\}| \\ &= \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} \frac{N}{\log^k N} (1 + o(1)). \end{aligned}$$

If $J_2(\omega) = 0$ then (10) has finite prime solutions. If $J_2(\omega) \neq 0$ then there are infinitely many primes P_1 such that P_2, \dots, P_k are primes.

To eliminate d from (10) we have

$$P_3 = 2P_2 - P_1, \quad P_j = (j-1)P_2 - (j-2)P_1, 3 \leq j \leq k$$

From (6) and (7) we have Jiang's function

$$J_3(\omega) = \prod_{3 \leq P < k} (P-1) \prod_{k \leq P} (P-1)(P-k+1) \neq 0$$

Since $J_3(\omega) \neq 0$ in (2) there are infinitely many pairs of P_1 and P_2 prime equations such that P_3, \dots, P_k are prime equations. Therefore we prove that there are infinitely many pairs of primes P_1 and P_2 such that P_3, \dots, P_k are primes.

From (8) we have the best asymptotic formula

$$\begin{aligned} \pi_{k-1}(N, 3) &= \left| \{P_1, P_2 \leq N : (j-1)P_2 - (j-2)P_1 \text{ prime}, 3 \leq j \leq k\} \right| \\ &= \frac{J_3(\omega) \omega^{k-2}}{2\phi^k(\omega)} \frac{N^2}{\log^k N} (1+o(1)) = \frac{1}{2} \prod_{2 \leq P < k} \frac{P^{k-2}}{(P-1)^{k-1}} \prod_{k \leq P} \frac{P^{k-2}(P-k+1)}{(P-1)^{k-1}} \frac{N^2}{\log^k N} (1+o(1)) \end{aligned}$$

Example 9. It is a well-known conjecture that one of $P, P+2, P+2^2$ is always divisible by 3. To generalize above to the k -primes, we prove the following conjectures. Let n be a square-free even number.

1. $P, P+n, P+n^2$,

where $3|(n+1)$.

From (6) and (7) we have $J_2(3) = 0$, hence one of $P, P+n, P+n^2$ is always divisible by 3.

2. $P, P+n, P+n^2, \dots, P+n^4$,

where $5|(n+b), b=2, 3$.

From (6) and (7) we have $J_2(5) = 0$, hence one of $P, P+n, P+n^2, \dots, P+n^4$ is always divisible by 5.

3. $P, P+n, P+n^2, \dots, P+n^6$,

where $7|(n+b), b=2, 4$.

From (6) and (7) we have $J_2(7) = 0$, hence one of $P, P+n, P+n^2, \dots, P+n^6$ is always divisible by 7.

4. $P, P+n, P+n^2, \dots, P+n^{10}$,

where $11|(n+b), b=3, 4, 5, 9$.

From (6) and (7) we have $J_2(11) = 0$, hence one of $P, P+n, P+n^2, \dots, P+n^{10}$ is always divisible by

11.

5. $P, P+n, P+n^2, \dots, P+n^{12}$,

where $13|(n+b), b=2, 6, 7, 11$.

From (6) and (7) we have $J_2(13) = 0$, hence one of $P, P+n, P+n^2, \dots, P+n^{12}$ is always divisible by

13.

6. $P, P+n, P+n^2, \dots, P+n^{16}$,

where $17|(n+b), b=3, 5, 6, 7, 10, 11, 12, 14, 15$.

From (6) and (7) we have $J_2(17) = 0$, hence one of $P, P+n, P+n^2, \dots, P+n^{16}$ is always divisible by

17.

7. $P, P+n, P+n^2, \dots, P+n^{18}$,

where $19|(n+b), b=4, 5, 6, 9, 16, 17$.

From (6) and (7) we have $J_2(19) = 0$, hence one of $P, P+n, P+n^2, \dots, P+n^{18}$ is always divisible by 19.

Example 10. Let n be an even number.

1. $P, P+n^i, i=1, 3, 5, \dots, 2k+1$,

From (6) and (7) we have $J_2(\omega) \neq 0$. Therefore we prove that there exist infinitely many primes P such that $P, P+n^i$ are primes for any k .

2. $P, P+n^i, i=2, 4, 6, \dots, 2k$

From (6) and (7) we have $J_2(\omega) \neq 0$. Therefore we prove that there exist infinitely many primes P such that $P, P+n^i$ are primes for any k .

Example 11. Prime equation $2P_2 = P_1 + P_3$

From (6) and (7) we have Jiang's function

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 2) \neq 0$$

Since $J_3(\omega) \neq 0$ in (2) there are infinitely many pairs of P_1 and P_2 prime equations such that P_3 is prime equations. Therefore we prove that there are infinitely many pairs of primes P_1 and P_2 such that P_3 is a prime.

From (8) we have the best asymptotic formula

$$\pi_2(N, 3) = \left| \{P_1, P_2 \leq N : P_3 \text{ prime}\} \right| = \frac{J_3(\omega)\omega}{2\phi^3(\omega)} \frac{N^2}{\log^3 N} (1 + o(1)).$$

In the same way we can prove $2P_2^2 = P_3 + P_1$ which has the same Jiang's function.

Jiang's function is accurate sieve function. Using it we can prove any irreducible prime equations in prime distribution. There are infinitely many twin primes but we do not have rigorous proof of this old conjecture by any method [20]. As strong as the numerical evidence may be, we still do not even know whether there are infinitely many pairs of twin primes [21]. All the prime theorems are conjectures except the prime number theorem, because they do not prove the simplest twin primes. They conjecture that the prime distribution is randomness [12-26], because they do not understand theory of prime numbers.

Acknowledgements

The Author would like to express his deepest appreciation to M. N. Huxley, R. M. Santilli, L. Schadeck and G. Weiss for their helps and supports.

References

1. Chun-Xuan Jiang, On the Yu-Goldbach prime theorem, Guangxi Sciences (Chinese) 3(1996), 91-2.
2. Chun-Xuan Jiang, Foundations of Santilli's isonumber theory, Part I, Algebras Groups and Geometries, 15(1998), 351-393.
3. ChunXuan Jiang, Foundations of Santilli's isonumber theory, Part II, Algebras Groups and Geometries, 15(1998), 509-544.
4. Chun-Xuan Jiang, Foundations Santilli's isonumber theory, In: Fundamental open problems in sciences at the end of the millennium, T. Gill, K. Liu and E. Trel (Eds) Hadronic Press, USA, (1999), 105-139.
5. Chun-Xuan Jiang, Proof of Schinzel's hypothesis, Algebras Groups and Geometries, 18(2001), 411-420.
6. Chun-Xuan Jiang, Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture, Inter. Acad. Press, 2002, MR2004c: 11001, <http://www.i-b-r.org/jiang.pdf>
7. Chun-Xuan Jiang, Prime theorem in Santilli's isonumber theory, 19(2002), 475-494.
8. Chun-Xuan Jiang, Prime theorem in Santilli's isonumber theory (II), Algebras Groups and Geometries, 20(2003), 149-170.

9. Chun-Xuan Jiang, Disproof's of Riemann's hypothesis, Algebras Groups and Geometries, 22(2005), 123-136. http://www.i-b-r.org/docs/Jiang_Riemann.pdf
10. Chun-Xuan Jiang, Fifteen consecutive integers with exactly k prime factors, Algebras Groups and Geometries, 23(2006), 229-234.
11. Chun-Xuan Jiang, The simplest proofs of both arbitrarily long arithmetic progressions of primes, preprint, 2006.
12. D. R. Heath-Brown, Primes represented by $x^3 + 2y^3$, Acta Math., 186 (2001), 1-84.
13. J. Friedlander and H. Iwaniec, The polynomial $x^2 + y^4$ captures its primes, Ann. Math., 148(1998), 945-1040.
14. E. Szemerédi, On sets of integers containing no k elements in arithmetic progressions, Acta Arith., 27(1975), 299-345.
15. H. Furstenberg, Ergodic behavior of diagonal measures and a theorem of Szemerédi on arithmetic progressions, J. Analyse Math., 31(1997), 204-256.
16. W. T. Gowers, A new proof of Szemerédi's theorem, GAFA, 11(2001), 465-588.
17. B. Kra, The Green-Tao theorem on arithmetic progressions in the primes: An ergodic point of view, Bull. Amer. Math. Soc., 43(2006), 3-23.
18. B. Green and T. Tao, The primes contain arbitrarily long arithmetic progressions, Ann. Math., 167(2008), 481-547.
19. T. Tao, The dichotomy between structure and randomness, arithmetic progressions, and the primes, In: Proceedings of the international congress of mathematicians (Madrid. 2006), Europ. Math. Soc. Vol. 581-608, 2007.
20. B. Green, Long arithmetic progressions of primes, Clay Mathematics Proceedings Vol. 7, 2007,149-159.
21. H. Iwaniec and E. Kowalski, Analytic number theory, Amer. Math. Soc., Providence, RI, 2004
22. R. Crandall and C. Pomerance, Prime numbers a computational perspective, Springer-Verlag, New York, 2005.
23. B. Green, Generalising the Hardy-Littlewood method for primes, In: Proceedings of the international congress of mathematicians (Madrid. 2006), Europ. Math. Soc., Vol. II, 373-399, 2007.
24. K. Soundararajan, Small gaps between prime numbers: The work of Goldston-Pintz-Yildirim, Bull. Amer. Math. Soc., 44(2007), 1-18.
25. Granville, Harald Cramér and distribution of prime numbers, Scand. Actuar. J, 1995(1) (1995), 12-28.
26. J.Friedlander and H.Iwaniec,Opera de Cribro, Colloq. Publication, vol.57, Amer. Math. Soc., (Providence), 2010. This book is outdated and medley.By using sieve methods they cannot prove the Goldbach conjecture and twin primes conjecture.

The Hardy-Littlewood prime k -tuple conjecture is false

Chun-Xuan Jiang
P. O. Box 3924, Beijing 100854, P. R. China
Jiangchunxuan@vip.sohu.com

Abstract: Using Jiang function we prove Jiang prime k -tuple theorem. We prove that the Hardy-Littlewood prime k -tuple conjecture is false. Jiang prime k -tuple theorem can replace the Hardy-Littlewood prime k -tuple conjecture.

(A) Jiang prime k -tuple theorem [1, 2].

We define the prime k -tuple equation

$$p, p + n_i, \tag{1}$$

where $2 | n_i, i = 1, \dots, k - 1$

we have Jiang function [1, 2]

$$J_2(\omega) = \prod_P (P-1 - \chi(P)) \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{i=1}^{k-1} (q + n_i) \equiv 0 \pmod{P}, \quad q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) < P-1$ then $J_2(\omega) \neq 0$. There exist infinitely many primes P such that each of $P+n_i$ is prime. If $\chi(P) = P-1$ then $J_2(\omega) = 0$. There exist finitely many primes P such that each of $P+n_i$ is prime. $J_2(\omega)$ is a subset of Euler function $\phi(\omega)$ [2].

If $J_2(\omega) \neq 0$, then we have the best asymptotic formula of the number of prime P [1, 2]

$$\pi_k(N, 2) = \left| \{P \leq N : P+n_i = \text{prime}\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} \frac{N}{\log^k N} = C(k) \frac{N}{\log^k N} \quad (4)$$

$$\phi(\omega) = \prod_P (P-1)$$

$$C(k) = \prod_P \left(1 - \frac{1 + \chi(P)}{P} \right) \left(1 - \frac{1}{P} \right)^{-k} \quad (5)$$

Example 1. Let $k = 2, P, P+2$, twin primes theorem.

From (3) we have

$$\chi(2) = 0, \quad \chi(P) = 1 \quad \text{if } P > 2, \quad (6)$$

Substituting (6) into (2) we have

$$J_2(\omega) = \prod_{P \geq 3} (P-2) \neq 0 \quad (7)$$

There exist infinitely many primes P such that $P+2$ is prime. Substituting (7) into (4) we have the best asymptotic formula

$$\pi_k(N, 2) = \left| \{P \leq N : P+2 = \text{prime}\} \right| \sim 2 \prod_{P \geq 3} \left(1 - \frac{1}{(P-1)^2} \right) \frac{N}{\log^2 N}. \quad (8)$$

Example 2. Let $k = 3, P, P+2, P+4$.

From (3) we have

$$\chi(2) = 0, \quad \chi(3) = 2 \quad (9)$$

From (2) we have

$$J_2(\omega) = 0 \quad (10)$$

It has only a solution $P = 3, P+2 = 5, P+4 = 7$. One of $P, P+2, P+4$ is always divisible by 3.

Example 3. Let $k = 4, P, P+n$, where $n = 2, 6, 8$.

From (3) we have

$$\chi(2) = 0, \chi(3) = 1, \chi(P) = 3 \quad \text{if } P > 3. \quad (11)$$

Substituting (11) into (2) we have

$$J_2(\omega) = \prod_{P \geq 5} (P-4) \neq 0 \quad (12)$$

There exist infinitely many primes P such that each of $P+n$ is prime. Substituting (12) into (4) we have the best asymptotic formula

$$\pi_4(N, 2) = \left| \{P \leq N : P + n = \text{prime}\} \right| \sim \frac{27}{3} \prod_{P \geq 5} \frac{P^3(P-4)}{(P-1)^4} \frac{N}{\log^4 N} \tag{13}$$

Example 4. Let $k = 5$, P , $P + n$, where $n = 2, 6, 8, 12$.

From (3) we have

$$\chi(2) = 0, \chi(3) = 1, \chi(5) = 3, \chi(P) = 4 \text{ if } P > 5 \tag{14}$$

Substituting (14) into (2) we have

$$J_2(\omega) = \prod_{P \geq 7} (P-5) \neq 0 \tag{15}$$

There exist infinitely many primes P such that each of $P + n$ is prime. Substituting (15) into (4) we have the best asymptotic formula

$$\pi_5(N, 2) = \left| \{P \leq N : P + n = \text{prime}\} \right| \sim \frac{15^4}{2^{11}} \prod_{P \geq 7} \frac{(P-5)P^4}{(P-1)^5} \frac{N}{\log^5 N} \tag{16}$$

Example 5. Let $k = 6$, P , $P + n$, where $n = 2, 6, 8, 12, 14$.

From (3) and (2) we have

$$\chi(2) = 0, \chi(3) = 1, \chi(5) = 4, J_2(5) = 0 \tag{17}$$

It has only a solution $P = 5$, $P + 2 = 7$, $P + 6 = 11$, $P + 8 = 13$, $P + 12 = 17$, $P + 14 = 19$. One of $P + n$ is always divisible by 5.

(B) The Hardy-Littlewood prime k -tuple conjecture[3-14].

This conjecture is generally believed to be true, but has not been proved (Odlyzko et al.1999).

We define the prime k -tuple equation

$$P, P + n_i \tag{18}$$

where $2 | n_i, i = 1, \dots, k-1$.

In 1923 Hardy and Littlewood conjectured the asymptotic formula

$$\pi_k(N, 2) = \left| \{P \leq N : P + n_i = \text{prime}\} \right| \sim H(k) \frac{N}{\log^k N}, \tag{19}$$

where

$$H(k) = \prod_P \left(1 - \frac{\nu(P)}{P} \right) \left(1 - \frac{1}{P} \right)^{-k} \tag{20}$$

$\nu(P)$ is the number of solutions of congruence

$$\prod_{i=1}^{k-1} (q + n_i) \equiv 0 \pmod{P}, \quad q = 1, \dots, P. \tag{21}$$

From (21) we have $\nu(P) < P$ and $H(k) \neq 0$. For any prime k -tuple equation there exist infinitely many primes P such that each of $P + n_i$ is prime, which is false.

Conjecture 1. Let $k = 2, P, P + 2$, twin primes theorem

From (21) we have

$$\nu(P) = 1 \tag{22}$$

Substituting (22) into (20) we have

$$H(2) = \prod_P \frac{P}{P-1} \tag{23}$$

Substituting (23) into (19) we have the asymptotic formula

$$\pi_2(N, 2) = \left| \{P \leq N : P + 2 = \text{prime}\} \right| \sim \prod_P \frac{P}{P-1} \frac{N}{\log^2 N} \quad (24)$$

which is false see example 1.

Conjecture 2. Let $k = 3, P, P + 2, P + 4$.

From (21) we have

$$\nu(2) = 1, \nu(P) = 2 \quad \text{if } P > 2 \quad (25)$$

Substituting (25) into (20) we have

$$H(3) = 4 \prod_{P \geq 3} \frac{P^2(P-2)}{(P-1)^3} \quad (26)$$

Substituting (26) into (19) we have asymptotic formula

$$\pi_3(N, 2) = \left| \{P \leq N : P + 2 = \text{prime}, P + 4 = \text{prime}\} \right| \sim 4 \prod_{P \geq 3} \frac{P^2(P-2)}{(P-1)^3} \frac{N}{\log^3 N} \quad (27)$$

which is false see example 2.

Conjecture 3. Let $k = 4, P, P + n$, where $n = 2, 6, 8$.

From (21) we have

$$\nu(2) = 1, \nu(3) = 2, \nu(P) = 3 \quad \text{if } P > 3 \quad (28)$$

Substituting (28) into (20) we have

$$H(4) = \frac{27}{2} \prod_{P > 3} \frac{P^3(P-3)}{(P-1)^4} \quad (29)$$

Substituting (29) into (19) we have asymptotic formula

$$\pi_4(N, 2) = \left| \{P \leq N : P + n = \text{prime}\} \right| \sim \frac{27}{2} \prod_{P > 3} \frac{P^3(P-3)}{(P-1)^4} \frac{N}{\log^4 N} \quad (30)$$

Which is false see example 3.

Conjecture 4. Let $k = 5, P, P + n$, where $n = 2, 6, 8, 12$

From (21) we have

$$\nu(2) = 1, \nu(3) = 2, \nu(5) = 3, \nu(P) = 4 \quad \text{if } P > 5 \quad (31)$$

Substituting (31) into (20) we have

$$H(5) = \frac{15^4}{4^5} \prod_{P > 5} \frac{P^4(P-4)}{(P-1)^5} \quad (32)$$

Substituting (32) into (19) we have asymptotic formula

$$\pi_5(N, 2) = \left| \{P \leq N : P + n = \text{prime}\} \right| \sim \frac{15^4}{4^5} \prod_{P > 5} \frac{P^4(P-4)}{(P-1)^5} \frac{N}{\log^5 N} \quad (33)$$

Which is false see example 4.

Conjecture 5. Let $k = 6, P, P + n$, where $n = 2, 6, 8, 12, 14$.

From (21) we have

$$\nu(2) = 1, \nu(3) = 2, \nu(5) = 4, \nu(P) = 5 \quad \text{if } P > 5 \quad (34)$$

Substituting (34) into (20) we have

$$H(6) = \frac{15^5}{2^{13}} \prod_{P > 5} \frac{(P-5)P^5}{(P-1)^6} \quad (35)$$

Substituting (35) into (19) we have asymptotic formula

$$\pi_6(N, 2) = \left| \{P \leq N : P + n = \text{prime}\} \right| \sim \frac{15^5}{2^{13}} \prod_{P>5} \frac{(P-5)P^5}{(P-1)^6} \frac{N}{\log^6 N} \quad (36)$$

which is false see example 5.

Conclusion. The Hardy-Littlewood prime k -tuple conjecture is false. The tool of additive prime number theory is basically the Hardy-Littlewood prime tuples conjecture. Jiang prime k -tuple theorem can replace Hardy-Littlewood prime k -tuple Conjecture. There cannot be really modern prime theory without Jiang function.

References

1. Chun-Xuan Jiang, Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture. Inter. Acad. Press, 2002, MR2004c:11001, (<http://www.i-b-r.org/docs/jiang.pdf>) (<http://www.wbabin.net/math/xuan13.pdf>).
2. Chun-Xuan Jiang, Jiang's function $J_{n+1}(\omega)$ in prime distribution. (<http://www.wbabin.net/math/xuan2.pdf>) (<http://vixra.org/pdf/0812.0004v2.pdf>)
3. G. H. Hardy and J. E. Littlewood, Some problems of 'Partition Numerorum', III: On the expression of a number as a sum of primes, Acta Math, 44(1923), 1-70.
4. B. Green and T. Tao, The primes contain arbitrarily long arithmetic progressions, Ann. Math., 167(2008), 481-547.
5. D. A. Goldston, S. W. Graham, J. Pintz and C. Y. Yildirim, Small gaps between products of two primes, Proc. London Math. Soc., (3) 98 (2009) 741-774.
6. D. A. Goldston, S. W. Graham, J. Pintz and C. Y. Yildirim, Small gaps between primes or almost primes, Trans. Amer. Math. Soc., 361(2009) 5285-5330.
7. D. A. Goldston, J. Pintz and C. Y. Yildirim, Primes in tulpes I, Ann. Math., 170(2009) 819-862.
8. P. Ribenboim, The new book of prime number records, 3rd edition, Springer-Verlag, New York, NY, 1995. PP409-411.
9. H. Halberstam and H.-E. Richert, Sieve methods, Academic Press, 1974.
10. Schinzel and W. Sierpinski, Sur certaines hypotheses concernant les nombres premiers, Acta Arith., 4(1958) 185-208.
11. P. T. Bateman and R. A. Horn, A heuristic asymptotic formula concerning the distribution of prime numbers, Math. Comp., 16(1962) 363-367
12. W. Narkiewicz, The development of prime number theory, From Euclid to Hardy and Littlewood, Springer-Verlag, New York, NY, 2000, 333-53.
13. B. Green and T. Tao, Linear equations in primes, To appear, Ann. Math.
14. T. Tao, Recent progress in additive prime number theory, <http://terrytao.files.wordpress.com/2009/08/prime-number-theory1.pdf>

Riemann Paper (1859) Is False

Chun-Xuan. Jiang
P. O. Box3924, Beijing 100854, China
Jiangchunxuan@vip.sohu.com

Abstract: In 1859 Riemann defined the zeta function $\zeta(s)$. From Gamma function he derived the zeta function with Gamma function $\bar{\zeta}(s)$. $\bar{\zeta}(s)$ and $\zeta(s)$ are the two different functions. It is false that $\bar{\zeta}(s)$ replaces $\zeta(s)$. After him later mathematicians put forward Riemann hypothesis (RH) which is false. The Jiang function $J_n(\omega)$ can replace RH.

AMS mathematics subject classification: Primary 11M26.

In 1859 Riemann defined the Riemann zeta function (RZF)[1]

$$\zeta(s) = \prod_p (1 - P^{-s})^{-1} = \sum_{n=1}^{\infty} \frac{1}{n^s}, \quad (1)$$

where $s = \sigma + ti$, $i = \sqrt{-1}$, σ and t are real, P ranges over all primes. RZF is the function of the complex variable s in $\sigma \geq 0, t \neq 0$, which is absolutely convergent.

In 1896 J. Hadamard and de la Vallee Poussin proved independently [2]

$$\zeta(1+ti) \neq 0 \quad (2)$$

In 1998 Jiang proved [3]

$$\zeta(s) \neq 0 \quad (3)$$

where $0 \leq \sigma \leq 1$.

Riemann paper (1859) is false [1] We define Gamma function [1, 2]

$$\Gamma\left(\frac{s}{2}\right) = \int_0^{\infty} e^{-t} t^{\frac{s}{2}-1} dt \quad (4)$$

For $\sigma > 0$. On setting $t = n^2 \pi x$, we observe that

$$\pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) n^{-s} = \int_0^{\infty} x^{\frac{s}{2}-1} e^{-n^2 \pi x} dx \quad (5)$$

Hence, with some care on exchanging summation and integration, for $\sigma > 1$,

$$\begin{aligned} \pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) \bar{\zeta}(s) &= \int_0^{\infty} x^{\frac{s}{2}-1} \left(\sum_{n=1}^{\infty} e^{-n^2 \pi x} \right) dx \\ &= \int_0^{\infty} x^{\frac{s}{2}-1} \left(\frac{\mathcal{G}(x) - 1}{2} \right) dx \end{aligned} \quad (6)$$

where $\bar{\zeta}(s)$ is called Riemann zeta function with gamma function rather than $\zeta(s)$,

$$\mathcal{G}(x) := \sum_{n=-\infty}^{\infty} e^{-n^2 \pi x} \quad (7)$$

is the Jacobi theta function. The functional equation for $\mathcal{G}(x)$ is

$$x^{\frac{1}{2}} \mathcal{G}(x) = \mathcal{G}(x^{-1}), \quad (8)$$

and is valid for $x > 0$.

Finally, using the functional equation of $\mathcal{G}(x)$, we obtain

$$\bar{\zeta}(s) = \frac{\pi^{\frac{s}{2}}}{\Gamma\left(\frac{s}{2}\right)} \left\{ \frac{1}{s(s-1)} + \int_1^{\infty} (x^{\frac{s}{2}-1} + x^{-\frac{s}{2}-\frac{1}{2}}) \cdot \left(\frac{\mathcal{G}(x) - 1}{2} \right) dx \right\}. \quad (9)$$

From (9) we obtain the functional equation

$$\pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) \bar{\zeta}(s) = \pi^{-\frac{1-s}{2}} \Gamma\left(\frac{1-s}{2}\right) \bar{\zeta}(1-s) \quad (10)$$

The function $\bar{\zeta}(s)$ satisfies the following

1. $\bar{\zeta}(s)$ has no zero for $\sigma > 1$;

2. The only pole of $\bar{\zeta}(s)$ is at $s = 1$; it has residue 1 and is simple;
3. $\bar{\zeta}(s)$ has trivial zeros at $s = -2, -4, \dots$ but $\zeta(s)$ has no zeros;
4. The nontrivial zeros lie inside the region $0 \leq \sigma \leq 1$ and are symmetric about both the vertical line $\sigma = 1/2$.

The strip $0 \leq \sigma \leq 1$ is called the critical strip and the vertical line $\sigma = 1/2$ is called the critical line.

Conjecture (The Riemann Hypothesis). All nontrivial zeros of $\bar{\zeta}(s)$ lie on the critical line $\sigma = 1/2$, which is false. [3]

$\bar{\zeta}(s)$ and $\zeta(s)$ are the two different functions. It is false that $\bar{\zeta}(s)$ replaces $\zeta(s)$, Pati proved that is not all complex zeros of $\bar{\zeta}(s)$ lie on the critical line: $\sigma = 1/2$ [4].

Schadeck pointed out that the falsity of RH implies the falsity of RH for finite fields [5, 6]. RH is not directly related to prime theory. Using RH mathematicians prove many prime theorems which is false. In 1994 Jiang discovered Jiang function $J_n(\omega)$ which can replace RH, Riemann zeta function and L-function in view of its proved feature: if $J_n(\omega) \neq 0$ then the prime equation has infinitely many prime solutions; and if $J_n(\omega) = 0$, then the prime equation has finitely many prime solutions. By using $J_n(\omega)$ Jiang proves about 600 prime theorems including the Goldbach's theorem, twin prime theorem and theorem on arithmetic progressions in primes[7,8].

In the same way we have a general formula involving $\bar{\zeta}(s)$

$$\int_0^\infty x^{s-1} \sum_{n=1}^\infty F(nx) dx = \sum_{n=1}^\infty \int_0^\infty x^{s-1} F(nx) dx$$

$$= \sum_{n=1}^\infty \frac{1}{n^s} \int_0^\infty y^{s-1} F(y) dy = \bar{\zeta}(s) \int_0^\infty y^{s-1} F(y) dy \tag{11}$$

where $F(y)$ is arbitrary.

From (11) we obtain many zeta functions $\bar{\zeta}(s)$ which are not directly related to the number theory.

The prime distributions are order rather than random. The arithmetic progressions in primes are not directly related to ergodic theory, harmonic analysis, discrete geometry, and combinatorics. Using the ergodic theory Green and Tao prove that there exist infinitely many arithmetic progressions of length k consisting only of primes which is false [9, 10, 11]. Fermat's last theorem (FLT) is not directly related to elliptic curves. In 1994 using elliptic curves Wiles proved FLT which is false [12]. There are Pythagorean theorem and FLT in the complex hyperbolic functions and complex trigonometric functions. In 1991 without using any number theory Jiang proved FLT which is Fermat's marvelous proof[7, 13].

Primes Represented by $P_1^n + mP_2^n$ [14]

(1) Let $n = 3$ and $m = 2$. We have

$$P_3 = P_1^3 + 2P_2^3$$

We have Jiang function

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 3 - \chi(P)) \neq 0$$

Where $\chi(P) = 2P - 1$ if $2^{\frac{P-1}{3}} \equiv 1 \pmod{P}$; $\chi(P) = -P + 2$ if $2^{\frac{P-1}{3}} \not\equiv 1 \pmod{P}$; $\chi(P) = 1$ otherwise.

Since $J_n(\omega) \neq 0$, there exist infinitely many primes P_1 and P_2 such that P_3 is a prime.

We have the best asymptotic formula

$$\pi_2(N, 3) = \left| \{P_1, P_2 : P_1, P_2 \leq N, P_1^3 + 2P_2^3 = P_3 \text{ prime}\} \right|$$

$$\sim \frac{J_3(\omega)\omega}{6\Phi^3(\omega)} \frac{N^2}{\log^3 N} = \frac{1}{3} \prod_{3 \leq P} \frac{P(P^2 - 3P + 3 - \chi(P))}{(P-1)^3} \frac{N^2}{\log^3 N}$$

$$\omega = \prod_{2 \leq P} P \quad \Phi(\omega) = \prod_{2 \leq P} (P-1)$$

where ω is called primorial,

It is the simplest theorem which is called the Heath-Brown problem [15].

(2) Let $n = P_0$ be an odd prime, $2 \mid m$ and $m \neq \pm b^{P_0}$.
we have

$$P_3 = P_1^{P_0} + mP_2^{P_0}$$

We have

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 3 - \chi(P)) \neq 0$$

where $\chi(P) = -P + 2$ if $P \mid m$; $\chi(P) = (P_0 - 1)P - P_0 + 2$ if $m^{\frac{P-1}{P_0}} \equiv 1 \pmod{P}$;

$\chi(P) = -P + 2$ if $m^{\frac{P-1}{P_0}} \not\equiv 1 \pmod{P}$; $\chi(P) = 1$ otherwise.

Since $J_n(\omega) \neq 0$, there exist infinitely many primes P_1 and P_2 such that P_3 is a prime.
We have

$$\pi_2(N, 3) \sim \frac{J_3(\omega)\omega}{2P_0\Phi^3(\omega)} \frac{N^2}{\log^3 N}$$

The Polynomial $P_1^n + (P_2 + 1)^2$ Captures Its Primes [14]

(1) Let $n = 4$, We have

$$P_3 = P_1^4 + (P_2 + 1)^2$$

We have Jiang function

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 3 - \chi(P)) \neq 0$$

Where $\chi(P) = P$ if $P \equiv 1 \pmod{4}$; $\chi(P) = P - 4$ if $P \equiv 1 \pmod{8}$; $\chi(P) = -P + 2$ otherwise.

Since $J_n(\omega) \neq 0$, there exist infinitely many primes P_1 and P_2 such that P_3 is a prime.

We have the best asymptotic formula

$$\pi_2(N, 3) = \left| \{P_1, P_2 : P_1, P_2 \leq N, P_1^4 + (P_2 + 1)^2 = P_3 \text{ prime}\} \right|$$

$$\sim \frac{J_3(\omega)\omega}{8\Phi^3(\omega)} \frac{N^2}{\log^3 N}$$

It is the simplest theorem which is called Friedlander-Iwaniec problem [16].

(2) Let $n = 4m$, We have

$$P_3 = P_1^{4m} + (P_2 + 1)^2$$

where $m = 1, 2, 3, \dots$

We have Jiang function

$$J_3(\omega) = \prod_{3 \leq P \leq P_1} (P^2 - 3P + 3 - \chi(P)) \neq 0$$

where $\chi(P) = P - 4m$ if $8m|(P-1)$; $\chi(P) = P - 4$ if $8|(P-1)$; $\chi(P) = P$ if $4|(P-1)$; $\chi(P) = -P + 2$ otherwise.

Since $J_3(\omega) \neq 0$, there exist infinitely many primes P_1 and P_2 such that P_3 is a prime. It is a generalization of Euler proof for the existence of infinitely many primes.

We have the best asymptotic formula

$$\pi_2(N, 3) \sim \frac{J_3(\omega)\omega}{8m\Phi^3(\omega)} \frac{N^2}{\log^3 N}$$

(3) Let $n = 2b$. We have

$$P_3 = P_1^{2b} + (P_2 + 1)^2$$

where b is an odd.

We have Jiang function

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 3 - \chi(P)) \neq 0$$

Where $\chi(P) = P - 2b$ if $4b|(P-1)$; $\chi(P) = P - 2$ if $4|(P-1)$; $\chi(P) = -P + 2$ otherwise.

We have the best asymptotic formula

$$\pi_2(N, 3) \sim \frac{J_3(\omega)\omega}{4b\Phi^3(\omega)} \frac{N^2}{\log^3 N}$$

(4) Let $n = P_0$, We have

$$P_3 = P_1^{P_0} + (P_2 + 1)^2$$

where P_0 is an odd. Prime.

we have Jiang function

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 3 - \chi(P)) \neq 0$$

where $\chi(P) = P_0 + 1$ if $P_0|(P-1)$; $\chi(P) = 0$ otherwise.

Since $J_3(\omega) \neq 0$, there exist infinitely many primes P_1 and P_2 such that P_3 is also a prime.

We have the best asymptotic formula

$$\pi_2(N, 3) \sim \frac{J_3(\omega)\omega}{2P_0\Phi^3(\omega)} \frac{N^2}{\log^3 N}$$

The Jiang function $J_n(\omega)$ is closely related to the prime distribution. Using $J_n(\omega)$ we are able to tackle almost all prime problems in the prime distributions.

Acknowledgements

The Author would like to express his deepest appreciation to R. M. Santilli, L. Schadeck and Chen I-wan for their helps and supports.

References

1. B. Riemann, *Über die Anzahl der Primzahlen unter einer gegebenen Grösse*, Monatsber Akad. Berlin, 671-680 (1859).
2. P. Bormein, S. Choi, B. Rooney, *The Riemann hypothesis*, pp28-30, Springer-Verlag, 2007.
3. Chun-Xuan. Jiang, *Disproofs of Riemann hypothesis*, *Algebras Groups and Geometries* 22, 123-136(2005). <http://www.i-b-r.org/docs/JiangRiemann.pdf>
4. Tribikram Pati, *The Riemann hypothesis*, arxiv:

- math/0703367v2, 19 Mar. 2007.
5. Laurent Schadeck, Private communication. Nov. 5. 2007.
 6. Laurent Schadeck, Remarques sur quelques tentatives de demonstration Originales de l'Hypothèse de Riemann et sur la possibilité De les prolonger vers une th éorie des nombres premiers consistante, unpublished, 2007.
 7. Chun-Xuan. Jiang, Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture, Inter. Acad. Press, 2002. MR2004c: 11001, <http://www.i-b-r.org/Jiang.pdf>
 8. Chun-xuan. Jiang, The simplest proofs of both arbitrarily long arithmetic progressions of primes, Preprint (2006).
 9. B. Kra, The Green-Tao theorem on arithmetic progressions in the primes: an ergodic point of view, Bull. Am. Math. Soc. 43, 3-23(2006).
 10. B. Green and T. Tao, The primes contain arbitrarily long arithmetic progressions. To appear, Ann. Math.
 11. T.Tao, The dichotomy between structure and randomness, arithmetic progressions, and the primes. In proceedings of the international congress of mathematicians (Madrid. 2006). Europ. Math. Soc. Vol.1, 581-609, 2007.
 12. Wiles, Modular elliptic curves and Fermat's last theorem, Ann. Math. 141, 443-551 (1995)
 13. Chun-Xuan. Jiang, Fermat's marvelous proofs for Fermat's last theorem, preprint (2007), submit to Ann. Math.
 14. Chun-Xuan. Jiang, Prime theorem in Santilli's isonumber theory (II), Algebras Groups and Geometries 20, 149-170(2003).
 15. D.R.Heath-Brown, Primes represented by $x^3 + 2y^3$. Acta Math. 186, 1-84(2001).
 16. J. Friedlander and H. Iwaniec, The polynomial $x^2 + y^4$ captures its primes. Ann. Math. 148, 945-1040(1998).

4/10/2016