

**New prime K-tuple theorem (4)**

$$P, P + (2j)^2 (j = 1, \dots, k)$$

Jiang, Chunxuan (蒋春暄)

Institute for Basic Research, Palm Harbor, FL34682-1577, USA

And: P. O. Box 3924, Beijing 100854, China (蒋春暄, 北京 3924 信箱, 100854)

[jiangchunxuan@sohu.com](mailto:jiangchunxuan@sohu.com), [cxjiang@mail.bcf.net.cn](mailto:cxjiang@mail.bcf.net.cn), [jcxuan@sina.com](mailto:jcxuan@sina.com), [Jiangchunxuan@vip.sohu.com](mailto:Jiangchunxuan@vip.sohu.com),  
[jcxxxx@163.com](mailto:jcxxxx@163.com)

**Abstract:** Using Jiang function we prove that for every positive integer  $k$  there exist infinitely many primes  $P$  such that each of  $P + (2j)^2$  is prime.

[Chun-Xuan, Jiang. **New prime K-tuple theorem (4)**  $P, P + (2j)^2 (j = 1, \dots, k)$ . *Academ Arena* 2016;8(2s): 6-6]. (ISSN 1553-992X). <http://www.sciencepub.net/academia>. 4. doi:10.7537/marsaj0802s1604.

**Keywords:** new; prime;  $k$ -tuple; theorem; Jiang Chunxuan; mathematics; science; number; function

**Theorem**

$$P, P + (2j)^2 (j = 1, \dots, k) \quad (1)$$

For every positive integer  $k$  there exist infinitely many primes  $P$  such that each of  $P + (2j)^2$  is prime.

**Proof.** We have Jiang function [1]

$$J_2(\omega) = \prod_P (P - 1 - \chi(P)) \quad (2)$$

$$\omega = \prod_P P$$

where

$\chi(P)$  is the number of solutions of congruence

$$\prod_{j=1}^k [q + (2j)^2] \equiv 0 \pmod{P} \quad (3)$$

where  $q = 1, \dots, P-1$

From (3) we have

If  $P < 2k$  then  $\chi(P) = (P-1)/2$ , if  $2k < P$  then  $\chi(P) = k$

From (3) and (2) we have

$$J_2(\omega) = \prod_{P=3}^{P < 2k} \frac{P-1}{2} \prod_{2k < P} (P-1-k) \neq 0 \quad (4)$$

We prove that for every positive integer  $k$  there exist infinitely many primes  $P$  such that each of  $P + (2j)^2$  is prime.

We have the best asymptotic formula [1]

$$\pi_{k+1}(N, 2) = \left| \left\{ P \leq N : P + (2j)^2 = \text{prime} \right\} \right| \sim \frac{J_2(\omega) \omega^k}{\phi^{k+1}(\omega) \log^{k+1} N} \quad (5)$$

The author takes a day to write this paper.

**References**

Chun-Xuan Jiang, Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture. *Inter. Acad. Press*, 2002, MR2004c:110011, (<http://www.i-b-r.org/docs/jiang.pdf>) (<http://www.wbabin.net/math/xuan13.pdf>)