

**New prime K-tuple theorem (5)**  
 $P_1, P_2, jP_1 + (j+1)P_2 (j=1, \dots, k)$

Jiang, Chunxuan (蒋春煊)

Institute for Basic Research, Palm Harbor, FL34682-1577, USA

And: P. O. Box 3924, Beijing 100854, China (蒋春煊, 北京 3924 信箱, 100854)

[jiangchunxuan@sohu.com](mailto:jiangchunxuan@sohu.com), [cxjiang@mail.bcf.net.cn](mailto:cxjiang@mail.bcf.net.cn), [jcxuan@sina.com](mailto:jcxuan@sina.com), [Jiangchunxuan@vip.sohu.com](mailto:Jiangchunxuan@vip.sohu.com),  
[jcxxxx@163.com](mailto:jcxxxx@163.com)

**Abstract:** Using Jiang function we prove that for every positive integer  $k$  there exist infinitely many primes  $P_1$  and  $P_2$  such that each of  $jP_1 + (j+1)P_2$  is prime.

[Chun-Xuan Jiang. **New prime K-tuple theorem (5)**  $P_1, P_2, jP_1 + (j+1)P_2 (j=1, \dots, k)$ . *Academ Arena* 2016;8(2s): 7-8]. (ISSN 1553-992X). <http://www.sciencepub.net/academia>. 5. doi:[10.7537/marsaj0802s1605](https://doi.org/10.7537/marsaj0802s1605).

**Keywords:** new; prime;  $k$ -tuple; theorem; Jiang Chunxuan; mathematics; science; number; function

**Theorem**

$$P_1, P_2, jP_1 + (j+1)P_2 (j=1, \dots, k) \quad (1)$$

For every positive integer  $k$  there exist infinitely many primes  $P_1$  and  $P_2$  such that each of  $jP_1 + (j+1)P_2$  is prime.

**Proof.** We have Jiang function [1, 2]

$$J_3(\omega) = \prod_P [(P-1)^2 - \chi(P)], \quad (2)$$

where  $\omega = \prod_P P$ ,

$\chi(P)$  is the number of solutions of congruence

$$\prod_{j=1}^k [jq_1 + (j+1)q_2] \equiv 0 \pmod{P}, \quad (3)$$

$q_i = 1, \dots, P-1, i = 1, 2$ .

From (3) we have

If  $P \leq k+1$  then  $\chi(P) = (P-1)(P-2)$ , if  $k+1 < P$  then  $\chi(P) = k(P-1)$ .

From (3)and (2) we have

$$J_3(\omega) = \prod_{P=3}^{P \leq k+1} (P-1) \prod_{k+1 < P} [(P-1)^2 - k(P-1)] \neq 0 \quad (4)$$

We prove that for every positive integer  $k$  there exist infinitely many primes  $P_1$  and  $P_2$  such that each of  $jP_1 + (j+1)P_2$  is prime.

We have the best asymptotic formula [1, 2]

$$\pi_{k+1}(N, 3) = |\{P_1, P_2 \leq N : jP_1 + (j+1)P_2 = \text{prime}\}| \sim \frac{J_3(\omega) \omega^k}{\phi^{k+2}(\omega)} \frac{N^2}{\log^{k+2}}, \quad (5)$$

where  $\phi(\omega) = \prod_P (P-1)$ .

**References**

- 
1. Chun-Xuan Jiang, Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture. Inter. Acad. Press,2002,MR2004c:110011, (<http://www.i-b-r.org/docs/jiang.pdf>) (<http://www.wbabin.net/math/xuan13.pdf>).
  2. Chun-Xuan Jiang, Jiang's function  $J_{n+1}(\omega)$  in prime distribution. (<http://www.wbabin.net/math/xuan2.pdf>) (<http://vixra.org/pdf/0812.0004v2.pdf>) The author takes a day to write this paper. <http://wbabin.net/xuan.htm#chun-xuan>.

4/27/2016