

Estimation of Natural Radionuclides in Grasses, Soils, and Cattle-dungs from a Cattle Rearing-Field at Mangoro-Agege, Lagos State, Nigeria.

Ilori, A.O. ^{1*} and Alausa, S.K. ²

¹ Department of Physical Sciences, Ondo State University of Science and Technology, Okitipupa, Nigeria.

iloriabiola99@gmail.com

² Department of Physics, Olabisi Onabanjo University, Ago-Iwoye, Ogun State, Nigeria. skalaus@yahoo.com

*Corresponding Author: iloriabiola99@gmail.com

Abstract: Estimating the levels of radiation in the environment is crucial in implementing appropriate controls for the sake of radiological protection. It was earlier reported that the natural radioactivity levels of soil in some locations in Agege were significantly higher than the world average values and as such grasses grown in such soils may be susceptible to high radioactivity levels through the plant-root uptake of radionuclides from such soils. Livestock that ingests such grasses may be susceptible to high radionuclides retention which may result in enhanced internal radiation exposure to human. Therefore, the aim of the study was to determine the radioactivity levels of ⁴⁰K, ²³⁸U and ²³²Th in the grass, soils, and dung and its radiological impacts. A well calibrated NaI (TI) and well-shielded detector couple to computer resident quantum multichannel analyzer was employed to determine the activity concentrations from the study area. The results obtained from the study area showed the radioactivity level in the soil were lower than the values earlier reported for a corresponding cattle-rearing field at Alabata, Abeokuta and also lower than the world average values. The activity concentrations of the radionuclides were similar to the reported values for Tehran-Iran but higher than some values obtained in some other literature. The transfer factor indicated that only a fractional amount of the radionuclides was transferred from the soil to the grass. The study indicated a possible low radiological effect on human feeding on cattle beef in the study area when compared to that earlier reported in other literature.

[Ilori, A.O. and Alausa, S.K. **Estimation of Natural Radionuclides in Grasses, Soils, and Cattle-dungs from a Cattle Rearing-Field at Mangoro-Agege, Lagos State, Nigeria.** *Cancer Biology* 2019;9(4):62-72]. ISSN: 2150-1041 (print); ISSN: 2150-105X (online). <http://www.cancerbio.net>. 9. doi:10.7537/marscbj090419.09.

Keywords: Radionuclides, transfer factor, grass, soil, cattle-dung, rearing field, Nigeria.

1.0 Introduction

Estimating the levels of radiation in the environment is crucial in implementing appropriate controls for the sake of radiological protection since the constituents that make up the environment contains various amount of radionuclides and their decay products. The major ways by which humans become exposed to these radiations are radiation from sources outside the body (external exposure), radionuclides that are ingested through consumption of food and water or as inhaled radioactive gases (internal exposure) ⁽¹⁾. The two main sources of exposure are cosmic rays that are released from outer space and from the surface of the sun and terrestrial radionuclides that occur in the earth crust in building materials and in the dust, water and food and the human body at large ⁽²⁾. Also, traces of radionuclides are found in water, air, soil, and human bodies. We inhale and ingest radionuclides every day of our lives and radioactive material has been ubiquitous on earth since its creation. The presence of natural radioactivity in soil results in internal and external exposure to humans ⁽³⁾. The most commonly encountered radionuclides that irradiate the human body through external exposure (primarily by gamma radiation) are

²³⁸U and ²³²Th, their radioactive decay products and ⁴⁰K.

Cosmic rays also offer natural external exposure of 13% with the earth's gamma radiation imparting a natural external radiation exposure of 15% ⁽⁴⁾. A small percentage dose may seem insignificant, but every effort must be made to monitor the environment for radiation in order to control the radiation dose released to a man from the environment. Cosmic radiation comes from both the primary energetic protons and alpha particles of extra-terrestrial origin that strike the earth's atmosphere and the secondary particles or cosmogenic radionuclides which are continuously generated by bombardment of stable nuclides in the atmosphere from these cosmic rays ⁽⁵⁾. At higher elevations, the amount of atmosphere shielding cosmic ray decreases and thus, the dose increases. Terrestrial radiation is natural sources of radiation in the ground, rocks, building materials, and drinking water supplies, they are often called "primordial". Some of the contributors to terrestrial sources are natural radium, Uranium, and Thorium. Even though terrestrial radiation (primordial radionuclides) are ubiquitous, their concentrations vary substantially with location. The distribution of the primordial radionuclides and

their progeny in natural ecosystems is influenced by many factors, including chemical properties of the nuclides, physical factors of the ecosystem, and physiological and ecological attributes of the biota. The naturally occurring radioactive materials are known as the largest sources of exposure to human (2).

The nuclei of elements exhibiting radioactivity are unstable and are found to be undergoing continuous disintegration, the decay process carries on until a daughter nucleus reaches stability. Consequently, the energy of the transformation can be released by the emission of nuclear particles and/or in terms of electromagnetic radiation. The unstable nuclei attempt to become more stable by emitting alpha, beta or gamma radiation. When an unstable parent nucleus emits an alpha or beta particle, it is left with a different number of protons and neutrons. This new nucleus is called a daughter nucleus.

The strength or intensity of the radioactivity is called the activity and is defined as the rate of nuclei number decaying. The probability per unit of time for the decay of given nucleus is a constant which can be commonly named as the disintegration or decay constant (λ) (6). The rate of radioactive decay which is related to the activity can be expressed by the fundamental law of radioactive decay by:

$$A = -dN/dt = \lambda N \quad (1)$$

Where A is the activity of an isotopically pure source, which is equal to the number, dN , of radioactive nuclei integrating in a given time, dt , and is proportional to the number, N , of radioactive nuclei at a time, t . λ is the decay constant and the negative sign indicates that the number of radioactive nuclei decreases when the time increases. The solution to equation 1 leads to the exponential laws of radioactive decay which may be written as (7):

$$N_t = N_0 e^{-\lambda t} \quad (2)$$

Where N_0 is the original number of nuclei present at time $t = 0$. The number of nuclei decreases exponentially with time. Usually, the rate of radioactive decay can be characterized in terms of a specific time frame known as the half-life (6). The half-life ($T_{1/2}$) is the time required for one-half of a certain number of active nuclei to disintegrate. The decay constant correlates with half-life by:

$$\lambda = \frac{0.693}{T_{1/2}} \quad (3)$$

The decay constant is also considered in terms of the mean life, τ which is described as the mathematical average time a radioactive nucleus is likely to survive before it decays. The relationship between the decay constant and the mean lifetime is:

$$\tau = \frac{1}{\lambda} \quad (4)$$

Thus, the mean lifetime can be expressed in terms of the half-life by using equation 3 and 4 as:

$$\tau = \frac{T_{1/2}}{0.693} \quad (5)$$

The half-life and mean lifetime for radionuclides can range from fractions of seconds to billions of years (8).

Naturally occurring radionuclides of terrestrial origin also referred to as primordial radionuclides, they are present in various degrees in the various components of the environment. These radionuclides with half-lives long enough comparable to the number of years the earth has been in existence as well as their progenies exist in substantial amount in rocks and soil. They therefore contribute significantly to population exposure. Radionuclides such as ^{40}K , ^{238}U and ^{232}Th which are present in trace amounts in soil represent the major source of external exposure due to gamma radiation (9).

Radionuclides in solution can then be incorporated through root hair and then to the root of grasses for onward transfer to the leaf system of the grasses that cattle feed upon. In most cases, this is facilitated by their chemical similarity with other element that the grasses usually depend on for growth. The ingestion of radionuclide through the grass by cattle is dependent on the concentration of radionuclides in the grasses consumed (10). It is therefore very important to assess the radiological safety of grasses consumed by cattle. This is because the ingestion of meats loaded with radionuclides has the potential of exposing human beings to high level of radiation dose. Furthermore, radionuclides with relatively long half-lives are considered human health risk as they can get into the human system through the food chain and thereby increase the radiation burden for many years (11).

Cattle products are important contributors to total food production. The addition of milk and meat provides protein, calcium, vitamins, and other nutrients that could be lacking in diets that are exclusively made up of staples such as cereals. The poor, in particular, use organic fertilizer from cattle operations, especially when rising petroleum prices make chemical fertilizers unaffordable. Cattle also store value and provide insurance for people who have no other financial markets available to them. Cattle trade provides the largest livestock market in the Nigeria. Millions of Nigerians made their livelihood from the beef enterprises as producer, marketers, and transporters. Others, as processors of beef products, feed millers, veterinary services, and, in agricultural pieces of machinery. It also generates a lot of revenue to the government through various forms of taxations. Nigeria plays a vital role in the livestock economy of Africa.

An important goal of the United Nations (UN) relating to sustainable food security is to assist

members' states in ensuring that people have access to food that is sufficient, nutritionally adequate and above all considered safe for human consumption⁽¹²⁾. The presence of radionuclide in soil above a certain threshold leads to contamination of grass since grasses derived their nutrients for growth from the topsoil on which they are grown.

The natural radioactivity levels in soils of some locations in Abeokuta were reported to be significantly higher than the world average⁽¹³⁾. Hence, the radioactivity levels in grasses and vegetation grown in such soils may be enhanced through the plant-root uptake of radionuclides and such, animals grazing on the grasses may have a high retention of natural radionuclides in their body. Also, the radioactivity levels in the soils from a cattle rearing-field at Alabata, Abeokuta was reported to be higher than the values earlier reported for Abeokuta and the world average values⁽¹⁴⁾. The activity concentration of radionuclides in the soil of Agege was reported to be significantly high⁽¹⁵⁾. Therefore, this study is significant in accessing the activity concentration of natural radionuclides in soil, grass and dung of cattle in a cattle-rearing field at Agege, Lagos state and evaluate the amount of radionuclides retained in the

cattle in order to access the likely health cancer risk due to ingestion of beef from the cattle in the area.

2.0 Materials and Methods

2.1 Geographical description of the study area

Mangoro-Agege, the study area, is a boundary city between Lagos and Ogun state sharing same geological setting as Abeokuta, which unconformably overlies the rocks of the Basement Complex to the Quaternary Deltaic sands⁽¹⁶⁾. Agege is directly underlain by the Benin Formation. The Benin Formation consists largely of sands/sandstones with lenses of shales and clays. The geologic succession in Lagos spans through the Cretaceous Abeokuta Formation.

Agege is a suburb and Local Government Area in Ikeja Division of Lagos State, Southwestern, Nigeria. It has geographical coordinates of 6°37'17" North and 3°19'33" East with a mean annual temperature of 27°C, it has a density (inh.per km²) of 41,071, land mass area (in km²) of 17km², annual rainfall of 1540mm and the population in the area is 459,939⁽¹⁷⁾. From the western part of Lagos state Nigeria, the cattle rearing field is at the boundary of Agege to Abeokuta expressway leading to Ikeja local government area.




Figure 1: Geological Map of Lagos State showing Agege (the study area)¹⁶.

2.2 Samples collection

A total of 20 soil samples, 20 grass samples, and 20 cattle dung samples were collected for the present study, these points of the collection were marked out using a Global Positioning System (GPS). Grass samples were uprooted from the grazing ground and the whole grass was collected, Soil samples were collected from depths 5cm at the same location where

grass samples were collected. The dung samples were collected at various locations within the grazing field where the cattle directly defecated. At the collection point, all samples were wrapped in separate black plastic bags and were well labeled with a paper masking tape. The samples were then transported to the laboratory for preparation. Below is the GPS representation of the location of samples.

Table 1: Some samples location mapped out by GPS

Samples	GPS (Location)	Samples	GPS (Location)
1	N 06 ⁰ 36.801 E 003 ⁰ 19.681	11	N 06 ⁰ 37.325 E 003 ⁰ 18.820
2	N 06 ⁰ 36.815 E 003 ⁰ 19.662	12	N 06 ⁰ 36.425 E 003 ⁰ 19.620
3	N 06 ⁰ 36.808 E 003 ⁰ 19.673	13	N 06 ⁰ 37.035 E 003 ⁰ 19.305
4	N 06 ⁰ 36.825 E 003 ⁰ 19.650	14	N 06 ⁰ 37.423 E 003 ⁰ 19.321
5	N 06 ⁰ 36.925 E 003 ⁰ 19.647	15	N 06 ⁰ 36.812 E 003 ⁰ 19.723
6	N 06 ⁰ 36.961 E 003 ⁰ 19.663	16	N 06 ⁰ 36.308 E 003 ⁰ 19.628
7	N 06 ⁰ 37.000 E 003 ⁰ 19.590	17	N 06 ⁰ 37.435 E 003 ⁰ 18.039
8	N 06 ⁰ 37.125 E 003 ⁰ 19.542	18	N 06 ⁰ 37.306 E 003 ⁰ 18.596
9	N 06 ⁰ 37.145 E 003 ⁰ 19.550	19	N 06 ⁰ 37.125 E 003 ⁰ 18.345
10	N 06 ⁰ 37.240 E 003 ⁰ 19.613	20	N 06 ⁰ 36.806 E 003 ⁰ 19.675

2.3 Samples preparation

Soils were well mixed after removing extraneous materials such as roots, pieces of stones and gravels. Samples were then weighted and dried into an electric oven at 110°C for 4 days until a constant dry weight was obtained. After crushing and mixing thoroughly, soil samples were shaken in a sieve shaker and were scaled in 200g each⁽¹⁸⁾.

Grass samples were cleared by fresh water for removing the dust and surface contaminations. All the samples were then dried under the sun and humidity condition for 2 days, they were then weighed in 200g each⁽¹⁹⁾. The samples were charred and then finely ground into fine powder, the ash samples were cooled at room temperature. The cooled ash samples were weighed⁽²⁰⁾.

Dung samples were dried for 4 days in an electric oven of 110°C to obtain a constant dry weight. The samples were then stored for about 30 days, to reach secular equilibrium between ²²⁶Ra (daughter of ²³⁸U and ²³²Th with their daughter nuclei, in a 250cm³ container with a cap and wrapped with thick vinyl tapes. This was done in order to allow radon and its short-lived progenies to reach secular radioactive equilibrium. Furthermore, all the containers where samples were kept were preserved airtight by plastic packets to ensure that ²²²Rn and ²²⁰Rn are confined within the volume⁽²¹⁾.

2.4 Detection of radiation by NaI (TI) crystal

When incoming radiation is incident on the crystal, it gives up its energy E completely to the

scintillator resulting in the production of N number of photons given by:

$$N = \frac{E_q}{\omega_0} \quad (6)$$

Where E_q is the luminescence quantum efficiency, which is the probability of a photoelectric interaction of the incident photon and ω_0 is the average energy of a single photon which is about 3.0eV for NaI (TI). These N photons impinge on the photocathode of the photomultiplier tube and are converted into photoelectrons, which are directed to the incident on the first dynode of the photomultiplier tube. The total number N_c of photoelectrons at the first dynode is given by:

$$N_c = \frac{E_q}{\omega_0} mc_{p,e}g_cG \quad (7)$$

Where G is the light collection efficiency of the photocathode which gives the function of photons that impinge on the photocathode which are converted into photoelectrons; $c_{p,e}$ is the photo-quantum efficiency of the window-cathode system, m is a factor between 0 and 1 depending on the degree of spectral matching between the scintillation spectrum and the spectral responses of the photocathode; g_c is the efficiency with which the first dynode collects the number of electrons arriving. These efficiencies are affected by a number of factors; G is a factor determined by self-absorption, reflection losses, light trapping, optical flaws and the optical geometry of the photocathode. In NaI (TI), G is usually made nearly unity by coating the director with a reflector like MgO thereby making self-absorption very small. The term $mc_{p,e}g_c$ depends in a complex manner, on the wavelength and the point

of the incident of the photons on the photocathode. Factor $c_{p,e}$ depends on the cathode material and its thickness while g_c depends on the structure of the dynode and its potential. The total number of electrons Q at the last dynode collected at the anode is given as:

$$Q = MN_i \quad (8)$$

Where M is the overall gain resulting from the k successive multiplications of N_c electrons at each dynode given by:

$$M = \prod_i^k m_i \quad (9)$$

Where m_i is the multiplication at the i^{th} dynode.

It is roughly proportional to the voltage between the dynodes. It can be observed from equations 6 and 7 that Q is a linear function of the energy E of the initial incident photon. Apart from the number of electrons given by equation 8, there are a number of electrons produced due to thermionic emission in the photomultiplier tube. The number of electrons with thermal energy greater than the work function of photocathode which is emitted as thermionic electrons is a function of temperature as given in the equation:

$$n_T = ATe^{\frac{Qe}{kT}} \quad (10)$$

Where T is the absolute temperature, e is the electronic charge, k is the Boltzmann constant, A and Q are characteristics of the cathode material.

2.5 Activity determination

A well calibrated NaI (TI) and a well-shielded detector coupled to a computer resident quantum MCA2100R Multichannel analyzer for 36,000s were employed. An empty container under identical geometry was also counted for the same time. The 1460KeV gamma-radiation of ^{40}K was used to determine the concentration of ^{40}K in the sample. The gamma transition energy of 1764.5KeV ^{214}Bi was used to determine the concentration of ^{238}U while the gamma transition energy of 2614KeV ^{208}Tl was used to determine the concentration of ^{232}Th while ^{137}Cs was detected by its 661.6KeV gamma transition.

$$C_s = \frac{C_\alpha}{P_\gamma \left(\frac{M_s}{V_s}\right) \epsilon_\gamma t_c} (BqKg^{-1}) \quad (11)$$

Where C_s is the sample concentration, C_α is the net peak energy, ϵ_γ is the efficiency of the detector for a γ -energy of interest, M_s/V_s is the sample mass per volume of soil, t_c is the total counting time and P_γ is the abundance of the γ -line in a radionuclide.

The efficiency calibration of the detector was done using a reference standard mixed source traceable to Analytical Quality Control Service (AQCS, USA), which has certified activities of the selected radionuclide and has a geometrical configuration identical to sample container. The standard sources contained ten known radionuclides. The energy calibration was also performed by using the peaks of the radionuclide present in the standard sources. The channel number is proportional to

energy; the channel scale was then converted to an energy scale. This produces an energy calibration curve, i.e. energy versus channel.

2.6 Transfer factor

The soil-to-grass transfer factor (TF) measured the transfer of radionuclides from the soil to grass. In soil, each radioactive element follows complex dynamics in which a part of its concentration is transported into the soil solution, while another part gradually becomes strongly bound to the particles of the soil. The portion of these radionuclides, which is in the soil solution, can be incorporated via the root into the grass⁽²¹⁾.

From observed activity concentrations of the radionuclides in the grass and in the corresponding soil, the transfer factor (TF) values were calculated according to the equation:

$$TF = \frac{A_g}{A_s} \quad (12)$$

Where: A_g = Activity of radionuclides in the grass (BqKg⁻¹ dry weight)

A_s = Activity of radionuclides in soil (BqKg⁻¹ dry weight)

The dry weight was preferred because the amount of radioactivity per kilogram dry weight is much less variable than the amount per unit of fresh weight. The soil-to-grass TF can be used as an index for the accumulation of trace elements by plants or the transfer of elements from the soil to the plant.

3.0 Results

3.1 Activity measurement in the soil samples

In the study, the radioactivity level in the soil ranged from 301.18 to 440.48 Bqkg⁻¹ with a mean value of 403.07±33.85 Bqkg⁻¹ for ^{40}K ; 10.19 to 13.05 Bqkg⁻¹ with a mean value of 11.47±0.75 Bqkg⁻¹ for ^{238}U and 9.12 to 11.97 Bqkg⁻¹ with a mean value of 10.44±0.75 Bqkg⁻¹ for ^{232}Th . The activity concentrations of ^{40}K , ^{238}U and ^{232}Th radionuclides in the soil samples were higher than the corresponding values in grass samples. This affirmed the general assertion that only a fractional part of the radionuclides in the soil is transferable to the plant. The activity concentrations of ^{40}K , ^{238}U and ^{232}Th were lower than the values of 411±341 Bqkg⁻¹ for ^{40}K ; 184±205 Bqkg⁻¹ for ^{226}Ra and 65±29 Bqkg⁻¹ for ^{238}U reported for Abeokuta⁽²²⁾. The values were also lower than the world average values of 410.0 Bqkg⁻¹ for ^{40}K ; 35.0 Bqkg⁻¹ for ^{238}U and 28.0 Bq kg⁻¹ for ^{232}Th ⁽²⁾. The low activity concentrations of radionuclides in the soil samples from the study may be attributed to the underlain Benin formation, which consists of largely sands/sandstones with lenses of shales and clays overlain by the Basement complex⁽¹⁶⁾. The ranges of activity concentration of the radionuclides are similar to the values of 12 to 31 Bqkg⁻¹ for ^{238}U ; 14 to 36

Bqkg⁻¹ for ²³²Th and 267 to 867 Bqkg⁻¹ for ⁴⁰K for Tehran-Iran⁽⁹⁾.

Table 2: Activity concentrations ²³²Th in soil samples from the study area

Samples	⁴⁰ K (Bqkg ⁻¹)	²³⁸ U (Bqkg ⁻¹)	²³² Th (Bqkg ⁻¹)
1	429.11±24.43	13.05±0.84	11.97±0.57
2	421.43±38.00	12.13±0.19	11.03±0.16
3	413.78±26.88	11.01±0.50	10.09±0.60
4	301.18±15.75	11.15±0.99	12.08±2.25
5	412.16±10.01	10.19±1.01	9.12±0.08
6	418.22±13.43	10.93±0.67	10.01±0.21
7	420.02±10.16	11.61±0.40	10.13±1.13
8	422.18±21.57	12.29±2.30	10.25±1.70
9	412.31±41.07	12.03±1.95	10.13±0.30
10	420.73±24.08	11.87±0.80	10.58±1.65
11	404.41±17.05	11.68±2.34	9.93±0.54
12	412.24±25.54	11.11±1.98	9.63±0.69
13	420.07±13.89	10.54±1.67	9.33±0.86
14	361.24±22.34	10.23±0.80	10.20±2.31
15	356.67±12.79	10.67±1.50	10.60±0.17
16	352.10±16.72	11.11±0.45	11.00±1.90
17	420.34±17.12	11.74±0.60	10.36±0.89
18	401.68±21.38	11.96±1.40	10.64±0.43
19	421.08±22.56	12.01±0.99	10.81±1.41
20	440.48±20.05	12.06±0.32	10.96±0.71
Mean ±σ	403.07±33.85	11.47±0.75	10.44±0.75

3.2 Activity measurement in the grass samples

The activity concentrations of ⁴⁰K, ²³⁸U and ²³²Th in the grass samples ranges from 202.41 to 99.57 Bqkg⁻¹ with a mean value of 115.46±21.68 Bqkg⁻¹ for ⁴⁰K; 11.21 to 8.44 Bqkg⁻¹ with a mean value of 10.06±0.75 Bqkg⁻¹ for ²³⁸U and 10.50 to 3.00 Bqkg⁻¹ with a mean value of 8.31±2.76 Bqkg⁻¹ for ²³²Th. ⁴⁰K activity concentration exhibited the highest values among other radionuclides as shown in Table 3. Grasses and other plants derive their nutrients from the topsoil on which they are grown and the presence of radionuclides in such soil above certain threshold encourages the soil-to-plant uptake of such radionuclides⁽¹²⁾. The average radioactivity concentrations in elephant grasses as 25.7±5.5 Bqkg⁻¹ for ²³⁸U and 33.4±3.9 Bqkg⁻¹ for ²³²Th reported for Ibadan⁽²²⁾. The mean radioactivity levels of 291.8±58.3 Bqkg⁻¹ for ⁴⁰K; 11.7±0.8 Bqkg⁻¹ for ²³⁸U and 9.3±2.0 Bqkg⁻¹ for ²³²Th reported in grasses⁽¹⁴⁾ were higher than the corresponding values in the present study. In the present study, the values of ⁴⁰K were lower while the values of ²³²Th were higher than the values reported for vegetables in Kuca, Jos⁽²³⁾. The range of ⁴⁰K activity concentrations in the study was lower when compared to the range of 201 to 684

Bqkg⁻¹ but the ²³²Th activity concentrations in the study were higher than the values in vegetables consumed in Jordan⁽²⁴⁾.

Table 3: Activity concentrations in grass samples from the study area

Samples	⁴⁰ K (Bqkg ⁻¹)	²³⁸ U (Bqkg ⁻¹)	²³² Th (Bqkg ⁻¹)
1	109.48±25.80	9.83±0.85	9.76±1.10
2	112.12±30.10	10.01±1.01	10.01±1.24
3	114.76±23.45	10.01±0.46	10.30±1.45
4	105.12±11.03	10.97±1.27	3.00±1.12
5	125.03±25.46	9.02±0.20	3.00±1.45
6	121.10±18.94	8.73±0.69	3.00±2.89
7	117.17±19.28	8.44±1.21	3.00±3.50
8	202.41±15.89	10.36±0.92	10.18±4.02
9	99.57±21.41	10.11±1.75	9.31±3.05
10	100.02±16.22	10.24±0.51	9.14±2.25
11	100.47±21.27	10.37±0.87	8.97±3.16
12	102.57±20.57	10.11±2.61	10.11±7.67
13	107.03±13.56	11.11±0.82	9.02±5.53
14	108.62±12.89	11.16±0.69	9.08±11.50
15	110.21±18.25	11.21±3.74	9.10±2.25
16	113.11±23.69	9.85±1.23	9.03±3.81
17	110.56±23.43	9.49±0.98	10.07±1.10
18	120.13±20.45	9.78±1.20	9.54±0.98
19	116.61±27.78	10.04±1.61	10.02±2.56
20	113.09±16.08	10.30±0.58	10.50±1.14
Mean ±σ	115.46±21.68	10.06±0.75	8.31±2.76

3.3 Activity measurement in the cattle dung samples

The activity concentration of ⁴⁰K, ²³⁸U and ²³²Th in the dung samples range from 115.21 to 225.44 Bqkg⁻¹ with a mean value of 184.90±40.92 Bqkg⁻¹ for ⁴⁰K; 10.13 to 13.17 Bqkg⁻¹ with a mean value of 11.50±0.78 Bqkg⁻¹ for ²³⁸U and 8.12 to 11.28 Bqkg⁻¹ with a mean value of 10.20±0.71 Bqkg⁻¹ for ²³²Th. The activity concentrations in cattle dung samples were higher than the corresponding values in the grass from the study area. This may be attributed to the ingestion of other sources of radionuclides such as the drinking water available to the cattle grazing in the study area. The activity concentrations of the radionuclides in water samples were reported to be significantly high in Agege⁽²⁵⁾. The ⁴⁰K value of 218.6±66.0 Bqkg⁻¹ obtained in Abeokuta⁽¹⁴⁾ was higher than that of the present study. However, the ²³⁸U and ²³²Th activity concentration values of 10.5±1.1 and 8.3±1.9 Bqkg⁻¹ obtained in Abeokuta⁽¹⁴⁾ were lower when compared to that obtained in the present study. The mean values for the differences between the dungs and grasses are 69.44±42.48 Bqkg⁻¹ for ⁴⁰K; 1.44±0.68 Bqkg⁻¹ for ²³⁸U and 1.89±2.61 Bqkg⁻¹ for ²³²Th.

Table 4: Activity concentrations in dung samples from the study area

Samples	^{40}K (Bqkg $^{-1}$)	^{238}U (Bqkg $^{-1}$)	^{232}Th (Bqkg $^{-1}$)
1	225.44±30.10	11.73±1.01	9.80±0.24
2	212.32±32.89	11.35±1.17	10.33±0.38
3	199.20±41.03	10.97±0.57	10.36±0.60
4	221.02±40.05	13.17±0.44	8.12±1.41
5	212.10±25.86	10.13±0.20	10.01±0.81
6	210.14±33.82	10.53±0.68	10.30±1.34
7	150.14±21.93	10.93±0.28	10.59±1.11
8	208.14±26.80	11.76±0.52	10.18±1.02
9	127.43±23.96	11.97±0.75	11.00±0.40
10	121.32±19.28	12.20±0.81	11.14±0.83
11	115.21±33.16	12.07±1.61	11.28±0.61
12	211.23±29.17	10.32±1.20	10.17±1.31
13	219.53±36.94	11.29±1.29	9.76±0.74
14	216.67±27.98	11.64±0.92	9.59±0.89
15	213.81±20.83	11.99±1.89	9.42±0.96
16	215.58±27.85	11.85±1.01	9.85±1.03
17	211.15±38.02	10.24±0.53	10.09±0.33
18	137.23±34.45	11.65±0.72	10.36±0.68
19	135.87±35.11	11.95±1.37	10.66±0.52
20	134.51±23.35	12.25±0.56	10.96±0.91
Mean ±σ	184.90±40.92	11.50±0.78	10.20±0.71




Fig 2: Mean distribution of radionuclides in all samples

3.4 Soil to grass transfer factor

The plants take in deposited radionuclides from the soil, commonly expressed as soil-to-plant transfer factor (TF), which is widely used for determining the radiological effect. This parameter is an environmental transfer model, which is useful in the prediction of radionuclide concentration in plants/agricultural crops for estimating dose impact to human. Generally, the soil-to-plant transfer of radionuclides depends on soil type, pH, solid/liquid distribution coefficient, exchangeable K^+ , and organic matter contents. The transfer factor TFs for ^{40}K , ^{238}U and ^{232}Th were determined and the results were presented in Table 5.

The mean values of the transfer factors TFs were 0.29 ± 0.05 , 0.88 ± 0.10 and 0.80 ± 0.27 for ^{40}K , ^{238}U and ^{232}Th respectively. The TFs values for ^{40}K in the samples were lower compared to that of ^{238}U and ^{232}Th . The TFs values for ^{232}Th in the grass samples were smaller than the values obtained for ^{238}U , this was in conformity with the result obtained from other literature⁽¹⁴⁾. The values obtained in this study were higher than the range of values of 0.056 for ^{226}Ra , 0.089 for ^{232}Th and 0.275 for ^{40}K obtained for Bangladesh⁽²⁶⁾. The values reported in Table 6 could be traced to other sources such as drinking water and feed given to cattle at the rearing field.

Table 5: Transfer ratio of ^{40}K , ^{238}U and ^{232}Th for soil to grass samples

Samples	TF (^{40}K)	TF (^{238}U)	TF (^{232}Th)
1	0.26	0.75	0.82
2	0.27	0.83	0.91
3	0.28	0.91	1.02
4	0.35	0.98	0.25
5	0.30	0.89	0.33
6	0.29	0.80	0.30
7	0.28	0.73	0.30
8	0.48	0.84	0.99
9	0.24	0.84	0.92
10	0.24	0.86	0.86
11	0.25	0.89	0.90
12	0.25	0.91	1.05
13	0.25	1.05	0.97
14	0.30	1.09	0.89
15	0.31	1.05	0.86
16	0.32	0.89	0.82
17	0.26	0.81	0.97
18	0.30	0.82	0.90
19	0.28	0.84	0.93
20	0.26	0.85	0.96
Mean $\pm\sigma$	0.29 \pm 0.05	0.88 \pm 0.10	0.80 \pm 0.27







Fig 3: Mean soil-grass transfer factor of radionuclides in samples

Table 6: Difference in the estimated activity concentration in dung and grass samples (concentration in dung – concentration in grass)

Samples	^{40}K (Bqkg^{-1})	^{238}U (Bqkg^{-1})	^{232}Th (Bqkg^{-1})
1	115.96	1.90	0.04
2	100.2	1.34	0.32
3	84.44	0.96	0.06
4	115.9	2.20	5.12
5	87.07	1.11	7.01
6	89.04	1.80	7.30
7	32.97	2.49	7.59
8	5.73	1.40	0.02
9	27.86	1.86	1.69
10	21.30	1.96	2.00
11	14.74	1.70	2.31
12	108.66	0.21	0.06
13	112.50	0.18	0.74
14	108.05	0.48	0.51
15	103.60	0.78	0.32
16	102.47	2.00	0.82
17	100.59	0.75	0.02
18	17.10	1.87	0.82
19	19.26	1.91	0.64
20	21.42	1.95	0.46
Mean $\pm\sigma$	69.44 \pm 42.48	1.44 \pm 0.68	1.89 \pm 2.61

Fig 4: Distribution of ⁴⁰K concentrations in soil, grass, and dung.Fig 5: Distribution of ²³⁸U concentrations in soil, grass, and dung.Fig 5: Distribution of ²³²Th concentrations in soil, grass, and dung.

4.0 Discussion

The mean radioactivity levels of 291.8 ± 58.3 Bqkg⁻¹ for ⁴⁰K; 11.7 ± 0.8 Bqkg⁻¹ for ²³⁸U and 9.3 ± 2.0 Bqkg⁻¹ for ²³²Th reported in grass samples reported for Abeokuta ⁽¹⁴⁾ were higher than the corresponding values in the present study. However, the ranges of activity concentration of the radionuclides are similar to the values obtained for Tehran-Iran ⁽⁹⁾. The range of ⁴⁰K activity concentrations in the study was lower

when compared to the values reported for some vegetables consumed in Jordan ⁽²⁴⁾. The activity concentrations in cattle dung samples were higher than the corresponding values in the grass from the study area and this may be attributed to the ingestion of other sources of radionuclides such as the drinking water available to cattle in the grazing area. The activity concentrations of the radionuclides in water was reported to be high in Agege ⁽²⁵⁾.

The transfer factor values for ^{40}K in the samples were lower compared to that of ^{238}U and ^{232}Th . The transfer factor values for ^{232}Th in the grass samples were lower than the values obtained for ^{238}U , this was in conformity with the result obtained in other literature. The values obtained in this study were higher than the range of values obtained for Chittagong city of Bangladesh⁽²⁶⁾.

The transfer factor indicated that only a fractional amount of the radionuclides was transferred from the soil to the grass in the study. The results obtained showed that Uranium concentrations in the grass samples from the study area have low concentrations compared to that of Thorium but Uranium has the highest transfer factor. This is because Uranium is relatively susceptible to be soluble whereas Thorium is easily absorbed by the soil. The study indicated a possible low radiological effect on human feeding on cattle beef in the study area when compared to that earlier reported for Abeokuta.

5.0 Conclusion

The activity concentration of ^{40}K , ^{238}U and ^{232}Th in the samples collected at cattle-rearing field area of Mangoro-Agege, Lagos Nigeria were measured using a well calibrated NaI (TI) and a well-shielded detector coupled to a computer resident quantum multichannel analyzer. The radioactivity distribution in all samples was shown in Tables 2-4 and the transfer factor of radionuclides from soil-to-grass in Table 5. The mean activity concentration of ^{40}K , ^{238}U and ^{232}Th for the present study were lower than the results obtained at a cattle-rearing field area of Alabata in Abeokuta, Ogun state Nigeria. The results obtained were however higher than some values obtained in some other literature. The activity concentration of radionuclides in the soil were lower than the world average values. The soil samples exhibited the highest radioactivity levels followed by the cattle dung and the least was measured in the grass samples. It could be concluded that the cattle ingest radionuclides from other sources such as drinking water in the area that may constitute to the higher values in dung than grass, this is not in conformation with the study at Alabata cattle-rearing field. The transfer factor indicated that only a fractional amount of the radionuclides was transferred from the soil to the grass in the study area.

After carrying out the measurement of activity concentration of radionuclides in the samples collected from the study area, the result showed that the concentration in the cattle dung samples was higher than the concentration in the grass samples. Therefore, it is recommended that drinking water and feed is given to the cattle should be studied to ascertain the radioactivity levels in the water within the rearing

field. This is to affirm sources of a higher level of radionuclides in the cattle dung than the grass.

6.0 Acknowledgement

The Authors wish to thank Dr. Ishola of the Department of Physics and technical staff of the Radiation Physics Research Laboratory both of Ladoko Akintola University, Ogbomoso, Oyo State, Nigeria for allowing us the research laboratory for the sample analysis.

References

- 1 UNSCEAR (2008). United Nation Scientific Committee on Sources and Effects of Ionizing Radiation, Report to the General Assembly, United Nations (Vol. 1, p. 223). New York.
- 2 UNSCEAR (2000). United Nations Scientific Committee on Sources and Effects of Ionizing Radiation, Report on Effects of Atomic Radiation submitted to the General Congress by United Nations Scientific Committee, Volume II, 12-15.
- 3 L'Annunziata, M.F. (2007). Radioactivity, Introduction, and History, Amsterdam: Elsevier, B.V.
- 4 W.H.O (2006). Guidelines for Drinking Water Quality, 3rd Edition, World Health Organization., Geneva (Vol. 1, pp. 197-207).
- 5 NCRPM (1975). Natural Background Radiation in the United States by the National Council on Radiation Protection and Measurement. NCRP Report No. 45, Washington, D.C.
- 6 Faires, R.A. and Boswell, G.G.J. (1981). Radioisotope Laboratory Techniques (4th Edition), London: Butterworth & co. (publishers) Ltd.
- 7 Lilley, J. (2001). Nuclear Physics, principles/application, Chichester, John Wiley and Sons Ltd.
- 8 Nuclear Science Guide (2003). A Guide to the Nuclear Science Wall Chart, Contemporary Physics Education Project (CPEP), Chapter 3.
- 9 Hafezi, S., Amidi, J., and Attarilar, A. (2005). The concentration of Natural Radionuclides in Soil and Assesment of External Exposure to the Public in Tehran-Iran. Iran Journal of Radiation Resources, 3(2), pp. 85-88.
- 10 Salbu, B. Lind, O.C. and Skipperud, L. (2004). Radionuclide specification and its relevance in environmental impact assessment. Journal of Environmental Rad., vol 2, 25-57.
- 11 Abu-khadral, S.A., Abdel-sabour, M.F., Abdel-Fattah, A.T. and Eissa, H.S. (2008). Transfer factor of natural radionuclides from Egyptian soils to roots and leaves of the wheat plant, 15-19, Cairo.

- 12 Jibiri, N.N., Farai, I.P. and Alausa, S.K. (2006). Activity concentration of ^{226}Ra , ^{228}Th and ^{40}K in different food crops from a high background radiation area in Bitsichi, Jos Plateau, Nigeria. *Journal of Environmental Radioactivity*, 90(1), pp. 29 – 36.
- 13 Farai, I.P. and Jibiri, N.N. (2000). Baseline study of terrestrial outdoor gamma dose rate levels in Nigeria. *Radiological Protection Dosimeter*, 88(3), 247-254.
- 14 Alausa, S.K., Olabamiji, A.O. and Kola Odunaike (2015). Estimation of natural radionuclides in cattle from a rearing-field at Alabata, Abeokuta, Ogun State, South Western, Nigeria. *Journal of the National Association of Mathematical Physics*, vol 29, 353-362.
- 15 Ilori, A.O., Alausa, S.K. and Ajanaku, O. (2017). Activity Concentrations, Absorbed Dose Rate and Annual Effective Dose of Surface Soils of Agege, Lagos State, Southwestern Nigeria. *Journal of the Nigerian Association of Mathematical Physics*, volume 42, July 2017, pp 357-360. @J. of NAMP.
- 16 Obiora, D.N and Onwuka, O.S. (2005). Groundwater exploration in Ikorodu, Lagos Nigeria: A surface geophysical in survey contribution. *The Pacific Journal of Science and Technology*, vol 6, No 1, pp 86-93.
- 17 FRN (2006). National Population Census Report, Federal Republic of Nigeria. web.archive.org. Retrieved 12 November 2018.
- 18 Abbady, E.G.A., Uosif, M.A.M. and El-Taher, A. (2005). Natural radioactivity and dose assessment for phosphate rocks from Wadi El-Mashash and El-Mahamid mines, Egypt. *J. Environ. Radiat.*, 84, 65-78.
- 19 Ababneh, A.M. (2009). Radioactivity concentrations in soil and vegetables from the northern Jordan rift valley and the corresponding dose estimates. *Radiat. Prot. Dosim.*, 134, 30-37.
- 20 IAEA (1989). Measurements of Radionuclides in Food and the Environment, in International Atomic Energy Agency Tech. 1989, IAEA: Vienna.
- 21 Alharbi, A., and El - Taher, A. (2013). A study on transfer factors of radionuclides from soil to plant. *Life Science Journal* (2). <http://www.lifesciencesite.com>. Retrieved 22 October 2018.
- 22 Jibiri, N.N. and Ajao, A.O. (2005). Natural activities of ^{40}K , ^{238}U and ^{232}Th in elephant grass (*Pennisetum purpureum*) in Ibadan metropolis, Nigeria. *Journal of Environmental Radioactivity*, 78(1), pp. 105, Nigeria.
- 23 Alausa, S.K. (2012). Radioactivity in farm soils and food crops grown in Jos and Abeokuta Nigeria and its associated cancer risks. A Doctorate Dissertation, University of Ibadan, Nigeria.
- 24 Jamal Al-Jundi (2015). ^{226}Ra , ^{228}Ra and ^{40}K activity concentration in some vegetable consumed in Jordan and resultant annual ingestion effective dose. *Journal of Indian Association for Radiation and Protection (IARP)*, vol. 38, 29-34.
- 25 Ojo, T.J. and Afolayan, B.O. (2013). Distribution of radionuclide concentration with proximity to the Lagoon in Lagos state, Southwestern Nigeria. *Advance in physics Theories and Applications*, vol. 15, pp 5 – 7.
- 26 Shyamal Ranjan Chakraborty, Razaul Azim, A.K.M. Rezaul Rahman and Rashmi Sarker (2013). Radioactivity concentration in soil and transfer factors of radionuclides from soil to grass and plants in the Chittagong city of Bangladesh. *Journal of Physical Science*, vol. 24 (1), 95-113.

12/23/2019