

Stem Cell

Hongbao Ma, George Chen

Department of Medicine, Michigan State University, East Lansing, MI 48824, USA, hongbao@msu.edu

Abstract: The definition of stem cell is “an unspecialized cell that gives rise to a specific specialized cell, such as a blood cell”. Embryonic stem cells are derived from the inner cell mass of blastocyst stage embryos. Somatic stem cells differentiate into only the cells the specific tissue wherein they reside. Stem Cell is the original of life. All cells come from stem cells. [The Journal of American Science. 2005;1(2):90-92].

Key words: DNA; gene; life; protein; stem cell

1 Introduction

Stem cell is the origin of an organism's life. Stem cells have the potential to develop into many different types of cells in life bodies, that are exciting to scientists because of their potential to develop into many different cells, tissues and organs. Stem cells can be used in the clinical medicine to treat patients with a variety of diseases (Daar, 2003). Serving as a repair system for the living body, the stem cells can divide without limit to replenish other cells as long as the living body is still alive. When a stem cell divides, each new cell has the potential to either remain a stem cell situation or become another type of cell with a more specialized function, such as a muscle cell, a red blood cell, a bone cell, a nerve cell, or a brain cell. Stem cell research is a typical and important topic of life science.

2 Definition of Stem Cells

The definition of stem cell is “an unspecialized cell that gives rise to a specific specialized cell, such as a blood cell” (Stedman's Medical Dictionary, 2002).

3 Characterization of Stem Cell

Stem cell is totipotent, that means it holds all the genetic information of the living body and it can develop into a mature cell. Stem cell is a single cell that can give rise to progeny that differentiate into any of the specialized cells of embryonic or adult tissue. The ultimate stem cells (fertilized egg) divide to branches of cells that form various differentiated tissues or organs. During these early decisions, each progeny cell retains totipotency. Through divisions and differentiations the embryonic stem cells lose totipotency and gain differentiated function. During normal tissue renewal in adult organs, tissue stem cells give rise to progeny that differentiate into mature functioning cells of that tissue. Stem cells losing totipotentiality are progenitor cells. Except for

germinal cells, which retain totipotency, most stem cells in adult tissues have reduced potential to produce different cells.

4 Sources of Stem Cells

Aristotle (384-322 BC) deduced that the embryo was derived from mother's menstrual blood, which was based on the concept that living animals arose from slime or decaying matter. This concept was accepted in western world for over 2000 years, and it controlled western philosophy for over 2000 years either. In 1855, Virchow supposed that all cells in an organism are derived from preexisting cells. Now we know that all the human cells arise from a preexisting stem cell – the fertilized egg, that come from the mating of a man and a woman naturally but now can be produced in the laboratory tube. The counter hypothesis of spontaneous generation was accepted until 1864, when the French scientist Louis Pasteur demonstrated that there would be no microorganisms' growing after sterilizing and sealing.

The animal body has an unlimited source of stem cells, almost. However, the problem is not in locating these stem cells, but in isolating them from their tissue source.

Five key stem cells have been isolated from human: (1) Blastocysts; (2) Early embryos; (3) Fetal tissue; (4) Mature tissue; (5) Mature cells that can be grown into stem cells.

Up to today, only stem cells taken from adults or children (known generically as "adult stem cells") have been used extensively and effectively in the treatment of degenerative diseases.

5 Embryonic Stem Cell

Embryonic stem cells hold great promise for treating degenerative diseases, including diabetes, Parkinson's, Alzheimer's, neural degeneration, and cardiomyopathies (Bavister, 2005). Embryonic stem

cells are derived from the inner cell mass of blastocyst stage embryos. Embryonic stem cells can replicate indefinitely. This makes it feasible to culture the cells on a large scaled for cell transplantation therapy in clinical application. Embryonic stem cells are pluripotent and have the potential to differentiate into all three germ layers of the mammalian body including the germ cells.

6 Somatic Stem Cell

Normally to say that somatic stem cells differentiate only into specific tissue cells wherein they reside. However, somatic stem cells can differentiate into cells other than those of their tissue of origin. Adult bone marrow, fat, liver, skin, brain, skeletal muscle, pancreas, lung, heart and peripheral blood possess stem or progenitor cells with the capacity to transdifferentiate. Due to this developmental plasticity, somatic stem cells may have potential in autologous regenerative medicine, circumventing problems like rejection and the ethically challenged use of embryocyte stem cells.

7 Isolation and Characterisation of Stem Cells

As the example, the following is describing the isolation and characterization of the putative prostatic stem cell, which was done by Bhatt, Brown, Hart, Gilmore, Ramani, George, and Clarke in 2003. The detail methods have been described by Bhatt, Brown, Hart, Gilmore, Ramani, George, and Clarke in the article "Novel method for the isolation and characterisation of the putative prostatic stem cell" in the journal Cytometry A in 2003 (Bhatt, 2003).

7.1 Prostatic tissue collection and culture

When using human tissue, formal consent by the donor must be obtained before tissue collection. Tissue sections are obtained under sterile conditions. Each individual tissue section is bisected with half being sent for histological analysis for diagnostic evaluation and the remainder used for tissue culture. After then, tissue sections are chopped and placed in collagenase type I at 200 U/ml in RPMI 1640 medium with 2% v/v FCS overnight on a shaking platform at 37°C. The digest is then broken down further by shaking in 0.1% trypsin in PBS with 1% BSA and 1 mM ethylenediaminetetraacetic acid (EDTA) for 15-20 min. The cell suspension is then washed three times in PBS with 1% BSA and 1 mM EDTA before resuspending in RPMI 10% v/v FCS. Prostate epithelial cells are separated from fibroblasts by differential centrifugation (360 g, 1 min without braking). This process produced a supernatant enriched for fibroblasts and a pellet enriched for epithelia. The epithelial cell

suspension is then spun on a metrizamide gradient (1.079 g/ml), and the cells are isolated from the interface (Bhatt, 2003).

7.2 Ber-EP4/ α_2 /CD45 labelling of cells

Isolated epithelial cells are labeled at ambient temperature with either anti-human integrin α_2 monoclonal antibody or Ber-EP4 antibody (8 μ g/ml in 1% BSA/PBS) for 30 min before the addition of the secondary antibody, RAMBO (2.6 μ g/ml in 1% BSA/PBS) for 30 min. After washing with PBS, the cells are incubated for 20 min in the dark with streptavidin PE-Cy7 (20 μ g/ml). Samples are then dual labelled with CD45-FITC (1 μ g/ml in 1% BSA/PBS) for 30 min (Bhatt, 2003).

7.3 Ber-EP4/ α_2 and Hoechst labelling for flow cytometry

Isolated epithelial cells are labelled at ambient temperature with anti-human integrin α_2 monoclonal antibody (8 μ g/ml in 1% BSA/PBS) for 30 min before the addition of the secondary antibody, RAMBO (2.6 μ g/ml in 1% BSA/PBS) for 30 min. After washing with PBS, the cells are incubated for 20 min in the dark with streptavidin PE-Cy7 (20 μ g/ml). Hoechst staining could be performed by using the protocol for HSC as described by Rupesh, et al (Bhatt, 2003). Briefly, epithelial cells are resuspended in Hoechst buffer (Hanks' balanced salts solution, 10% FCS, 1% D-glucose, and 20 mM HEPES) and warmed to 37°C. Hoechst 33342 is then added to give a final concentration of 2 μ M and the cells incubated at 37°C for 2 h. Fifteen min before the end of incubation, the cells are labelled with monoclonal anti-human Ber-EP4 directly conjugated to FITC (8 μ g/ml). The cells are then washed in ice-cold Hoechst buffer before resuspending in ice-cold Hoechst buffer containing propidium iodide (PI) at 20 ng/ml (Bhatt, 2003).

7.4 Flow cytometry isolation of the SP fraction

Flow cytometry is carried out using a Becton Dickinson FACS Vantage SE flow cytometer. Hoechst 33342 is excited with an argon ion, ultraviolet-enhanced laser at 350 nm, and its fluorescence is measured with a 424/44 BP filter (Hoechst BLUE) and a 675DF20 BP optical filter (Hoechst RED; Omega Optical, Brattleboro VT). A 640 LP dichroic mirror is used to separate the emission wavelengths. PI fluorescence is also measured through the 675DF20 BP (having been excited at 350 nm). A second argon ion laser is used to excite the additional fluorochrome PE-Cy7 at 488 nm. PE-Cy7 is measured using a 787RDF40 (Omega Optical) filter (Bhatt, 2003).

7.5 Cell cycle characterisation of SP fraction

Epithelial cells are isolated and all fractions are resuspended in Hoechst buffer and warmed to 37°C. Hoechst 33342 is then added to give a concentration of 2 μ M and incubated at 37°C for 45 min. Pyronin Y (250 ng/ μ l) is added to each tube, and the samples are incubated for 45 min. Monoclonal anti-human Ber-EP4 FITC (8 μ g/ml) is added as appropriate 15 min before the end. After this, ice-cold Hoechst buffer is added immediately and the samples are washed then resuspended in ice-cold Hoechst buffer. The samples are analysed immediately by flow cytometry. Flow cytometry is performed using a modification of the method described above. Cells under study are selected by positive labelling for Ber-EP4 FITC before being analysed for Hoechst and Pyronin Y staining. These cells are then analysed by plotting the Hoechst profile on the x-axis and Pyronin Y along the y-axis in a linear scale (Bhatt, 2003).

7.6 Cytokeratin phenotype studies

Samples are processed as above, divided into two fractions, and labelled with either cytokeratin 8 or 14 indirectly conjugated to PE-Cy5. Samples are then dual labelled with Ber-EP4 FITC and integrin α_2 PE-CY7. Flow cytometry is performed as described and analysed on forward (FSC) and side (SSC) scatter (Bhatt, 2003).

8 Application of Stem Cells in Clinical Medicine

There are over four thousand registered diseases specifically linked to genetic abnormalities. Although stem cells are unlikely to provide powerful treatment for these diseases, they are unique in their potential application to these diseases.

Indeed, in many research projects, scientists have demonstrated that stem cells can be used to replenish or rejuvenate damaged cells within the immune system of the human body and that damaged stem cells can repair themselves and their neighbors. For example, in what is regarded as the first documented case of successful gene-therapy "surgery", scientists at the Necker Hospital for Sick Children in Paris of French succeeded in treating two infants diagnosed with Severe Combined Immunodeficiency Disease, a life-threatening degenerative disease caused by defects on the male (X) chromosome. With the identification of stem cell plasticity several years ago, multiple reports raised hopes that tissue repair by stem cell transplantation could be within reach in the near future (Kashofer, 2005). In cardiovascular medicine, the

possibility to cure heart failure with newly generated cardiomyocytes has created the interest of many researchers (Condorelli, 2005). Gene clone techniques can be widely used in the stem cell researches and applications (Ma, 2004).

9 Debates on Stem Cell Research

There are a lot of debates on the stem cell research. Stem cell research is a high-tech question and the people involved in this rebates should have certain scientific knowledge on the stem cell. It is OK for the politicians or religionists to show their opinions on any topic they are interested in, but not suitable for them to make decisions (or make laws) that will significantly influence the scientific research as this field the politicians or religionists are not specialized. Such as, it is not suitable for the American President George W. Bush to show the power in the stem cell research. It is scientists' job. When politics and science collide, science should do scientific way, rather political way. Major ethical and scientific debates surround the potential of stem cells to radically alter therapies in health care (Williams, 2005).

Correspondence to:

Hongbao Ma
B410 Clinical Center, Michigan State University
East Lansing, MI 48824, USA
Telephone: (517) 432-0623
Email: hongbao@msu.edu

References

- [1] Bavister BD, Wolf DP, Brenner CA. Challenges of primate embryonic stem cell research. *Cloning Stem Cells* 2005;7(2):82-94.
- [2] Bhatt RI, Brown MD, Hart CA, Gilmore P, Ramani VAC, George NJ, Clarke NW. Novel method for the isolation and characterisation of the putative prostatic stem cell. *Cytometry A*. 2003;54(2):89-99.
- [3] Condorelli G, Peschle C. Stem cells for cardiac repair: state of the art. *Front Biosci* 2005;10:3143-50.
- [4] Daar AS, Sheremeta L. The science of stem cells: ethical, legal and social issues. *Exp Clin Transplant*. 2003;1(2):139-46.
- [5] Kashofer K, Bonnet D. Gene Therapy Progress and Prospects: Stem cell plasticity. *Gene Ther*. 2005 (Epub ahead of print).
- [6] Ma H. Technique of Animal Clone. *Nature and Science* 2004;2(1):29-35.
- [7] Stedman's Medical Dictionary. The American Heritage®. Houghton Mifflin Company. <http://dictionary.reference.com/search?q=stem%20cell>. 2002.
- [8] Williams D. Stem cells in medical technology. *Med Device Technol* 2005;16(3):9-11.