

Assessment the Ameliorative Effect of Pomegranate and Rutin on Chlorpyrifos-ethyl-Induced Oxidative Stress in Rats

Mahgoub M. Ahmed¹, Nashwah I.Zaki²

¹Molecular Drug Evaluation Dep. and ²Physiology Dep.
National Organization for Drug Control & Research (NODCAR), Egypt
drosfscience@ymail.com

Abstract: Oxidative stress is one of the possible mechanisms resulted from organophosphate toxicity. Therefore, the aim of this study is to evaluate the *in vivo* effects of chlorpyrifos-ethyl (CE; 16.4 mg / kg / day body weight), on the serum and tissues antioxidant system of male albino rat and the efficacy of pomegranate peel extract (P; 500 mg/ kg/ day body weight) and rutin (R; 50 mg/ kg/ day body weight) as polyphenols to antagonize this response. The parameters were cholinesterase (ChE), acid phosphatase (ACP) and protein thiol (PrTh) in serum. Levels of malondialdehyde (MDA) as a marker of lipid peroxidation (LPO), reduced glutathione (GSH), glutathione-S-transferase (GST), and catalase (CAT) were estimated in liver, brain and kidney tissues. In addition, the activities of lysosomal enzymes (acid phosphatase, cathepsin D and RNase) in the liver were measured as early apoptosis marker. Administration of CE orally by gavage for two weeks induced a significant increase in serum ACP activity, LPO levels and liver lysosomal enzymes. Associated inhibitions in serum ChE activity and PrTh level were detected to CE exposure. Also, results showed significant decreases in GSH content, GST and CAT activities in liver, brain and kidney. Supplementation with P or R to treated animals was significantly ($P < 0.01$) attenuated the toxicity and oxidative stress evoked by CE. [Nature and Science. 2009;7(10):49-61]. (ISSN: 1545-0740).

Keywords: Oxidative stress, chlorpyrifos-ethyl, antioxidant system, pomegranate peel extract, rutin, acid phosphatase, cathepsin D, RNase

Introduction

Organophosphates (OPs) act mainly as acetylcholinesterase inhibitors (AChE) and can be an indicator of chronic toxicity of Ops (Tinoco and Halperine, 1998). Oxidative stress is one of the possible mechanisms that could be involved in the OPs toxicity. Oxidative stress is known to be a key factor in several diseases and was reported as a result of OPs exposure in human and experimental animals (Abou-Donia, 2003; Abdollahi et al., 2004; Milatovic et al., 2006; Dettbarn et al., 2006). Chlorpyrifos is a member of the most commonly used organophosphorus insecticide. As a result of widespread use, residues of chlorpyrifos have been detected in the air (Cattani et al., 2001) and in the crops (Sun et al., 2006; Atif Randhawa et al., 2007) which considered a risk for living organisms (Zhao et al., 2006). Chlorpyrifos, in particular chlorpyrifos-ethyl (CE), resulted in deleterious effects including hepatotoxicity, genotoxicity, teratogenicity, immunotoxicity as well as neurochemical and neurobehavioural alterations (Thrasher et al., 1993; Bagchi et al., 1995; Song et al., 1998; Dam et al., 1999; Gomes et al., 1999; Hunter et al., 1999). Previous studies has been shown that there is a correlation between acetylcholine inhibition and lipid peroxidation

level following subchronic exposure to OPs (Ranjbar et al., 2002; Akhgari et al., 2003). Exposure to chlorpyrifos increased levels of lipid peroxides in the rat liver, kidney, brain, and erythrocytes (Bagchi et al., 1995; Gultekin et al., 2001; Verma and Srivastava, 2001; Oncu et al., 2002 ; Tuzmen et al., 2008) and altered antioxidant enzymes in rat blood, liver, and lung (Bebe and Panemangalore, 2003). Moreover, administration of CE to pregnant rats induced oxidative stress and altered antioxidant system in liver, kidney, brain, and fetus (Zama et al., 2007).

Lysosomes are membrane bound structures that contain hydrolytic enzymes capable of degrading most of the cellular constituents. They are essential for controlled intracellular pathways such as autophagy, heterophagy and endocytosis. Apoptosis or programmed cell death which follows from moderate oxidative stress is preceded by partial lysosomal rupture and such lysosomal destabilization seems to be an initial event in apoptosis caused by a variety of other agents (Brunk et al., 2001). Exposure to organophosphorus insecticides has been shown to inhibit all cytoplasmic proteases and some of the lysosomal proteases in the liver tissue; the major site of insecticides metabolism (Mantle, 1997).

Polyphenols are one of the most abundant groups in plant which have a high antioxidant activity. Phenolic compounds are a biologically active group of phytochemicals. They are classified according to their chemical structure into flavonoids, phenolic acids, coumarins, and tannins (Tapiero et al., 2002; Mennen et al., 2005). Because of its potent anti-oxidant activity, pomegranate considers one of the commonly used natural anti-oxidants. Pomegranate fruit, juice, and peel extracts is a rich source of polyphenols and hence possess a potent antioxidant properties (Gil et al., 2000; Noda et al., 2002; Murthy et al., 2002; Singh et al., 2002). The effectiveness and safety of its isolated antioxidants have been tested (Cedra et al., 2003a, b). Murthy et al., 2002 added that methanolic extract of the peel has shown a higher anti-oxidant potential than that of seeds and could prevent CCl₄-induced hepatotoxicity. Recently, study by Elhalwagy et al., 2008 shown that supplementation with (60 mg/animal) green tea polyphenols, partially attenuate oxidative stress resulted from the toxic effect of fenitrothion insecticide, on the liver and kidney of rats. Also, tea polyphenols have been shown to protect against liver injury in animals intoxicated with chlorpyrifos insecticide (Khan and Kour 2007).

Flavonoids are a family of phenolic compounds (Harborne, 1986). It has become increasingly popular in terms of health protection because they possess a remarkable spectrum of biochemical and pharmacological activities. Flavonoids affect basic cell function such as growth, differentiation and apoptosis. Also, they were shown to be potent antioxidant because of their radical-scavenging activity; ability to complex heavy metal ion and to antagonize a broad spectrum of enzymes such as tyrosine protein kinase (Akiyama et al., 1987; Hollman et al., 1996; Knekt et al., 2002; Mira et al., 2002). Rutin, a flavonoid, has shown pharmacological benefits including anti-tumor (Deschner et al., 1991), anti-inflammatory (Aleksandrov et al., 1986), anti-diarrhoeal (Di Carlo et al., 1993), anti-mutagenic (Bear and Teel, 2000), myocardial protecting (Pozin et al., 1996), immunomodulator (Chen et al., 2000) and hepatoprotective activities (Janbaz et al., 2002). Literatures revealed that rutin increase of antioxidant capacity in the kidney of normal rats (Gao et al., 2002) as well as in a liposomal model (Nagasawa et al., 2003). Moreover, treatments of diabetic rats with rutin inhibit lipid peroxides while total protein and reduced glutathione were increased (Kamalkannan and Stanely, 2006). Based on our knowledge on free radicals and their involvement

in several diseases (Hogg, 1998) and the dependence on traditional medicine to replace ineffective medications, in this context, more research should be conducted to investigate the effectiveness of natural antioxidants (Madhavi and Salumkhe, 1995). The aim of this study was to investigate the ameliorative effect of pomegranate peel extract and rutin on toxicity, oxidative stress and apoptosis induced by chlorpyrifos-ethyl intoxication in male rats.

Material and methods

Plant Extraction

Pomegranate fruit peel was purchased from local market, dried and powdered. Amount of 500 g of the powdered plant material was extracted three times with ethanol (80%). The extracts were filtered, concentrated and freeze dried. The residue yielded was stored at 4°C for further analysis.

Chemicals

Chlorpyrifos-ethyl (CE) [O,O-diethyl-O-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 40% CE emulsion concentrate (Caribo™, Egychem) was purchased from local market and reconstituted to 1% solution in distilled water. Rutin, reduced glutathione, 1-chloro-2,4-dinitrobenzene (CDNB), bovine serum albumin and 5,5-dithiobis (2-nitrobenzoic acid) (DTNB), were purchased from Sigma (St Louis, MO, USA). All solvents used were HPLC grade (Merck, Darmstadt, Germany).

Animal

Sprague Dawley rats (200±30g) were obtained from the animal house of the National Organization for Drug Control and Research (NODCAR), Egypt. The animals were kept under standard laboratory conditions of light/dark cycle (12h/12h) and temperature (25 ± 2°C). They were provided with a nutritionally adequate standard laboratory diet.

Experimental Design

Sixty rats were treated according to the standard procedures laid down by OECD guidelines 407 (1992) repeated dose 28 days oral toxicity study in rodents. They are randomly allocated to six groups of six rats each as follow: Group 1 received distilled water by gavage orally and acts as control. Group 2 received CE (16.4 mg / kg / day bw) orally by gavage.

Group 3 received P (500 mg / kg / day bw) and CE (16.4 mg / Kg / day bw) orally by gavage
Group 4 received P (500 mg / kg / day bw) orally by gavage.

Group 5 received R (50 mg / kg / day bw) and CE (16.4 mg / Kg / day bw) orally by gavage.

Group 6 received R (50 mg / kg / day bw) orally by gavage.

Sampling

The animals were sacrificed by cervical decapitation after one and two weeks of exposure. Blood was collected and the separated serum was used for the estimation of ACP, ChE activities and PrTh content. Liver, brain and kidney tissues were removed quickly, washed in cold isotonic saline and homogenized in 50 mM phosphate buffer (pH 7) using an electronic homogenizer to prepare 10 % W/V homogenate. The homogenate was centrifuged at 3000 rpm for 10min at 4 °C by cooling centrifuge (Sigma 3K 30) to separate the nuclear debris. The supernatant were used for biochemical analysis.

Isolation of liver lysosomal fraction

A portion of liver homogenate was centrifuged at 4°C and 1000 xg for 10 min. The supernatant was centrifuged at 16.000 xg for 20 min to obtain lysosomal fraction. The residual pellets were then resuspended in phosphate buffer, pH 7.4 (Galvin Jr. et al., 1980).

Biochemical Analysis

The procedure used for the determination of cholinesterase activity in serum is a modification of Ellman et al. (1961) method as described by Gorun et al. (1978). Enzymatic activity of serum ACP was determined according to Moss (1984) using ready made kits by QCA, Spain. While serum protein-SH was measured by spectrophotometric method using 5'-5'-dithio-bis-2-nitrobenzoic acid according to Motchink et al., 1994. Lipid peroxidation (LPO) was measured by estimation of thiobarbituric acid reactive substances (TBARS) by the method of Ohkawa et al. (1979). Reduced glutathione (GSH) and glutathione -S-transferase (GST) contents in tissues were measured by the methods of Ellman (1959) and Habig et al. (1974), respectively. Catalase (CAT) activity was estimated by the method of Takahara et al. (1960). Protein was assayed by the method of Lowry et al (1951) using bovine serum albumin as standard. The activities of lysosomal enzymes (acid phosphatase, cathepsin D and RNase) were

measured according to the method of Barrett and Heath (1977) and Gianetto and De Duve (1955).

Statistical Analysis

Data were expressed as mean \pm standard deviation (SE). Differences in experimental groups were determined by one-way analysis of variance (ANOVA) followed by Dunnett's *t* test. $P < 0.01$ was considered to be statistically significant.

Results

Serum parameters

The present data revealed that chlorpyrifos ethyl-treated rats showed a significant ($P < 0.01$) increase in the activity of Acid phosphatase (ACP) when compared with the normal rats, while associated inhibition in ChE activity and Prth level were detected. Supplementation with pomegranate peel extract significantly countered this effect; while supplementation with R significantly ($P < 0.01$) restored ACP activity near the control values (Table 1).

LPO level

Exposures to 16.4 mg/kg chlorpyrifos-ethyl up to two weeks elicited a significant increase in lipid peroxidation of liver, brain and kidney of rat as measured by the estimation of thiobarbituric acid reactive substances (TBRAS). After one week, the increase in LPO levels ranged from 256.3%, 167.6% and 219.6%, in the liver, brain and kidney respectively. While after two weeks, the levels estimated as 391.9%, 317.1% and 252.1%, in the same tested tissues. Co-administration of pomegranate or rutin significantly ($P < 0.01$) modulated this increase. However, treatment with P and R showed a noticeably decrease in LPO levels comparing to the CE-treated group. Thus this study shows that both of the P and R protected cells from oxidative damage by decreasing LPO levels elevated as a result of CE treatment as showed in Figures 1, 2 and Table 2.

Table (1): Effect of pomegranate peel extract or rutin on serum activities of some enzyme as well as protein-thiol level induced by chlorpyrifos ethyl intoxication.

Groups	Time	Parameters		
		ACP (U/L)	ChE ($\mu\text{mol SH/hr/ml}$)	Prth ($\mu\text{mol/L}$)
C	W1	41.05 \pm 2.07	247.02 \pm 15.12	926.47 \pm 12.18
	W2	39.98 \pm 1.41	308.76 \pm 17.34	955.59 \pm 40.58
CE	W1	50.70 \pm 1.12a	174.84 \pm 7.45a	494.41 \pm 35.18a
	W2	49.27 \pm 1.72a	191.52 \pm 7.27a	535.00 \pm 17.92a
P+CE	W1	34.60 \pm 1.79b	176.82 \pm 15.58	836.47 \pm 67.00b
	W2	36.54 \pm 2.13b	250.2 \pm 19.51b	836.76 \pm 66.99b
P	W1	37.43 \pm 1.58b	212.34 \pm 10.16b	1044.62 \pm 78.53b
	W2	40.32 \pm 1.51b	256.68 \pm 14.14b	1025.59 \pm 55.6b
R+CE	W1	38.64 \pm 2.19b	176.88 \pm 5.22	824.12 \pm 32.61b
	W2	36.20 \pm 0.59b	187.8 \pm 10.24	544.41 \pm 31.01
R	W1	37.41 \pm 1.22b	225.96 \pm 11.11b	922.06 \pm 10.43b
	W2	37.94 \pm 2.24b	253.44 \pm 10.38b	836.18 \pm 46.88b

n=6; values are expressed as mean \pm SE

a indicates significant difference against control group at P <0.01

b indicates significant difference against chlorpyrifos-ethyl treated group (CPF) at P <0.01

Table (2): Effect of pomegranate peel extract or rutin on some oxidative stress parameters induced by chlorpyrifos ethyl intoxication in kidney.

Groups	Time	Parameters			
		LPO	GSH	CAT	GST
C	W1	1.40 \pm 0.09	4.73 \pm 0.15	172.10 \pm 1.92	199.80 \pm 3.30
	W2	1.20 \pm 0.10	4.86 \pm 0.16	175.20 \pm 2.60	200.30 \pm 2.01
CE	W1	5.10 \pm 0.12a	2.40 \pm 0.06a	109.70 \pm 3.61a	115.00 \pm 3.62a
	W2	6.10 \pm 0.23a	1.96 \pm 0.10a	98.20 \pm 2.21a	97.60 \pm 4.83a
P+CE	W1	2.70 \pm 0.13b	3.94 \pm 0.08b	138.10 \pm 1.34b	156.90 \pm 5.13b
	W2	2.50 \pm 0.13b	4.12 \pm 0.21b	145.20 \pm 2.61b	176.20 \pm 5.98 b
P	W1	0.96 \pm 0.05b	5.00 \pm 0.07b	169.80 \pm 2.70b	197.40 \pm 4.65b
	W2	0.90 \pm 0.06b	5.28 \pm 0.11b	172.30 \pm 4.91b	201.10 \pm 3.12b
R+CE	W1	2.10 \pm 0.09b	3.62 \pm 0.17b	129.70 \pm 2.75b	172.40 \pm 1.84b
	W2	1.90 \pm 0.11b	3.92 \pm 0.15b	132.40 \pm 4.27b	184.10 \pm 3.24b
R	W1	0.94 \pm 0.09b	5.24 \pm 0.12b	170.50 \pm 1.87b	190.80 \pm 5.66b
	W2	0.80 \pm 0.06b	5.44 \pm 0.15b	175.30 \pm 5.30b	200.80 \pm 4.93b

* Significant difference is indicated by ^aP <0.01 when compared with control group (C). ^bP <0.01 indicates significant difference when compared with chlorpyrifos-methyl-treated group (CPF).

n=6; values are expressed as mean \pm SE

a indicates significant difference against control group at P <0.01

b indicates significant difference against chlorpyrifos-ethyl treated group (CPF) at P <0.01 ;

GSH (nmol/ mg protein), GST (nmol/min/ mg protein), LPO (nmol/ mg protein), CAT (nmol/min/ mg protein)

Table (3): Effect of pomegranate peel extract or rutin on some lysosomal enzymes induced by chlorpyrifos ethyl intoxication in liver.

Groups	Time	Parameter		
		ACP (nmol/min/ mg protein)	Cathepsin D (U/ mg protein)	RNase II (U/ mg protein)
C	W1	0.48±0.01	35.6±1.14	0.36±0.02
	W2	0.51±0.02	36.9±0.89	0.38±0.01
CE	W1	0.98±0.04a	74.0±1.75a	1.07±0.04a
	W2	1.34±0.14a	82.1±2.22a	1.42±0.08a
P+CE	W1	0.68±0.02b	52.1±1.33b	0.72±0.01b
	W2	0.63±0.02b	49.7±0.93b	0.61±0.01b
P	W1	0.45±0.02b	35.4±1.86b	0.38±0.02b
	W2	0.53±0.01b	36.1±1.49b	0.39±0.01b
R+CE	W1	0.64±0.02b	53.1±0.83b	0.71±0.01b
	W2	0.53±0.03b	48.4±2.3b	0.61±0.02b
R	W1	0.45±0.02b	33.8±1.47b	0.38±0.01b
	W2	0.50±0.01b	35.5±1.38b	0.37±0.02b

n=6; values are expressed as mean±SE

a indicates significant difference against control group at P <0.01

b indicates significant difference against chlorpyrifos-ethyl treated group (CE) at P <0.0




Figure 1: Effect of pomegranate peel extract (P) or rutin (R) on LPO content in liver of rats-treated with chlorpyrifos-ethyl (CE).




Figure 2: Effect of pomegranate peel extract (P) or rutin (R) on LPO content in brain of rats-treated with chlorpyrifos-ethyl (CE).

GSH Level

In order to evaluate endogenous antioxidant system, glutathione levels were examined. Figures 3, 4 and table 2 showed the decrease in GSH levels in the same previous tissues after treatment with CE recording its

maximum value at the second week. While administration of P extract or R along with CE significantly increase GSH levels ($p < 0.01$) compared to CE-group.




Figure 3: Effect of pomegranate peel extract (P) or rutin (R) on GSH level in liver of rats-treated with chlorpyrifos-ethyl (CE).




Figure 4: Effect of pomegranate peel extract (P) or rutin (R) on GSH level in brain of rats-treated with chlorpyrifos-ethyl (CE).

* Significant difference is indicated by ^a $P < 0.01$ when compared with control group (C). ^b $P < 0.01$ indicates significant difference when compared with chlorpyrifos-methyl-treated group(CE).

CAT activity

Highlights the activity of the scavenger enzyme (CAT) in the same tissues, it was clearly demonstrated that tissues CAT inhibited significantly ($P < 0.01$) as a result of CE treatment when compared to the controls. Administration of P or R (500 mg/ kg , 50 mg/ kg respectively) along with CE caused a significant increase ($P < 0.01$) in CAT activity as compared to CE-treated rats and restored its activity near that of the control (Figures 5, 6 and table 2).




Figure 5: Effect of pomegranate peel extract (P) or rutin (R) on CAT activity in liver of rats-treated with chlorpyrifos-ethyl (CE).




Figure 6: Effect of pomegranate peel extract (P) or rutin (R) on CAT activity in brain of rats-treated with chlorpyrifos-ethyl (CE).

* Significant difference is indicated by ^a $P < 0.01$ when compared with control group (C). ^b $P < 0.01$ indicates significant difference when compared with chlorpyrifos-methyl-treated group(CE).

GST activity

A consistent and significant decrease was observed in GST activity (GSH dependent antioxidant enzyme) in the tissues of rats treated with CE. Supplementation of P or R significantly countered this decrease which restored GST values close to the control level as shown in Figures 7, 8 and table 2.




Figure 7: Effect of pomegranate peel extract (P) or rutin (R) on GST activity in liver of rats-treated with chlorpyrifos-ethyl (CE).




Figure 8: Effect of pomegranate peel extract (P) or rutin (R) on GST activity in brain of rats-treated with chlorpyrifos-ethyl (CE).

* Significant difference is indicated by ^a $P < 0.01$ when compared with control group (C). ^b $P < 0.01$ indicates significant difference when compared with chlorpyrifos-methyl-treated group(CE).

Lysosomal Enzymes activities

Effect of CE treatment on liver lysosomal enzymes (acid phosphatase, cathepsin D, and RNase II) was presented in Table (3). It concluded that CE treatment increase the activities of the three enzymes significantly ($P<0.01$) compared to control group. Administration of P or R along with CE significantly ($P<0.01$) ameliorated the effect of CE when compared to CE-group.

Discussion

Accumulated data from *in vitro* and *in vivo* studies shown that the primary mechanism of action and most acutely life threaten effect of OPs insecticides are related to accumulation of acetylcholine within the cholinergic synapses resulting inhibition of acetyl cholinesterase by active oxon metabolites (Milesso et al., 1998; Karanth et al., 2006; Gokcimen et al., 2007; Elhalwagy and Zaki, 2009). The present study conform the pervious finding, CE treatment reduced ChE activity as a marker of Ops toxicity. Pomegranate peel extract and rutin combined treatments restored ChE activity near to control level indicating their ameliorating effect. Oxidative stress biomarker is apparent in blood in association with AChE inhibition (Banerjee et al., 1999; Akhgari et al., 2003; Abdollahi et al., 2004; Shadnia et al., 2007). In line with the pervious studies we found decrease in protein thiol due to CE administration, which may be due to increased degradation of protein or increased consumption of this antioxidant in stress environment which confirm the role of OPs in disruption of body total antioxidant capacity (Ranjbar et al., 2002; Teimouri et al., 2006; Soltaninejad et al., 2007). Our results indicated that supplementations with P or R can reduce both the toxicity and oxidative stress of CE treatment. Recent literature pointed to the role of OPs in interference with metabolism of carbohydrate, biosynthesis of protein as well as respiration of mitochondria. Increased serum activity of ACP was also shown to CE treatment indicated damage to any or all of the organs producing this enzyme. While its activity was significantly lower in the groups co-treated with P or R, indicating their protective effects on the damage induced by CE on organs producing this enzymes, especially liver. This finding agrees with those (Khan et al., 2006; Elhalwagy et al., 2008; Gawish and Elhalwagy, 2009) using phenolic antioxidant against organophosphate toxicity.

In addition, pervious studies have been shown a correlation between inhibition of AChE and lipid peroxidation following subchronic and chronic exposure to OP (Ranjbar et al., 2002; Akhgari et al., 2003). Lipid peroxidation has considered one of the molecular pathways involved in the toxicity of OPs

(Datta et al., 1992). Organophosphates have been suggested to induce lipid peroxidation *in vivo* by enhancement of MDA production (Debnath and Mandal, 2000; Gultekin et al., 2001; Altuntas et al., 2002; Altuntas and Delibas, 2002; Oncu et al., 2002; Sharma et al., 2005). The present study shows that CE affected lipid peroxidation and defense mechanisms in rats. The obtained results show that CE may have properties to induce oxidative stress indicated by enhancement of MDA production, decrease in GSH content, GST and CAT activities in rat tissues. The increase of free radicals and lipid peroxidation may result from the inhibition of GSH levels induced by CE toxicity. The present findings are in agreement with other investigations indicating that accumulation of lipid peroxides has been resulted after exposure to acute dose of cholrpyrifos in rat liver (Bagshi et al., 1995), kidney (Oncu et al., 2002), brain (Gultekin et al., 2001), and erythrocytes (Verma and Srivastava, 2001). However, repeated doses increased LPO levels as well as antioxidant enzymes in blood, liver, and lung of rat (Ahmed et al., 2000; Bebe and Panemangalore, 2003; Akhgari et al., 2003). *In vitro* studies have been reported that accumulation of lipid peroxidation in human erythrocytes (Gultekin et al., 2000) and PC12 cells (Qiao et al., 2005). Supplementation with antioxidants effectively suppressed the oxidative damage induced by OPs (Ahmed et al., 2000; Gupta, 2001; Gultekin et al., 2001; Karaöz et al., 2002; Oncu et al., 2002; Camkayali et al., 2005; Elhalwagy et al., 2008). Antioxidant activity of pomegranate is referred to its polyphenolic capacity such as ellagic acid and ellagitannis (Seeram et al., 2005), which may suggest its role as an electron donor in scavenging free radicals (Kaur et al., 2006). Previous investigation revealed the ability of pomegranate fruit extract (Sudheesh and Vijayalakshmi, 2005; Noda et al., 2002) and peel extract (Singh et al., 2002) to suppress lipid peroxidation. Pretreatment with pomegranate flower for a period of one week significantly protected against oxidative damage and hepatic injury induced by Ferric nitrilotriacetate (Fe-NTA) by modulation of LPO levels and GSH content as well as antioxidant enzymes CAT and GST (Kaur et al., 2006). The same result has been obtained using the peel extract in mice treated with CCl_4 (Murthy et al., 2002). Ingestion of pomegranate juice for 4 weeks reduced hepatic oxidative stress (Faria et al., 2007) and improved the antioxidant enzyme activity in elderly subjects (Guo et al., 2008). Oral administration of pomegranate fruit extract inhibit LPO and increase GSH content as well as CAT activity in liver, kidney, and heart (Sudheesh and Vijayalakshmi, 2005). Previous studies revealed the effect of rutin on the activity on gene expression

of antioxidant enzymes in different experimental models both *in vivo* and *in vitro* (Lores-Arnaiz et al., 1995). In this context, treatment with rutin increased the activity of enzymatic antioxidants and also levels on non-enzymatic antioxidants in liver, kidney and brain of CE intoxicated rats. Inhibition of MDA levels in the group treated with both of CE and rutin was referred to the ability of rutin to transfer electrons and free radicals (Ferrali et al., 1997) in addition to activation of anti-oxidants enzymes (Elliott et al., 1992). However administration of rutin alone did not show any significant effect. The same finding has been obtained after pretreatment with rutin to isoproterenol-treated rats for 42 days (Karthick and Stanely Mainzen, 2006). Also, it has been reported that rutin has effectively reversed the biochemical, behavioral, and neurochemical changes in rat treated with haloperidol (Bishnoi et al., 2007) and improved the antioxidants enzymes system in human hepatoma cell line (Hep G2) by inhibition MDA levels and increasing CAT activity and therefore preventing or delay oxidative damage and its adverse effects (Alia et al., 2006). Moreover the antioxidant enzyme status was increased after rutin feeding in normal liver and in diabetic liver and kidney (Kamalakkannan et al., 2006).

In the present study CE intoxicated rats showed significant increase in the liver lysosomal enzymes activity (acid phosphatase, cathepsin D and RNase II). We suggested that reactive oxygen species (ROS) generated from CE pesticide may be responsible for the release of lysosomal enzymes, as a result decrease in membrane integrity and the leakage of enzymes from the enclosed sacs. These lead to intracellular dysfunction, disruption of potential substrates and organelles such as mitochondria, sarcolemma etc. (Kennett and Weglicki, 1978 and Mayanskaya et al., 2000). Also, the present study showed that induction of lysosomal enzymes was associated with a decrease in serum protein thiol level. Early study by Teimouri et al., 2006 confirms our finding and concludes that ROS can induce oxidation of critical sulfhydryl (SH) groups in protein and DNA, which will alter cellular integrity and function. The phospholipids-rich lysosomal membrane is a potential site of free radical attack subsequently causing loss of membrane stability. However, Brunk et al., 2001 added that apoptosis or programmed cell death is preceded by partial lysosomal rupture and such lysosomal destabilization seems to be an initial event in apoptosis. Treatment with pomegranate peel extract or rutin were able to decrease the release of lysosomal enzymes which could be due to the membrane stabilizing affect of P and R on the lysosomal membrane. The antioxidant properties of P and R scavenge the oxygen free

radicals and preservation of cellular viability serving secondarily to preserve lysosomes, thereby retaining near normal functioning of the lysosomes. Indeed, cathepsin D is lysosomal proteases possibly involved in autophagic of discrete areas of cytoplasm, myofibrillary and mitochondria proteins (Zak et al., 1976). Cell death causing by oxidants may be initiated by lysosomal rupture and that this latter event may involve intralysosomal iron which catalyze Fenton-type chemistry and resultant peroxidative damage to lysosomal membrane. The antioxidant activity of rutin and pomegranate peel extract stopped this reaction by iron chelating activity. According to our results, pomegranate peel extract and rutin have shown abilities to preserve the activity of anti-oxidant enzymes and lysosomal membrane which may be referred to its role in modulating the levels of H₂O₂ and O₂. GST over expression after co-treatment with the extract, and CE may be attributed to the increase in oxidative damage. Furthermore, the contents of the extract have been suggested to induce the de novo synthesis of anti-oxidant enzymes by acting as several loci in the metabolic pathway.

Correspondence to:

Mahgoub M. Ahmed
Molecular Drug Evaluation Dep. and ² Physiology Dep.
Emaildrofscience@ymail.com

Conclusion

The present study highlights the protective role of pomegranate and rutin against OPs pesticides but the mechanism involved is still unclear. Further work is required to clarify how the plants extract works to enhance the antioxidant enzymes or their gene expression.

References

1. Abdollahi M, Ranjbar A, Shadnia S, Nikfar S and Rezaie OE. Pesticides and oxidative stress: A review. *Med. Sci. Monit* 2004; 10:141-147.
2. Abou-Donia M B, Abdel-Rahman A A, Goldstein L B and et al. Sensorimotor deficits and increased brain nicotinic acetylcholine receptors following exposure to chlorpyrifos and/or nicotine in rats. *Arch Toxicol* 2003; 77:452-458.
3. Ahmed R S, Seth V, Pasha S T and Banerjee B D. Influence of dietary ginger (*Zingiber Officinale* Rose) on oxidative stress induced by malathion in rats. *Food Chem. Toxicol* 2000; 38:443-450.
4. Akhgari M, Abdollahi M, Kebryaezadeh A, Mosseini R and Sabzevari O. Biochemical evidence for free radical-induced lipid peroxidation as a mechanism for toxicity of malathion in blood and liver of rats. *Human Exp. Toxicol* 2003; 22: 205-211.
5. Akiyama T, Ishida J, Nakagawa S, Ogawara H, Watanabe S, Itoh N, Shibuya M and Fukami Y.

- Genistein, a specific inhibitor of tyrosine-specific protein kinases. *J. Biol. Chem.* 1987; 262: 5592-5595.
6. Aleksandrov P N, Speranskaia T V, Bobkov Iu G, Zagorevskii V A and Zykov D A. Effect of rutin and esculamineon models of a septic inflammation. *Farmakol Toksiko* 1986; 149: 84-86.
 7. Alia M, Mateos R, Ramos S, Le cumberri E, Bravo L, Goya L. Influence of quercetin and rutin on growth and antioxidant defense system of a human hepatoma cell line (HepG2). *Eur. J. Nutr.* 2006; 45: 19-28.
 8. Altuntas I and Delibas N. The effects of Fenthion on lipid peroxidation and some liver enzymes: the possible protective role of vitamins E and C. *Turkish Journal of Medical Sciences* 2002; 32: 293-297. Altuntas I, Delibas N and Sutcu R. The effects of organophosphate insecticide methidathion on lipid peroxidation and antioxidant enzymes in rat erythrocytes: Role of vitamins E and C. *Hum. Exp. Toxicol.* 2002; 21: 681-685.
 9. Atif Randhawa M, Muhammad Anjum F, Ahmed A and Saqib Randhawa M. Field incurred chlorpyrifos and 3, 5, 6-trichloro-2-pyridinol residues in fresh and processed vegetables. *Food Chemistry* 2007; 103:1016-1023.
 10. Bagchi D, Bagchi M, Hassoun E A and Stohs S J. In vitro and in vivo generation of reactive oxygen species, DNA damage and lactate dehydrogenase leakage by selected pesticides. *Toxicology* 1995; 104: 129-140.
 11. Banerjee B D, Seth V, Bhattacharya A, Pasha S T, Chakraborty A K. Biochemical effects of some pesticides on lipid peroxidation and free-radicalscavengers. *Toxicol Lett.* 1999; 107: 33-47.
 12. Barrett AJ, Heath MF. In Lysosomes: A laboratory handbook ed. Dingle JT (North Holland Publishing Co. Amsterdam) 1977; pp 19-127.
 13. Bear W L and Teel R W. Effects of citrus flavonoids on the mutagenicity of heterocyclic amines and on cytochrome P450 1A2 activity. *Anticancer Res.* 2000; 20: 3609-3614.
 14. Bebe F N and Panemangalore M. Exposure to low doses of endosulfan and chlorpyrifos modifies endogenous antioxidants in tissues of rats. *J. Environ. Sci. Health B* 2003; 38(3): 349-363.
 15. Bishnoi M, Chopra K and Kulkarni S K. Protective effect of rutin, a polyphenolic flavonoid against haloperidol-induced orofacial dyskinesia and associated behavioural, biochemical and neurochemical changes. *Fundam Clin Pharmacol.* 2007; 21(5): 521-9.
 16. Brunk U T, Neuzil J and Eaton J W. Lysosomal involvement in apoptosis. *Redox Report* 2001; 6: 91-97.
 17. Camkayali I, Demirag K, Eris O, Ersoz B and Moral A R. The effect of N-acetylcysteine on oxidative stress in organophosphate poisoning model. *Adv. Ther.* 2005; 22:107-116.
 18. Cattani M, Cena K, Edwards J and Pisaniello D. Potential dermal and inhalation exposure to chlorpyrifos in Australian pesticide workers. *The Annals of Occupational Hygiene* 2001; 45: 299-308.
 19. Cerda B, Ceron J J, Tomas-Barberan F A and Espin J C. Repeated oral administration of high doses of the pomegranate ellagitannin punicalagin to rats for 37 days is not toxic. *J. Agric. Food Chem.* 2003 a; 51: 3493-3501.
 20. Cerda B, Llorach R, Ceron J J, Espin J C and Tomas-Barberan F A Evaluation of the bioavailability and metabolism in the rat of punicalagin, an antioxidant polyphenol from pomegranate juice. *Eur. J. Nutr.* 2003 b; 42:18-28.
 21. Chen S, Gong J, Liu F and Mohammed U. Naturally occurring polyphenolic antioxidants modulate IgE-mediated mast cell activation. *Immuology* 2000; 100: 471-480.
 22. Dam K, Garcia S J, Seidler F J and Slotkin T A. Neonatal chlorpyrifos exposure alters synaptic development and neuronal activity in cholinergic and catecholaminergic pathways. *Brain Res Dev Brain Res.* 1999; 116: 9-20.
 23. Datta J, Gupta J, Sarkar A and Sengupta D. Effects of organophosphorous insecticide phosphomidon on antioxidant defense components of human erythrocyte and plasma. *Indian J. Exp. Biol.* 1992; 30: 65-67.
 24. de Nigris F, Williams-Ignarro S, Lerman L O, Crimi E, Botti C, Mansueto G, D'Armiento F P, De Rosa G, Sica V, Ignarro L J and Napoli C. Beneficial effects of pomegranate juice on oxidation-sensitive genes and endothelial nitric oxide synthase activity at sites of perturbed shear stress. *Proceedings of the National Academy of Sciences*, 2005; USA. 102: 4896-4901.
 25. Debnath D and Mandal K. Study of quinalphos (an environmental oestrogenic insecticide) formulation (Ekalux 25 E.C.)-induced damage of the testicular tissues and antioxidant defense systems in Sprague-Dawley albino rats. *J Appl Toxicol.* 2000; 20 (3): 197-204.
 26. Deschner E E, Ruperto J, Wong G and Newmark H L. Quercetin and rutin as inhibitors of azoxymethanol-induced colonic neoplasia. *Carcinogenesis* 1991; 12: 1193-1196.
 27. Dettbarn W F, Milatovic D and Gupta R C. Oxidative stress in anticholinesterase-induced excitotoxicity. In: R.C. Gupta, Editor, *Toxicology of Organophosphate and Carbamate Compounds*, Elsevier, Amsterdam. 2006; pp. 511-532.
 28. Di Carlo G, Autore G, Izzo A A, Maiolino P, Mascolo N, Viola P, Diurno M V and Capasso F. Inhibition of intestinal motility and secretion by flavonoids in mice and rats: structure-activity relationships. *J Pharm Pharmacol.* 1993; 45:1054-1059.
 29. Elhalwagy M E A, Darwish N S and Zaher E M. Prophylactic effect of green tea polyphenol against liver and kidney injury induced by fenitrothion insecticides. *Pesti.Bio.Physio.* 2008; 91: 81-89.
 30. Elhalwagy M E A and Zaki N I. Comparative study on pesticide mixture of organophosphorus and pyrethroid in commercial formulation. *Environ. Toxicol. Pharmacol* 2009; 28:219-224.
 31. Elliott A J, Scheiber S A, Thomas C and Pardini R S. Inhibition of glutathione reductase by flavonoids. *Biochem Pharmacol.* 1992; 44:1603-1608.

32. Ellman G L. Tissue sulfhydryl groups. *Arch. Biochem. Biophys.* 1959; 82: 72- 77.
33. Ellman G L, Courtney K D, Andres V and Featherstone R M. A new and rapid colorimetric determination of cholinesterase activity. *Biochem. Pharmacol.*, 1961; 7: 88-95.
34. Faria A, Monteiro R, Mateus N, Azevedo I and Calhau C. Effect of pomegranate (*Punica granatum*) juice intake on hepatic oxidative stress. *Eur. J. Nutr.* 2007; 46:271- 278.
35. Ferrali M, Signofrini C, Caciotti B, Sugherini L, Ciccoli D, Giachetti D and Comporti M. Protection against oxidative damage of erythrocyte membranes by the flavonoid quercetin and its relation to iron chelating activity. *FEBS Letters* 1997; 416: 123–139.
36. Galvin Jr M J, Parks D L and McRee D I. Microwave irradiation and in vitro release of enzymes from hepatic lysosomes. *Radiat Environ Biophys.* 1980; 18(2): 129-136.
37. Gao Z, Xu H and Huang K. Effects of rutin supplementation on antioxidant status and iron, copper and zinc contents in mouse liver and brain. *Biol Trace Elem Res.* 2002; 88: 271–279.
38. Gawish A and Elhalwagy M E A. Which can attenuate hepatotoxicity induced by pesticides mixture natural or synthetic phenolic antioxidant. *Nature and Science* 2009; 7 (5): 29-44.
39. Gianetto R and De Duve C. Tissue fractionation studies. 4-Comparative study of binding of acid phosphatase, beta glucuronidase and cathepsin by rat liver paraticle. *Biochem. J.* 1955; 59: 433- 438.
40. Gil M I, Tomas-Barberan F A, Hess-Pierce B, Holcroft D M and Kader A A. Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. *Journal of Agricultural and Food Chemistry* 2000; 48: 4581– 4589.
41. Gokcimen A, Gulle K, Demirin H, Bayram D, Koca, A and Altuntas I. Effect of diazinon at different doses on rat liver and pancreas tissues. *Pestic. Biochem Physiol.* 2007; 87:103–108.
42. Gomes J, Dawodu A H, Llyod O, Revitt D M and Anilal S V. Hepatic injury and disturbed amino acid metabolism in mice following prolonged exposure to organophosphorus pesticides. *Hum Exp Toxicol* 1999; 18:33-37.
43. Gorun V, Proinov I, Baltescu V, Balaban G and Barzu O. Modified Ellman procedure for assay of cholinesterase in crude enzymatic preparation. *Anal. Biochem.* 1978; 86: 324-326.
44. Gultekin F, Delibas N, Yasar S and Kiline I. In vivo changes in antioxidant systems and protective role of melatonin and a combination of vitamin C and vitamin E on oxidative damage in erythrocytes induced by chlorpyrifos-ethyl in rats. *Arch. Toxicol.* 2001; 75: 88-96.
45. Gultekin F, Ozturk M and Akdogan M. The effect of organophosphate insecticide chlorpyrifos-ethyl on lipid peroxidation and antioxidant enzymes (in-vitro). *Arch. Toxicol.* 2000; 74:533-538.
46. Guo C, Wei J, Yang J, Xu J, Pang W and Jiang Y. Pomogranate juice is potentially better than apple juice in improving antioxidant function in elderly subjects. *Nutr. Res.* 2008; 28: 72-77.
47. Gupta R C. Depletion of energy metabolites following acetylcholinesterase in leptazol-induced status epilepticus and protection by antioxidants. *Neurotoxicol.* 2001; 22: 271-282.
48. Habig W H, Pabet J and Jakoby W B. Glutathione S-transferases: the first enzymatic step in mercapturic acid formation, *J. Biol. Chem.* 1974; 249: 7130–713.
49. Harborne J B. Plant phenolics. In: E.A. Bell and B.V. Charlwood (eds). *Encyclopedia of plant physiology*, Vol. 8. Secondary plant products. Springer Verlag, Berlin. 1986, pp: 329–395.
50. Hollman, P C H, Hertog M G L and Katan M B. Role of dietary Flavonoids in protection against cancer and coronary heart disease. *Biochem Soc.Trans.*, 1996; 24:785–789.
51. Hunter D L, Lassiter T L and Padilla S. Gestational exposure to chlorpyrifos: comparative distribution of trichloropyridinol in the fetus and dam. *Toxicol Appl Pharmacol.*, 1999; 158:16-23.
52. Janbaz K H, Saeed S A and Gilani A H. Protective effect of rutin on paracetamol and CCl4 induced hepatotoxicity in rodents. *Fitoterapia* 2002; 73:557–563.
53. Kamalakkannan N and Stanely Mainzen P. Rutin improve the antioxidant status in streptozotocin-induced diabetic rat tissues. *Molecular and Cellular Biochemistry* 2006; 293: 211–219.
54. Karanth S, Liu J, Mirajkar N, Pope C. Effects of acute chlorpyrifos exposure on in vivo acetylcholine accumulation in rat striatum. *Toxicol Appl Pharmacol.* 2006; 216 (1):150-156.
55. Karaöz E, Gultekin F, Akdogan M, Oncu M and Gokcimen A. Protective role of melatonin and a combination of vitamin C and vitamin E on lung toxicity induced by chlorpyrifos-ethyl in rats. *Exp. Toxicol. Pathol.* 2002; 54: 97-108.
56. Karthick M and Stanely Mainzen P. Preventive effect of rutin, a bioflavonoid, on lipid peroxides and antioxidants in isoproterenol-induced myocardial infarction in rats. *J Pharm Pharmacol.* 2006; 58 (5): 701-707.
57. Kaur G, Jabbar Z, Athar M, Alam S. *Punica granatum* (pomegranate) flower extract possesses potent antioxidant activity and abrogates Fe-NTA induced hepatotoxicity in mice. *Food and Chemical Toxicology* 2006; 44: 984 - 993.
58. Kennett F F and Weglicki W B. Effect of well-defined ischemia on myocardial lysosomal and microsomal enzymes in a canine model. *Circ. Res.* 1978; 43: 750-758.
59. Khan S M. Protective effect of black tea extract on the levels of lipid peroxidation and antioxidant enzymes in liver of mice with pesticide-induced liver injury. *Cell Biochem and funct* 2006; 24: 327-332.
60. Khan S M and Kour G. Subacute oral toxicity of chlorpyrifos and protective effect of green tea extract. *Pesti.Bio.Physio.* 2007; 89:118 -123.
61. Knekt P, Kumpulainen J, Jarvinen R, Rissanen H, Heliovaara M, Reunanen A, Hakulinen T and Aromaa

- A. Flavonoid intake and risk of chronic diseases. *Am. J. Clin. Nutr.* 2002; 76: 560–568.
62. Lores-Arnaiz S, Llesuy S, Curtin J C and Boveris A. Oxidative stress by acute acetaminophen administration in mouse liver. *Free Radic. Biol. Med.* 1995; 19: 303- 310.
 63. Lowry O H, Rosebrough N J, Farr A L and Randall R J. Protein measurement with Folin phenol reagent. *J. Biol. Chem.* 1951; 193: 265 –275.
 64. Madhavi D L and Salunkhe D K. Toxicological aspects of food antioxidants. In: D.L., Madhavi; S.S., Deshpande and Salunkhe, D.K. (Eds.) Food antioxidants, Dekker, New York. 1995; pp. 267.
 65. Mantle D. Effect of pirimiphos methyl on proteolytic enzyme activities in rat heart, kidney, brain and liver tissue in vivo. *Clin Chim Acta* 1997; 262: 89-97.
 66. Mayanskaya S D, Mayanskaya N N, Efremov A V and Yakobson G S. Activity of lysosomal apparatus in rat myocardium during experimental coronary myocardial damage. *Bull Exp.Mod.* 2000; 129: 530-532.
 67. Mennen L I, Walker R, Bennetau-Pelissero C and Scalbert A. Risks and safety of polyphenol consumption. *Am J Clin Nutr.* 2005; 81:326S–329S.
 68. Milatovic D, Gupta R C and Aschner M. Anticholinesterase toxicity and oxidative stress. *Scientific World J.* 2006; 6: 295-310.
 69. Mileson B E, Chambers J E, Chen W L, Dettbran W and Ehrich M. Common mechanism of toxicity: a case study of organophosphorus pesticides. *Toxicol. Sci.* 1998; 41: 8-20.
 70. Mira L, Fernandez M T, Santos M, Rocha R, Florencio M H and Jennings K R. Interactions of Flavonoids with iron and copper ions: a mechanism for their antioxidant activity. *Free Radic. Res.*2002; 36:1199–1208.
 71. Moss D W. In Methods of enzymatic analysis, Bergmeyer, H.U. (Ed),Veriag-Chemie 3rd edition 1984; V4: 92-106.
 72. Motchnik A P, Frei B, Ames N B. Measurement of antioxidants in human blood plasma: Protein thiols. In: Packer L, editor. Oxygen radicals in biological systems. Methods in Enzymology, Academic Press: California; 1994. p. 234 (D): 273-4.
 73. Murthy K NC, Jayaprakasha G K and Singh R P. Studies on antioxidant activity of pomegranate peel extract using in vivo models. *Journal of Agricultural and Food Chemistry* 2002; 50: 4791– 4795.
 74. Nagasawa T, Tabata N, Ito Y, Nishizawa N, Aiba Y and Kitts D D. Inhibition of glycation reaction in tissue protein incubations by water soluble rutin derivative. *Mol Cell Biochem.* 2003; 249: 3–10.
 75. Noda Y, Kaneyuka T, Mori A and Packer L. Antioxidant activities of pomegranate fruit extract and its anthocyanidins: delphinidin, cyanidin, and pelargonidin. *Journal of Agricultural and Food Chemistry* 2002; 50: 166–171.
 76. Ohkawa H, Ohishi N and Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* 1979; 95: 351–358.
 77. Oncu M, Gultekin F, Karaöz E, Altuntas I and Delibas N. Nephrotoxicity in rats induced by chlorpyrifos-ethyl and ameliorating effects of antioxidants. *Human Exp. Toxicol.* 2002; 21: 223-230.
 78. Organization for Economic cooperation and development (OECD). chairman's Report of the meeting of the ad hoc working group of experts on systemic short term and (delayed) neurotoxicity. 1992.
 79. Pozin V M, Skuratovskaia SG and Pocheptsova G A. Changes in the vascular wall and ischemic damages to the myocardium in reversible episodes of heart muscle ischemia. *Fiziologicheskii Zhurnal.*1996; 42:10–16.1
 80. Qiao D, Seider F J and Slotkin T A. Oxidative neurotoxicity of nicotine and chlorpyrifos. *Toxicol. Appl. Pharmacol.* 2005; 206: 17-26.
 81. Ranjbar A, Pasalar P and Abdollahi M. Induction of oxidative stress and acetylcholinesterase inhibition in organophosphorous pesticide manufacturing workers. *Hum. Exp. Toxicol.* 2002; 21:179-182.
 82. Seeram N, Adams L, Hennig S, Niu Y, Zhan Y, Nair M and Heber D. In vitro anti-proliferative, apoptotic and antioxidant activities of punicalagin, ellagic acid and a total pomegranate tannin extract are enhanced in combination with other polyphenols as found in pomegranate juice. *J. Nutr. Biochem.* 2005; 16: 360-367.
 83. Shadnia S, Dasgar M, Taghikhani S, Mohammadirad A, Khorasani R, Abdollahi M. Protective effects of α -tocopherol and N-acetyl-cysteine on diazinon-induced oxidative stress and acetylcholinesterase inhibition in rats. *Toxicology Mechanisms and Methods*, 2007; 17(2): 109-115.
 84. Sharma Y, Bashir S, Irshad M, Nag T C and Doqra T D. Dimethoate-induced effects on antioxidant status of liver and brain of rat following subchronic exposure. *Toxicology* 2005; 215 (3):173-181.
 85. Singh R P, Murthy K N C and Jayaprakasha G K. Studies on the antioxidant activity of pomegranate (*Punica granatum*) peel and seed extracts using in vitro models. *Journal of Agricultural and Food Chemistry* 2002; 50: 81–86.
 86. Soltaninejad K, Shadnia S, Afkhami-Taghipour M, Saljooghi R, Mohammadirad A and Abdollahi, M. Blood $^{2-}$ glucuronidase as a suitable biomarker at acute exposure of severe organophosphorus poisoning in human. *Hum Exp Toxicol* 2007; 26: 963
 87. Song X, Violin J D, Seidler F J and Slotkin T A. Modeling the developmental neurotoxicity of chlorpyrifos in vitro: macromolecule synthesis in PC12 cells. *Toxicol Appl Pharmacol.*1998;15:182-191.
 88. Sudheesh S and Vijayalakshmi N R. Flavonoids from *Punica granatum* potential antiperoxidative agents. *Fitoterapia* 2005; 67: 181-186.
 89. Sun F, Wong S S, Li G C and Chen S N. A preliminary assessment of consumer's exposure to pesticide residues in fisheries products. *Chemosphere* 2006; 62: 674–680.
 90. Takahara S, Hamilton B M, Nell J V, Ogura Y and Nishimura E T. Hypocatalasemia, a new genetic carrier state. *J. Clin. Invest.* 1960; 29: 610- 619.
 91. Tapiero H, Tew K D, Ba G N and Mathe G. Polyphenols: do they play a role in the prevention of

- human pathologies?. *Biomed Pharmacother.* 2002; 56: 200-207.
92. Teimouri F, Amirkabirian N, Esmaily H, Mohammadirad A, Ali ahmadi A, Abdollahi M. Alteration of hepatic cells glucose metabolism as a non cholinergic detoxification mechanism in counteracting diazinon induced oxidative stress. *Hum.Exp.Toxicol* 2006; 25: 697-703.
93. Thrasher J D, Madison R and Broughton A. Immunologic abnormalities in humans exposed to chlorpyrifos: preliminary observations. *Arch Environ Health.* 1993; 48: 89-93.
94. Tinoco R, Halperine D. Poverty production and health: inhibition of erythrocyte cholinesterase via occupational exposure to organophosphate insecticides in Chipas, Mexico. *Arch Environ Health* 1998; 53: 29-35.
95. Tuzmen N, Candan N, Kays E and Demiryas N. Biochemical effects of chlorpyrifos and deltamethrin on altered antioxidative defense mechanisms and lipid peroxidation in rat liver. *Cell Biochem.Funct.* 2008; 26:119-124.
96. Verma R S and Srivastava N. Chlorpyrifos induced alterations in levels of thiobarbituric acid reactive substances and glutathione in rat brain. *Indian J. Exp. Biol.* 2001; 39: 174-177.
97. Zak R, Martin A F, Reddy M K, Rabinowitz M. Control of protein balance in hypertrophied cardiac muscle. *Circ Res* 1976; 38 (5 suppl):1145.
98. Zama D, Meraihi Z, Benayssa W, Benayache F, Benyache S and Vlietinck A J. Chlorpyrifos- induced oxidative stress and tissue damage in the liver, kidney, brain and fetus in pregnant rats: The protective role of the butanolic extract of *Paronychia argentea* L. *Indian J. Pharmacol.* 2007; 39: 145 -150.
99. Zhao Q, Dourson M and Gadagbui B. A review of the reference dose for chlorpyrifos. *Regul Toxicol Pharmacol.* 2006; 44(2):111-24.

7/5/2009