

Repellent effect of leaves essential oils from *Eucalyptus globules* (Mirtaceae) and *Ocimum basilicum* (Lamiaceae) against two major stored grain insect pests of Coleopterons

Bhuwan Bhaskar Mishra^{1,2}, S.P. Tripathi¹ and C.P.M. Tripathi²

1. P.G. Department of Zoology, M.G.P.G. College, Gorakhpur, 273 001, India

2. Department of Zoology, D.D.U, Gorakhpur University, Gorakhpur, 273 009, India

b2mishra123@gmail.com

Abstract: The essential oils were extracted by water distillation method from two medicinal plants *Eucalyptus globules* (Mirtaceae) and *Ocimum basilicum* (Lamiaceae) leaves and its repellent activity was investigated against two major economic important stored-grain insect pests red flour beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae) and rice weevil *Sitophilus oryzae* L. (Coleoptera: Curculionide) adults. The results indicated that the repellency of *E. globules* and *O. basilicum* was 9.16 ± 0.30 and 8.50 ± 0.22 for *T. castaneum* and 8.66 ± 0.33 and 8.16 ± 0.30 for *S. oryzae*. The repellency of both insect pests increased with concentration from 0.05% to 0.40% at exposure time of 4 h. These observations confirmed that the essential oils which were extracted from these two medicinal plants leaves have more repellency against both insect pests.

[Bhuwan Bhaskar Mishra, S.P. Tripathi and C.P.M. Tripathi. **Repellent effect of leaves essential oils from *Eucalyptus globules* (Mirtaceae) and *Ocimum basilicum* (Lamiaceae) against two major stored grain insect pests of Coleopterons.** Journal of Nature and Science 2012;10(2):50-54]. (ISSN: 1545-0740). <http://www.sciencepub.net>. 9

Key words: Repellency; insect pests; rice weevil; flour beetle; essential oils

Introduction

Rice weevil *Sitophilus oryzae* L. (Coleoptera: Curculionide) and red flour beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae) are very common pest infesting on many flours, cereals, meal, grains etc. They have world-wide distribution and are among the most economically important stored product pests (Aitken, 1975; Weston and Rattlingourd, 2000; Pugazhvendon et al., 2009). These stored grain insect pests have been damaging our economy by infesting agricultural stored products. According to an estimate these are responsible for worldwide loss of 10-40% in the stored-grain annually (Matthews, 1993).

For the control of these insect pests many synthetic insecticides and fumigants are used, but their widespread use led to some serious problems including development of insect resistance to insecticides (Zettler and Cuperus, 1990; Ribeiro et al., 2003), toxic residues in food, toxicity to consumers and increasing cost of application (Sighamony et al., 1990). The uncontrolled use of these synthetic pesticides caused great hazards for environment and consumers due to residual property (White, 1995). However, there is an urgent need to develop safe alternatives that are of low cost, convenient to use and environmentally friendly (Hassanali et al., 1990; Jember et al., 1995). The plants volatile essential oils extracted from aromatic plants have insecticidal properties could be considered as alternative insecticides (Shaaya et al., 1991; Shukla et al., 2008). Many plant volatile

essential oils and their constituents have been studied to possess potential as alternative compound and gaining tremendous importance for the management of stored products and these are ecologically safe and biodegradable (Rajendran and Sriranjini, 2008; Batish et al., 2008; Cosmi et al., 2009). The essential oils are volatile with high insecticidal efficiency and very low persistence. The active compounds present in essential oils are specific to particular insect groups (Huang et al., 1997) and not to mammals (Isman, 2000), many of them are not dangerous to human.

Eucalyptus globules (Mirtaceae) and *Ocimum basilicum* (Lamiaceae) are two important medicinal plants. The essential oils of different species of both plants leaves and their constituents have pesticidal properties against insect pests (Nagssoum et al., 2007). Therefore, the purpose of present work was to investigate the repellent properties of leaves essential oils of *Eucalyptus globules* and *Ocimum basilicum* against stored grain insect pests.

Material and Methods

Isolation of essential oils

Essential oils were extracted from leaves of *Eucalyptus globules* and *Ocimum basilicum*. The leaves were collected from the local area of Gorakhpur, district of Uttar Pradesh, India during March to June 2011. The leaves were dried in absence of sunlight at room temperature ($30 \pm 5^{\circ}\text{C}$) and grounded by domestic mixer. The dried powdered material was hydro-distilled in Clevenger

apparatus continuously for five hours to yield essential oils. The oils were collected in glass containers and kept in appendorff tubes at 5°C till their use.

Insect rearing

Red flour beetle *T. castaneum* and rice weevil *S. oryzae* were used to determine the repellent activity of essential oils. The insects were reared on wheat flour and grain in laboratory at 30±2°C, 75±5% RH and at photoperiod of 10:14 (L: D) hours.

Repellency

The repellency test used was adopted after (McDonald et al., 1970; Talukder and Howse, 1993, 1994). Four solutions of 0.05%, 0.10%, 0.20% and 0.40% of essential oils were prepared by dissolving essential oils in acetone. Whatman no. 1 filter papers were cut into two equal halves one half of each dish was treated with essential oil solution as uniform as possible by using micro pipette. The other half of the filter paper was treated with acetone only. The essential oil treated and acetone treated filter papers halves were dried to evaporate the solvent completely. Essential oil treated and acetone treated half-dishes were then attached lengthwise, edge-to-edge with adhesive tape and placed at the bottom in glass petri dish (height 15 mm × radius 45 mm). Ten adults of insects were released at the centre of the petri dishes and then petri dishes were covered and kept in dark. Six replicates were set for each concentration of essential oils. Number of the insects on both treated and untreated halves were recorded after four hours in mild light.

Data analysis

All the values obtained during observation were represented as mean ± SE. Chi-square test was applied to establish the repellent activity of the essential oils tested (Sokal and Rohlf, 1973).

Result

The repellent action of these two essential oils was studied against *T. castaneum* and *S. oryzae*. Data in (Table-1) showed that at highest concentration 0.40%, the essential oil of *Eucalyptus globules* had strong repellent action (9.16±0.30) and (8.66±0.33) against *T. castaneum* and *S. oryzae*. Similarly, at highest concentration 0.40% the *Ocimum basilicum* essential oil showed (8.50±0.22) and (8.16±0.30) against *T. castaneum* and *S. oryzae*, respectively. The repellent activity of *Eucalyptus globules* and *Ocimum basilicum* progressively increase with increase in concentration against both stored-grain insect pests. The chi-square test showed a significant (P<0.05) repellency of the both plants leaves against both

stored grain insect pests but *E. globules* had strong repellency in comparison to *O. basilicum* against *T. castaneum* and *S. oryzae*.

Discussion

Plant products especially in the form of essential oils having considerable potential as insecticidal compounds are gaining tremendous importance in recent years. In our present study the leaves essential oil of *Eucalyptus globules* and *Ocimum basilicum* was significantly (P<0.05) repellent at very low concentration against the flour beetle and rice weevil and join a series of some other essential oils which have similar insecticidal effects on the management of pests and provides a scientific relation for their use in control of stored-grain insect pests (Hassanali et al., 1990; Bekele et al., 1996; Bouda et al., 2001). The repellent effect of volatile essential oils may have an important implication in traditional post harvest storage system and have potential activity for their local availability make it attractive candidate in management of stored-grain insect pests.


Similar observation on the other plants extracts effect on several insect pests have been reported that the essential oils of *Schzygium aromaticum*, *Aegle marmelos*, *Corriandrum sativum* and *Citrus reticulata* extracted by a water distillation method showed strongly repellency against *S. oryzae* and *T. castaneum* even at low concentration but its repellency was more marked towards *S. oryzae* (Mishra and Tripathi, 2011). The essential oil of *Piper nigrum* (L.) repelled the adults of *T. castaneum* at low concentration (Upadhyay and Jaiswal, 2007). The plant volatile essential oils isolated from *Trachyspermum ammi*, *Anethum graveolens*, *Nigella sativa* show insecticidal activity against *T. castaneum* (Chaubey, 2007), while the leaf essential oil of *Melaleuca cajuputi* had different insecticidal activity against *Sitophilus zeamais* and *T. castaneum* but 100% repellency was only occurred in *T. castaneum* (Ko et al., 2009). The essential oil derived from Citrus peels possesses pesticidal activity against both insect pests *T. castaneum* and *S. oryzae* (Mishra et al., 2011). The effect of essential oil of *Ocimum grattissimum* leaves on *S. zeamais* was assessed for repellency and toxicity. The oil was found to be moderately repellent to the *S. zeamais* (Aswalam et al., 2008). Similarly, the ethyl acetate extract of *Citrullus colocynthis* showed highest repellent activity in lower concentration against *Callosobruchus maculates* (Prabhu Seenivasan et al., 2004).

Essential oils and their constituents of different *Eucalyptus species* (Mirtaceae) and *Ocimum species* (Lamiaceae) show many insecticidal activity against stored-grain insect pests (Keita et al., 2001; Jirovetz


et al., 2005; Brito et al., 2006; Ngassoum et al., 2007). The main component present in essential oils of *Eucalyptus species* are 1-8 cineole and in *Ocimum basilicum* are eugenol which showed different insecticidal properties against insect pests (Lee et al., 1997; Aswalam et al., 2008).

All the above findings clearly support the result of the present study. The mode of action of both essential oils is yet to be confirmed but the repellency of adults stored-grain insect pests may be due to the suffocation and inhibition of different biosynthetic processes of the insect metabolism (Don-Perdo, 1989).

Therefore, essential oils from leaves of *Eucalyptus globules* and *Ocimum basilicum* may be recommended as cheap, easily available at farmer level, eco-friendly with low mammalian toxicity and good alternative to synthetic insecticides. It could further reduce the application of the synthetic chemicals.



1,8 Cineole



Eugenol

Table 1: Repellency caused by *Eucalyptus globules* against adults of *Tribolium castaneum* and *Sitophilus oryzae* after four hours in filter paper test

Insect pests	Con. (%) vol: vol	Mean number of insects untreated±SE	Mean number of insects treated ±SE	χ^2 value P<0.05 (df=5)
<i>Tribolium castaneum</i>	0.05%	5.66±0.21	4.33±0.21	0.072 ^{NS}
	0.10%	7.33±0.33	2.66±0.33	1.058 ^S
	0.20%	8.00±0.25	2.00±0.25	1.800 ^S
	0.40%	9.16±0.30	0.83±0.30	3.461 ^S
<i>Sitophilus oryzae</i>	0.05%	5.16±0.30	4.83±0.30	0.005 ^{NS}
	0.10%	6.83±0.30	3.16±0.30	0.669 ^S
	0.20%	7.66±0.33	2.33±0.33	1.415 ^S
	0.40%	8.66±0.33	1.33±0.33	2.679 ^S

Adults of *T. castaneum* and *S. oryzae* were used in filter paper repellency assay. For each concentration of essential oils six replicate were carried out and 10 adults were used per replicate. Mean of untreated and treated halves in filterpaper repellency assay.

a. Not significant (NS) as the calculated values of χ^2 were less than the table values at probability levels (99%). Significant (S) at probability levels (99%).

Table 2: Repellency caused by *Ocimum basilicum* against adults of *Tribolium castaneum* and *Sitophilos oryzae* after four hours in filter paper test

Insect pests	Con. (%) vol: vol	Mean number of insects untreated ±SE	Mean number of insects treated ±SE	χ^2 value P<0.05 (df=5)
<i>Tribolium castaneum</i>	0.05%	5.33±0.21	4.66±0.21	0.021 ^{NS}
	0.10%	7.00±0.25	3.00±0.25	0.800 ^S
	0.20%	7.50±0.22	2.50±0.22	1.250 ^S
	0.40%	8.50±0.22	1.50±0.22	2.450 ^S
<i>Sitophilus oryzae</i>	0.05%	5.16±0.30	4.83±0.30	0.005 ^{NS}
	0.10%	6.83±0.30	3.16±0.30	0.669 ^S
	0.20%	7.16±0.40	2.83±0.40	0.933 ^S
	0.40%	8.16±0.30	1.83±0.30	1.997 ^S

Adults of *S. oryzae* and *T. castaneum* were used in filter paper repellency assay. For each concentration of essential oils six replicate were carried out and 10 adults were used per replicate. Mean of untreated and treated halves in filterpaper repellency assay.

a. Not significant (NS) as the calculated values of χ^2 were less than the table values at probability levels (99%). Significant (S) at probability levels (99%).




Figure 1. Repellent activity of the essential oil from *Eucalyptus globules* against 2 major stored-grain insect pests.




Figure 2. Repellent activity of the essential oil from *Ocimum basilicum* against 2 major stored-grain insect pests.

Acknowledgement

Authors are grateful to the authorities of Mahatma Gandhi P.G. College, Gorakhpur for providing lab facilities and University Grants Commission, New Delhi for providing financial assistance (U.G.C.No.35-65/2008[SR]) to carry out the investigation.

Correspondence to:

S.P. Tripathi
P.G. Department of Zoology, M.G.P.G. College,
Gorakhpur, 273 001, India
Telephone: 091-9335709553
Email- b2mishra123@gmail.com

Reference

- Aitken AD. Insect Travelers. I: Coleoptera .Techn.Bull., 31. H.M.S.O.London 1975:pp. 190.
- Asawalam EF, Emosairue SO, Hassanali A. Essential oil of *Ocimum grattisimum* (Labiatae) as *Sitophilous zeamails* (Coleoptera: Curculionidae) protectant. African Journal of Biotechnology. 2008;7: 3771-3776.
- Batish DR, Singh HP, Kohli RK, Kaur S. Eucalyptus essential oil as a natural pesticide. Forest Ecological Management. 2008; 256: 2166–2174.
- Bekele AJ, Obeng-Ofori D, Hassanali A Evaluation of *Ocimum suave* (Willd) as a source of repellent, toxicants and protectants in storage against three stored product insect pests. International Journal of Pest Management. 1996; 42(2): 139-142.
- Bouda H, Tapondjo A, Fotem DA, Gumedzoe MYD Effect of essential oils from leaves of *Ageratum conyzoides*, *Lantana Camara* and *Chromolaena odorata* on the mortality of *Sitophilus zeamais* (Coleoptera, Curculionidae). Journal of Stored Products Research. 2001; 37(2): 103-109.
- Brito JP, Baptistussi RC, Funichelo M, Oliveira JEM, Bortoli SA Effect of essential oils of *Eucalyptus* spp. under *Zabrotes subfasciatus* (Both., 1833) (Coleoptera: Bruchidae) and *Callosobruchus maculatus* (Fabr., 1775) (Coleoptera: Bruchidae) in two beans species. Bol Sanidad Veg Plagas. 2006; 32(4): 573-580.
- Chaubey MK Insecticidal activity of *Trachyspermum ammi* (Umbelliferae), *Anethum graveolens* (Umbelliferae) and *Nigella sativa* (Ranunculaceae) against stored-product beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). African Journal of Agriculture Research. 2007; 2(11): 596-600.
- Cosimi S, Rossi E, Cioni PL, Canale A Bioactivity and qualitative analysis of some essential oils from Mediterranean plants against stored-product pests: evaluation of repellency against *Sitophilus zeamais* Motschulsky, *Cryptolestes ferrugineus* (Stephens) and *Tenebrio molitor* (L.). Journal of Stored Products Research. 2009; 45(2): 125–132.
- Don-Perdo KN Mechanism of the action of the some vegetable oils against *Sitophilus zeamais* (Motsch) (Coleoptera: Curculionidae) on wheat. Journal of Stored Products Research. 1989; 25: 217-223.
- Hassanali A, Lwande W, Ole-Sitayo N, Moreka L, Nokoe S, Chapya A Weevil repellent constituents of *Ocimum suave* leaves and *Eugenia caryophylla* cloves used as grain protectant in parts of East Africa. Disc. Inno. 1990; 2: 91-95.
- Huang Y, Tin JM, Kini RM, Ho SH. Toxic and antifeedent action of nutring oil against *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch Journal of Stored Products Research. 1997; 35:289-298.
- Isman MB. Plant essential oils for pest and disease management. Crop Protection. 2000; 19: 603-608.
- Jembere B, Obeng-Ofori D, Hassanali A, Nyamasyo GNN. Products derived from the leaves of *Ocimum kilimanndscharium* (Labiatae) as post-harvest grain

- protectants against the infestation of three major stored product insect pests. Bulletin of Entomological Research. 1995: 85:361- 367.
14. Jirovetz L, Buchbauer G, Ngassoum MB, Ngamo LT, Adjoudji O. Combined investigation of the chemical composition of essential oils of *Ocimum gratissimum* and *Xylopia aethiopica* from Cameroon and their insecticidal activities against stored maize pest *Sitophilus zeamais*. *Ernährung*. 2005; 29(2): 55-60.
 15. Keita MS, Vincent C, Schmit J, Arnason JT, Belanger A. Efficacy of essential oil of *Ocimum basilicum* and *O. gratissimum* applied as an insecticidal fumigant and powder to control *Callosobruchus maculatus*. *Journal of Stored Products Research*. 2001; 37: 339-349.
 16. Ko K, Juntarajumong W, Chandrapatya A. Repellency, Fumigant and Contact Toxicities of *Litsea cubeba* (Lour.) Persoon against *Sitophilus zeamais* Motschulsky and *Tribolium castaneum* Herbst. *Kasetsart Journal (Natural Science)*. 2009; 43(1):56-63.
 17. Lee S, Tsao R, Peterson C, Coats JR. Insecticidal activity of monoterpenoids to western corn root worm (Coleoptera: Chrysomelidae), two spotted spider mite (Acari: Tetranychidae) and Housefly (Diptera: Muscidae). *Journal of Economic Entomology*. 1997; 90: 883-892.
 18. Matthews GA. Insecticide application in the stores. In Matthews, G.A. and Hislop, E.C. (eds.). *Application technology for crop protection*. CAB, London. 1993: pp. 305-315.
 19. McDonald LL, Guy RH, Speirs RD. Preliminary evaluation of new candidate materials as toxicants, repellents and attractants against stored-product insects. Agricultural Research Service, US Department of Agriculture, Washington DC. 1970: Marketing Research Report No. 882.
 20. Mishra BB, Tripathi SP. Repellent activity of plant derived essential oils against *Sitophilus oryzae* (Linnaeus) and *Tribolium castaneum* (Herbst). *Singapore Journal of Scientific Research*. 2011; 1 (2):173-178.
 21. Mishra BB, Tripathi SP, Tripathi CPM. Contact Toxicity of essential oil of *Citrus reticulata* fruits peels against stored grain pests *Sitophilus oryzae* (Linnaeus) and *Tribolium castaneum* (Herbst). *World Journal of Zoology*. 2011; 6 (3):307-311.
 22. Ngassoum MB, Tinkeu LSN, Ngatanko L, Taponjou LA, Lognay G, Malaisse F, Hance T. Chemical composition, insecticidal effect and repellent activity of essential oils of three aromatic plants, alone and in combination, towards *Sitophilus oryzae* L. (Coleoptera: Curculionidae). *Nat. Prod. Comm*. 2007; 2(12): 1229-1232.
 23. Prabhu Seenivasan S, Jayakumar M, Raja N, Ignacimuthu S. Efficacy of bitter apple (*Citrullus colocynthis*) seed extracts against pulse beetle, *Callosobruchus maculatus* Fab. (Coleoptera: Bruchidae). *Entomon*. 2004; 29(1):81-84.
 24. Pugazhvendon SR, Elumalai K, Ronald Ross P, Sundararajan M. Repellent activity of chosen plant species against *Tribolium castaneum*. *World Journal of Zoology*. 2009; 4(3): 188-190.
 25. Rajendran S, Sriranjini V. Plant products as fumigants for stored product insect control. *Journal of Stored Products Research*. 2008; 43(2): 126-135.
 26. Ribeiro BM, Guedes RNC, Olivira EE, Santos JP. Insecticidal resistance and synergism in Brazilian populations of *Sitophilus zeamais* (Coleoptera: Curculionidae). *Journal of Stored Products Research*. 2003; 39:21-31.
 27. Shaaya E, Ravid U, Paster N, Juven B, Zisman U, Pistarev V. Fumigant toxicity of essential oils against four major stored product insects. *Journal of Chemical Ecology* 1991; 17: 499-504.
 28. Shukla J, Tripathi SP, Chaubey MK. Toxicity of *Myristica fragrans* and *Illicium verum* essential oils against flour-beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *Electronic Journal of Environmental Agricultural and Food Chemistry*. 2008; 7(7): 3059-3064.
 29. Sighamony S, Anees I, Chandrakala T, Kaiser Jamil S. Indigenous plant products as grain protectants against *Sitophilus Oryzae* (L) and *Rhyzopertha dominica* (F). *Journal of Stored Products Research*. 1990; 22: 21-23.
 30. Sokal RR, Rohlf FJ. *Introduction to biostatistics*. Freeman WH, San Francisco. 1973: pp. 165, 231, 289.
 31. Talukder FA, Howse PE. Deterrent and insecticidal effects of extracts of pithraj, *Aphanamixis polystachya* (Meliaceae), against *Tribolium castaneum* in storage. *Journal of Chemical Ecology*. 1993; 19: 2463-2471.
 32. Talukder FA, Howse PE. Efficacy of pithraj (*Aphanamixis polystachya*) seed extracts against stored-product pests. *Proc. Int. Working Conf. on Stored-product Protection*. 1994; 2: 848-852.
 33. Upadhyay RK, Jaiswal G. Evaluation of biological activities of *Piper nigrum* oil against *Tribolium castaneum*. *Bulletin of Insectology*. 2007; 60(1):57-61.
 34. Weston PA, Rattlingourd PA. Progeny production by *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Oryzaephilus surinamensis* (Coleoptera: Silvanidae) on maize previously infested by *Sitotroga cerealella* (Lepidoptera: Gelechiidae). *Journal of Economic Entomology*. 2000; 93: 533-5.
 35. White L. *Chemical control. Integrated management of insects in stored products*. Dekker, Inc; New York. Basel. Hong Kong. 1995: pp. 287- 330.
 36. Zettler JL, Cuperus GW. Pesticide resistance in *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Rhyzopertha dominica* (Coleoptera: Bostrichidae) in wheat. *Journal of Economic Entomology*. 1990; 83: 1677- 1681.