



## Urban church forests for urban temperature regulation: Implications the role of managing and incorporating urban green area in urban planning

Tulu Tura

Mr. Tulu Tolla Tura PhD student at Centre for Environment Science, College of life science, Addis Ababa University, Ethiopia.

Po. Box 1176

Phone: [+251-0911907038](tel:+251-0911907038)

[tulutollatura@gmail.com](mailto:tulutollatura@gmail.com)

Addis Ababa, Ethiopia

**Abstract:** The global surface temperature shows an increment of  $0.5 \pm 0.1^\circ\text{C}$  per decade and  $1.05 \pm 0.3^\circ\text{C}$  per century from 1880-2014 with greater increases in cities than non-urban areas. Global communities are shifting towards urbanization due to various factors. Urbanization has caused lack of stable condition for dwellers due to environmental and anthropogenic factors such as land cover changes. Urban temperature rising is the main factors hindering urban dwellers at global level due to insufficient green areas. Social institutions are playing important role in urban greening and urban climate regulation. Ethiopian Orthodox Tewahido Church has long history in biodiversity conservation and greening role in urban and rural parts of the country. However, there is a research gap in Ethiopia regarding the role of urban green area in the church yards in urban temperature regulation. Therefore, the study evaluated the role of church managed forests in Addis Ababa in regulating surface temperature. Surface temperature inside four church forest at a buffer radius of 0-50m, 50-100m, 100-200m and 200-500m estimated using Landsat image thermal band 6 of 1986, 2000 and 2010 and ground measurement by ambient thermometer at 10:00am, 12:30am and 3:00pm local time. The ground measurement was done in order to validate satellite image analysis. Plant species diversity, DBH, H, HC, BH and BA was measured. There were 1167 trees in the four studied churches. The mean temperatures of the studied sites were  $22.5 \pm 0.1$ ,  $23.25 \pm 0.2$ ,  $24 \pm 0.6$ ,  $24.6 \pm 1.1$  and  $25.5 \pm 2.2^\circ\text{C}$  on site, 0-50m, 50-100m, 100-200m and 200-500m respectively for 1986 images;  $23.2 \pm 0.5$ ,  $23.3 \pm 1.0$ ,  $24.3 \pm 2.1$ ,  $24.8 \pm 2.2$  and  $25.5 \pm 1.8^\circ\text{C}$  on site, 0-50m, 50-100m, 100-200m and 200-500m respectively for 2000 images and  $23.2 \pm 0.3$ ,  $23.27 \pm 0.2$ ,  $23.7 \pm 1.6$ ,  $24 \pm 1.4$  and  $24.7 \pm 1.3^\circ\text{C}$  on site, 0-50m, 50-100m, 100-200m and 200-500m respectively for 2010 images. The results illustrated a significant influence of green area on urban temperature in the buffering radius and imply the possibility of regulating urban temperature by planning urban green area in appropriate radius interval. Therefore, the study indicated that church forests in particular, social institution and urban green area in general have significant role in urban temperature regulation.

[Tulu Tura. **Urban church forests for urban temperature regulation: Implications the role of managing and incorporating urban green area in urban planning.** *Nat Sci* 2023,23(6):35-50]. ISSN 1545-0740 (print); ISSN 2375-7167(online). <http://www.sciencepub.net/nature> 06.doi:[10.7537/marsnsj210623.06](https://doi.org/10.7537/marsnsj210623.06).

Key words: Urban green area, Church forest, urban temperature, green area buffer, and urban planning

### 1. Introduction

The global surface temperature is increasing by  $0.5 \pm 0.1^\circ\text{C}$  per decade and  $1.05 \pm 0.3^\circ\text{C}$  per century from 1880-2014 on average (NOAA, 2014) with greater increases in cities than any terrestrial parts (Ackerman, 1985; Chandler, 1962; Oke, 1973; Moreno Garc'ia, 1994). Cities and other built up areas natural climatic condition is altered due to changes in land use land cover and other anthropogenic activities (Krayenhoff and Voogt, 2010). These alterations have an effect on the absorption of solar radiation, surface temperature, evaporation rates, storage of heat, wind turbulence, modification in the near atmosphere over cities and causes temperature difference between urban and rural

settings (Mitchell, 1961; Mallick J. *et al.*, 2008). The radical changing of urban land is increasing through time due to global community shifting towards urbanization (Smith *et al.*, 2009). Around half of the world's human population lives in urban areas and anticipation of urbanization will increased by 70% of the present world urban population by 2030 (Smith *et al.*, 2009).

Urban heat islands impact urban residents in several ways, including modify energy demand (higher in summer and lower in winter), causes thermal stress on inhabitants, and increase air pollution formation rates (Oke, 1997). Urban temperature is a concern because it causes health hazard such as heat syncope or fainting;

heat edema or swelling of dependent parts such as the legs; and heat tetanias as a result of heat induced hyperventilation with other effects include heat cramps, heat exhaustion vomiting, weakness, and mental status changes (Frumkin, 2002). The most acute heat related conditions is heat stroke (Frumkin, 2002). Generally, temperature in excess of 23 °C is associated with heat related stress and causes increase in summer deaths (WHO, 2008).

Land use planning and building strategies to regulate temperature within cities is influence the wellbeing as well as the environmental impact of heating and cooling of urban dwellers (Taha *et al.*, 1988; Saito, 1990; Saito, 1991). Therefore, studying urban green area role for local climate regulation is an important aspect in order to create sustainable and health ecosystem for the dwellers (Tratalos *et al.*, 2007;

Whitford *et al.*, 2001). Particularly, in the limited urban land social institutions role on urban greening is potential option. However, there is no published accessible research on urban green area in general social institution role in particular of urban temperature regulation in Ethiopia.

## 2. Materials and Methods

### 2.1. Description of Study Site

The study area, Addis Ababa City, is located in the central part of Ethiopia. Addis Ababa has an elevation between 3023 and 2054 meter above sea level at geographical location of 09°05' N to 09°00' N latitudes and 38°45' E to 38°42' E longitude. The city receives annual mean rainfall of 1128 mm and mean monthly temperature of 16.25 °C. Addis Ababa city is center of political and diplomatic mission with population of 3.5 million (CSA, 2007).




Fig 1: Location map of Addis Ababa and studied churches

### 2.2. Sampling Design

Sampling was designed based on elevation of the churches under investigation in order to control biasness due to temperature change along elevation

variation. Four churches were selected in the same elevation range of varying forest area coverage and tree species.

Table 1: Elevation, area and location of the studied church

Selected church	Elevation	Area	Location
Menbera Tsebaoti Kidist Selassie	2468	1.815	09 <sup>0</sup> 01 N and 38 <sup>0</sup> 46E
Manbera Mengisit Kidisit Mariam	2470	0.2	0998822N and 047375 E
Miskaye Hizunan Medhanialem	2510	0.5	1000252N and 0473843E
Geneta Tsige Kidus Giorgis	2473	0.765	09 <sup>0</sup> 02N and 38 <sup>0</sup> 45E

### 2.3. Species Identification

In order to investigate species diversity and growth impact on urban temperature regulation; the living trees in the studied churches were identified and recorded. Plant identification was done in the field using Flora of Ethiopia and Eritrea volume 2, part 1 and part 2 (Edwards *et al.*, 1995 and 2000); Flora of Ethiopia volume 3 (Hedberg and Edwards, 1989). For those species difficult to identify in the field, fresh specimens were collected and then pressed properly for identification at National Herbarium of Addis Ababa University.

### 2.4. Temperature Measurement and Land Cover Change Analysis

Primary data were generated from extraction of satellite image and ground measurement. Ground field data collection was conducted inside the church compound and buffering 50m, 100m, 200m and 500m from the existing green spaces at 10:00 am, 12:30 am and 3:00 pm local time using ambient thermometer in order to validate thermal band satellite image.

Satellite imageries and ancillary data were collected in order to identify historical and recent land cover change. Land cover change analysis helps as to generate normalized difference in vegetation index and estimating surface temperature. Garmin GPS was used to verify the data generated from satellite image with ground. ERIDAS imagine 2010; ArcGIS 10.2, Global Mapper 11, 3DEM and Easy GPS software was used for image processing and mapping purpose.

The image data was obtained from United State Geological Survey of Landsat thermal Band 6 1986, 2000 and 2010. The images has been acquired at the end of December and beginning of January between 10:00-11:00 am which is dry season in Ethiopia and high acquisition period with less environmental intrusion.

The images were selected in order to represent urban policy impact on urban green area and land cover change during the consecutive three political regime;

1986 Derg regime, 2000 Ethiopia People Republic Democratic Front (EPRDF) more focus on rural development and 2010 EPRDF focus on rural and urban development.

However, Land Sat 2014 image was not used due to the band width was smaller than that of Landsat TM and ETM<sup>+</sup> images of 1986 and 2000 which may cause overstatement in normalized difference vegetation index and temperature estimation. In the meantime, study area visualization has been done using Ikonos 2006 and Google earth image at various times for detail analysis.

Digital elevation model (DEM) of the study area has been developed using 20 m horizontal and 16 m vertical absolute precision Shuttle Radar Topography Mission (SRTM) of NASA satellite image with 30 meter resolution.

### 2.5. Image Precision Test and Validation

Band selection carried out toward the most precision information from remotely sensed data via analysis of reflectance properties of objects or features. Consequently, for the purpose of attaining accurate and validated image processing; Landsat images through ERDAS IMAGINE 2011 software, spectral reflectance curves and histogram behavior of the bands of satellite images were integrated an analysis of detailed feature records on the ground at the time of data acquisition and different vegetation types in the sites. Lillesand and Kiefer (1994) suggested that image interpreters should have good power of observations, imagination and careful understanding of the phenomenon being studied as well as knowledge of the geographic region under study. As a result, digital image enhancement and interpretation techniques were used.

True Color Composite and False Color Composite were produced to increase the visual interpretability of the images and the amount of information that can be visually interpreted from the data. Digital image enhancement techniques such as contrast stretching, band ratios and normalized difference vegetation index

analysis were done. Finally, image analyses result were validated with ground measurement.

**2.6. Data Analysis**

Thematic mapper and enhanced thematic mapper data process has been done through plunging ArcGIS remote sensing tool boxes in Landsat Thermal Band 6. The tool helps to estimate surface temperature of a given area and converts pixel values of thermal bands at spectral sensor radiance by conversion coefficient (Walaweder *et al.*, 2012).

*Step1. Conversion of Digital Number to Spectral Radiance (L<sub>s</sub>)*

$$L_s = Gain * Digital\ Number + Bias$$

Where

$$L_s = Spectral\ Radiance$$

$$Gain\ and\ Bias = Conversion$$

*coefficient*

Inverted plank's law and specific calibration constant (K1 and K2) transforms spectral radiance to sensor brightness temperature (Walaweder *et al.*, 2012)

*Step2. Conversion of Spectral Radiance to Temperature in Kelvin*

$$T_B = \frac{K_2}{\ln\left(\frac{K_1}{L_s} + 1\right)}$$

Where

*K1= Calibration Constant 1*

*K2 = Calibration Constant 2*

*TB = Surface Temperature in degree Kelvin*

And

Temperature in degree Kelvin (°K) converted to degree Celsius (°C) (Walaweder *et al.*, 2012)

*Step3. Conversion of Kelvin to Celsius*

$$TC = TB - 273$$

Where

*TB= Surface Temperature in degree Kelvin*

*TC= Surface Temperature in degree Celsius*

*273= Conversion factor from Kelvin to Celsius*

Basal area of trees in the study sites were calculated with the formula:

*BA = 0.785 DBH<sup>2</sup>, Where, DBH is the diameter of trees at breast height and 0.785 is conversion factors*

When the image processing completed the digital model images were displayed in the Arc Map software with applied color ramp scheme. Statistical software (R) was applied to analysis mean, standard deviation and correlation between temperature and different parameters considered in the study.

**3. Results and Discussion**

**3.1. Elevation and Land Use Classification**

The elevation classification of Addis Ababa city at 100 m interval was categorized in eleven (11) classes with minimum 2054m and maximum 3023m above sea level. This elevation difference with other factors may induce a large temperature variation in the city. However, there is no significant temperature difference due to elevation in the selected studied sites. Because all the churches under studies were found in 2401m - 2500 m above sea level elevation range.




Fig 2: Elevation Classification of Addis Ababa City

Land cover classification of the sites categorized with vegetation and without vegetation. The land cover model illustrated some parts of the city have better green area, some have partial green area and the others have completely urban downtown fig 3.




Fig 3: Digital Model of Land Cover in the study sites

The green color indicates green area and red color indicates urban downtown. The yellowish color is the jumble of sparse green area, building and other urban infrastructure. The appearance of varies color in the land cover digital model is due to reflectance of receiving objects during satellite image acquisition.

**3.2. Plant Species and Growth Distinctiveness of Study Sites**

There were 1167 trees in the study sites. The numbers of indigenous tree species were higher than exotic species in all the study sites. However, the numbers of exotic tree species were higher at *Menbere Tsebaot Kidist Silasse* than other study sites. The areas of green

parts in the studied church compound were directly related with species density. As green area increase species density linearly increase. The number of exotic tree species increases as area of green space increase table 2. Increasing green area size is directly influence the radiance of temperature regulation (Jauregui, 1990 and 1991).

Table 2: Type and number of plant species in the study sites

Study sites	Area	Species				Total density (n <sub>Q</sub> )
		Indigenous		Exotic		
		Species (n <sub>Q</sub> )	Density (no)	Species (n <sub>Q</sub> )	Density (n <sub>Q</sub> )	
Menbere Tsebaot Kidist Silasse	1.8	9	355	7	58	442
Manbera Mengisit Kidisit Mariam	0.2	4	78	1	2	80
Miskaye Hizunan Medhanialem	0.5	8	280	4	25	305
Genete Tsige Kidus Giyorgis	0.7	6	305	5	35	340
<b>Total plant species</b>						<b>1167</b>

The influence of green area on urban temperature regulation is mainly based on the size of green area, types of plant species, diversity of plant species and growth stages of plants in the green area. It is directly related with absorption and transmission of a certain percentage of the incoming solar radiation, Evapotranspiration (movement of water to the air from different surfaces such as the soil, canopy interception, and water, and conversion of water within the leaf to water vapor which is then released to the atmosphere through the stomata); thus cooling the leaf and the surrounding local microclimate (Kotzen, 2003;

Nakaohkubo and Hoyano, 2011; Hunter Block *et al.*, 2012; Berry *et al.*, 2013). This mechanism induces an important role in the water cycle, and contributes to the provision of ecosystem services by trees (Georgi and Dimitriou, 2010).

Trees produce oxygen, intercept airborne particulates, and reduce smog, enhancing a community's respiratory health and canopy directly contributes to meeting a city's regulatory clean air requirements (Emmanuel, 2005). This function is strongly influenced by type of species, diversity and growth stage of trees.

Table 3: Growth characteristics of plants in the study sites

Study sites	H (m)	DBH(cm)	HC (m)	BH (m)	Basal Area
<i>Menbere Tsebaot Kidist Silasse</i>	32.6 ± 5.12	58.3 ± 14.2	15.4 ± 12.4	7.4 ± 6.5	0.27 ± 0.01
<i>Manbera Mengisit Kidisit Mariam</i>	22.2 ± 3.3	47.3 ± 10.3	17.4 ± 7.6	4.7 ± 5.6	0.18 ± 0.02
<i>Miskaye Hizunan Medhanialem</i>	20.7 ± 6.7	35.7 ± 25.2	12.3 ± 12.5	8.2 ± 5.8	0.10 ± 0.05
<i>Genete Tsige Kidus Giorgis</i>	23.8 ± 4.5	60.2 ± 12.7	20.3 ± 19.7	5.6 ± 5.8	0.28 ± 0.01

Trees in the study sites have a range of growth stage. The mean height of trees in each sites were 32.6±5.12m, 22.2±3.3m, 20.7±6.7m and 23.8±4.5m for *Menbere Tsebaot Kidist Silasse*, *Manbera Mengisit Kidisit Mariam*, *Miskaye Hizunan Medhanialem* and *Genete Tsige Kidus Giorgis* respectively. Trees height at *Miskaye Hizunan Medhanialem* church have highest standard deviation and *Manbera Mengisit Kidisit Mariam* church have lowest standard deviation than other study sites. This height variation indicates growth difference and diversity of species. Diameters at breast height (DBH) of trees in the study sites were illustrated analogous trends of height. Crown height (HC), basal

height (BH) and basal area of trees in the study sites were designated irregular standard deviation table 3.

### 3.3. Temperature Influence Range of Studied Churches Forest

The digital thermal model of the study sites looks like similar manifestation in visual. However, the surround environment is not similarly for the entire study site. Some of the study sites were interfaced with green area and the other urban downtown. The interfacing environment of the study sites and buffering zone has impact on the green area temperature regulation range (Taha *et al.*, 1988; Saito, 1990; Saito, 1991).




Fig 3: Digital Thermal Model 2010 Image of the Study Site

The mean surface temperature of 2010 Landsat Thermal Band 6 image of entire study sites illustrated 34.99 °C highest and 13.44 °C lowest respectively fig 3. The temperature variation designated the difference between green and non green area entire study site with other environmental and anthropogenic factors. The green area appears green color in the digital thermal model with lowest temperature record and the red color

indicates devoid of vegetation with highest temperature record.

The mean maximum and minimum surface temperature of 2010 Landsat Thermal Band 6 image on sites and buffering 0-50m, 50-100m, 100-200m and 200-500m was 30.6 °C and 17.2 °C respectively. There was significant difference between temperatures of entire city and buffering zone of studied sites fig 4.




Fig 4: Digital Thermal Model of 2010 image onsite and 0-500m buffering zone of study sites

The mean surface temperature of 2000 landsat thermal band 6 image of entire study sites illustrated 42°C highest and 13.5°C lowest respectively fig 5. The temperature variation designated the difference between green and non green area on the entire study sites. The green area appears green color in the digital thermal model with lowest temperature record and the red color indicates devoid of vegetation with highest temperature record. The lowest temperature recorded at

urban covered with green area and the highest temperature is observed in downtown.

The highest mean temperature value is significantly differing from other regimes (2010 and 1986) considered satellite images. This might be due to varies factors related with image acquisition process which causes distinct values overall. However, lowest mean temperature was not significant differing from 2010 and 1986 images.




Fig 5: Digital Thermal Model 2000 image of the Study Sites

The mean maximum and minimum surface temperature of 2000 Landsat Thermal Band 6 image on site and buffering 0-50m, 50-100m, 100-200m and 200-500m is 32.6 °C and 16.1 °C respectively. There was significant

temperature difference between 2010 and 2000 image analysis both in lowest and highest point fig 6. But the temperature difference in the buffering zone was not significant for 2010 and 2000 images.




Fig 6: Digital Thermal Model of 2000 Image of Onsite and 0-500m Buffering Zone of Study Sites



The mean surface temperature of 1986 landsat thermal band 6 image of entire study sites illustrated 36.6 °C highest and 14.4 °C lowest respectively fig 7. The temperature variation designated the difference between green and non green area entire study site. The green area appears green color in the digital thermal model with lowest temperature record and the red color

indicates devoid of vegetation with highest temperature record.

The highest temperature acquired in 1986 image was greater than 2010 and lower than 2000. But the lowest temperature acquired was almost the same except very slight decimal point difference.




Fig 7: Digital Thermal Model 1986 Landsat Image of the study site

The mean maximum and minimum surface temperature of 1986 Landsat Thermal Band 6 image on site and buffering 0-50m, 50-100m, 100-200m and 200-500m is

29 °C and 17.6 °C respectively. There was significant difference between temperatures of enter city and the buffering zone of studied sites fig 8.




Fig 8: Digital Thermal Model of 1986 image of onsite and 0-500m buffering zone of the study sites

The mean entire city green area temperature ranges in the three political regimes were not regular trend. Even if the three political regimes were different police and strategies regarding urban, its impacts on studied churches forest were no significant. This indicates

social institutions not significantly influenced by policy and strategy of political philosophy followed when compared with the rest urban green area table 4 and fig 9.

Table 4: Mean surface temperature in °C inside churches compound and buffering zone of three political regimes

Year	Onsite	0-50	50-100	100-200	200-500
1986	22.5 ± 0.1	23.25 ± 0.2	24 ± 0.6	24.6 ± 1.1	25.5 ± 2.2
2000	23.2 ± 0.5	23.37 ± 1.0	24.3 ± 2.1	24.8 ± 2.2	25.5 ± 1.8
2010	23.2 ± 0.3	23.27 ± 0.2	23.7 ± 1.6	24 ± 1.4	24.7 ± 1.3




Fig 9: Green area temperature influence range in the three political regimes

The mean temperature in each studied churches compound and buffering zone in the three political regimes were not uniform. *Menbera Tsebaot Kidisit Selassie* illustrates the lowest ( $19.3 \pm 0.01$  °C) in 1986 and highest ( $21.7 \pm 0.2$  °C) in 2010; *Menbera Mengisit Kidisit Mariam* lowest ( $24 \pm 0.4$  °C) in 2010 and highest ( $24.5 \pm 0.8$ °C) in 2000; *Miskaye Hizunan*

*Medhanialem* lowest ( $22.8 \pm 0.1$ °C) in 1986 and highest ( $24.2 \pm 0.3$  °C) in 2010 and *Geneta Tsige Kidus Giorgis* lowest ( $23 \pm 0.3$ °C) in 2010 and highest in ( $24 \pm 0.6$  °C) in 2000 table 5 and fig 10. This illustrates green area size; type of tree species and growth stages of trees affect the temperature regulation range of green area in the three political regimes.

Table 5: Mean surface temperature in °C of each studied church compounds of three political regimes

Selected church	1986	2000	2010
Menbera Tsebaot Kidisit Selassie	19.3 ± 0.01	21 ± 0.2	21.7 ± 0.2
Menbera Mengisit Kidisit Mariam	24.2 ± 0.2	24.5 ± 0.8	24 ± 0.4
Miskaye Hizunan Medhanialem	22.8 ± 0.1	23.4 ± 0.4	24.2 ± 0.3
Geneta Tsige Kidus Giorgis	23.5 ± 0.1	24 ± 0.6	23 ± 0.3




Fig 10: Mean surface temperature in °C of each studied church compounds of three political regimes

The mean temperature recorded for Menbera Tsebaot Kidisit Silase and Geneta Tsige Kidus Giorgis was linearly increases as distance from green area increases. However, the mean temperature of Miskaye Hizunan

Medhanialem and Menbera Mengisit Kidisit Mariam was irregular it might be due to the neighboring environment in the buffering zone of study sites table 6 and fig 11.

Table 6: Mean surface temperature in °C inside church compound and buffering zone of ground measurement using ambient thermometer

Studied Churches	Site	0-50 m	50-100 m	100-200m	200-500m
Menbera Tsebaot Kidisit Selassie	20.50 ± 0.2	21.80 ± 0.2	22.70 ± 0.5	24.20 ± 1.2	25.3 ± 2.1
Menbera Mengisit Kidisit Mariam	21.30 ± 0.4	22.20 ± 1.1	22.50 ± 2.0	25.50 ± 2.3	25.2 ± 1.6
Miskaye Hizunan Medhanialem	20.7 ± 0.3	22.00 ± 0.3	22.90 ± 1.3	25.70 ± 1.4	24.8 ± 1.3
Geneta Tsige Kidus Giorgis	20.80 ± 0.3	22.10 ± 0.1	23.00 ± 0.4	25.90 ± 1.1	25.3 ± 2.3




Fig 11: Mean surface temperature in °C inside church compound and buffering zone of ground measurement using ambient thermometer

The mean surface temperature recorded in side studied churches compound and buffering zone of 0-50 m, 50-100m, 100-200m and 200-500m landsat thermal band 6 images 2010, 2000 and 1986 was increase as distance from green area increases. The mean surface

temperature increment was not uniform for all images in the three political regimes. This indicates the neighboring land cover change in the three political regimes buffering of study sites table 7.

Table 7: Mean surface temperature in °C with standard deviation of buffering zone in the three political regimes

Church	Year	0-50		50-100		100-200		200-500	
		Mean	St.dev	Mean	St.dev	Mean	St.dev	Mean	St.dev
Menbera Tsebaot Kidisit Selassie	2010	22	0.2	22.5	0.6	23.4	1	25	1.5
	2000	23.7	1	23.3	2.3	23.6	2.3	25	2
	1986	20	0.3	23.8	0.5	22.2	1.2	24.4	1.3
Menbera Mengisit Kidisit Mariam	2010	25	0	25.3	0.7	25.7	1	26	1
	2000	26.2	0.35	26.2	1.3	26.6	1.6	26	1.5
	1986	25	0	25.2	0.4	25	0.6	24.6	1
Miskaye Hizunan Medhanialem	2010	24.6	0	24.4	4	24.4	1.6	24.5	1.5
	2000	23	1	24.7	1.8	25	2	25	2
	1986	23.4	0	21	0.6	24.2	1.2	24	1.3
Geneta Tsige Kidus Giyorgis	2010	23.6	0.6	23.7	1	25.3	1.4	27	1.3
	2000	25.2	2.5	25.6	2.8	26.5	3	27.8	1.6
	1986	23.9	0.6	23.3	0.3	24.5	1.3	26	1.3

The correlation of temperature with area was strong negative which indicate strong relation. As green area increases temperature decrease in all buffering zone.

Area and total trees in the study sites have positive relationship. As area increase numbers of total, indigenous and exotic trees are increases table 8.

Table 8: Correlation of temperature in °C with area, tree diversity and total species

	Area	Onsite	0 - 50	50 - 100	100 - 200	200 - 500	Total Trees	Indigenous	
Exotic									
Area	1.00	-0.80	-0.94	-0.70	-0.91	-0.24	0.84	0.78	0.90
On site	-0.80	1.00	0.93	0.98	0.86	0.29	-0.98	-0.96	-0.95
0 - 50	-0.94	0.93	1.00	0.89	0.98	0.42	-0.91	-0.94	-0.93
50 - 100	-0.70	0.98	0.89	1.00	0.84	0.42	-0.92	-0.98	-0.87
100 - 200	-0.91	0.86	0.98	0.84	1.00	0.60	-0.81	-0.93	-0.82
200 - 500	-0.24	0.29	0.43	0.42	0.60	1.00	-0.11	-0.54	-0.10
Total Trees	0.84	-0.98	-0.91	-0.92	-0.81	-0.10	1.00	0.89	0.99
Indigenous	0.78	-0.96	-0.94	-0.98	-0.93	-0.54	0.89	1.00	0.86
Exotic	0.90	-0.95	-0.93	0.87	-0.82	-0.08	0.99	0.86	1.00

The correlation of height with DBH, HC, BH and BA was positive. It was a linear relationship as height increases DBH, HC, BH and BA are increase. DHB was negatively correlated with BH and HC. It indicates weak correlation of DHB with BH and HC in the sites. The correlation of temperature onsite, 0-50, 50-100, 100-200 and 200-500 was negatively correlated with H, DBH, BH and BA except 200-500 with BA and DBH. This indicates the relationship of temperature with H, DBH, BH and BA was indirect strong relationship except 200-500 with BA and DBH. The correlation of

temperature with crown height was positive which indicates as crow height increase temperature also increases. This was indirect negative relationship of crown height on temperature regulation. Trees having large crown height regulate less surface temperatures than short crown height table 9. The result illustrated strong relationship between species diversity, area of forest coverage and growth stages of plants temperature of urban. Therefore, urban green area role for urban temperature regulation was clearly visualized.

Table 9: Correlation of temperature in °C with height (H), diameter at breast height (DBH), crown height (HC), basal height (BH) and basal area (BA)

	On.site	0-50	50-100	100-200	200-500	H	DBH	HC		
BH										
BA										
On.site	1.00	0.93	0.98	0.86	0.29	-0.59	-0.21	0.35	-0.82	-0.25
0-50	0.93	1.00	0.89	0.98	0.43	-0.82	-0.30	0.37	-0.73	-0.33
50-100	0.98	0.89	1.00	0.84	0.42	-0.48	-0.01	0.51	-0.91	-0.05
100-200	0.86	0.98	0.84	1.00	0.59	-0.79	-0.15	0.52	-0.76	-0.19
200-500	0.29	0.43	0.42	0.59	1.00	-0.16	0.63	0.95	-0.70	0.61
H	-0.60	-0.81	-0.48	-0.79	-0.16	1.00	0.66	0.05	0.22	0.68
DBH	-0.21	-0.29	-0.01	-0.15	0.63	0.66	1.00	0.77	-0.39	0.99
HC	0.35	0.37	0.51	0.51	0.95	0.05	0.77	1.00	-0.81	0.74
BH	-0.82	-0.73	-0.91	-0.76	-0.70	0.22	-0.39	-0.81	1.00	-0.35
BA	-0.25	-0.33	-0.05	-0.19	0.60	0.68	1.0	0.74	-0.35	1.00

The result of this study coincides with studies conducted at different countries on different climatic zones. Spatial scales of cooling the surface temperature within a green space may be 15–20 °C lower than that of the surrounding urban area giving rise to 2–8 °C cooler air temperatures and a cooling effect that extends out in to the surrounding area (Taha *et al.*, 1988; Saito, 1990; Saito, 1991). The mean near surface air temperature in the shade was less 0.8-1.7°C than the ambient air temperature (min 0.4, max. 4.5°C) depending on the background climate, measurement method and type of surface cover when measured at 25-35°C air temperature (Hodder and Parson, 2007; Thorsson *et al.*, 2007; Kántor *et al.*, 2013).

Susca *et al.*(2011) finding show that, New York City on average, those areas with less vegetation were hotter than those with more vegetation, the difference between the hottest and the coolest areas around 2°C. Chapultepec Park in Mexico City of 500 ha size was reduced temperature by 1.7°C in the radius of 2 km (Jauregui, 1990–91). A leeward breeze park of 60 ha size was reduced temperature by 1.5°C in the radius of 1 km (Asaeda and Abu, 1998). Benjamin Park in Haifa, Israel of 0.5 ha size was reduced temperature by 1.5°C in the radius of 150m (Givoni, 1998). Green space Kumamoto City, Japan of 0.24 ha size was reduced temperature by 1–2°C in the radius of 20 m (Saito, 1990 and 1991). Honjo and Takakura (1990 and 1991) suggested that a 100 m wide green space cools to a distance of 300 m, 400 m wide green space cools to a distance of 400 m and recommended that green spaces should be no more than 300 m apart for optimum cooling within a neighborhood.

In addition to the role of green space size, the extent of the cooled area around a green space is influenced by the type and composition of vegetation in the green space. Potchter *et al.*(2006) found that high and wide canopy trees and higher tree/shrub coverage resulted in cooler parks compared to the surroundings. Similarly, Yu and Hien (2006) reported that the ambient temperature in a park was strongly correlated to the density of plants. Chang *et al.* (2007) showed that paved surfaces were equal or more than 50% of the land coverage parks were on average warmer than their surroundings. The other study result indicates wind strength and direction can affect the size of the cooled area around a green space (Yu and Hien, 2006; Zouli, *et al.*, 2009).

In a hot summer the impact of direct radiation on human comfort has emphasized importance that can be described by mean radiant temperature value. Mean radiant temperature is the most important parameter in the human energy balance during summer conditions, since most human bioclimatic thermal indices have special interest in this parameter (Hodder and Parson

2007; Thorsson *et al.*, 2007; Kántor *et al.*, 2013). This fundamentally affects the energy balance of the body and thus the degree of physiological stress (Shashua-Bar *et al.*, 2010). Urban trees on the modification of urban human comfort especially in shading upshot by alteration of radiation energy balance and human bioclimatic impacts by reducing the short wave radiation impact of a human body (Ali- Toudert and Mayer, 2007; Lee *et al.*, 2013).

#### 4. Conclusion and Recommendation

The urban climate can be effectively modified by altering the amounts of heat energy absorbed, stored and transferred by adopting cooling strategies. The local climate regulation is a valuable service for the urban inhabitants by greening because it reduces the urban heat island effect. The green area size, plant density and plant growth stage have strong correlation with urban green area temperature regulation. The political regimes policy and strategy did not significant affect social institutions, however, it affects significantly the enter city green area. The green area influence range is not uniform for all study sites might be due to the influence of interlock environment in the buffering zone. However, the result strongly agrees with literature values in the different climatic zone with varying size of green area, plant density and growth stages. Therefore, researchers should consult urban managers/administration on how to develop sustainable and health urban with sufficient green in our vicinity.

#### 5. Reference

- [1] Krayenhoff, E. S. and Voogt, J. A. (2010). Impacts of urban albedo increase on local air temperature at daily through annual time scales: Model results and synthesis of previous work, *J. Appl. Meteorol. Climatol.* 49, 1634-1648.
- [2] Chandler, T. J. (1962). London's Urban Climate. *Geogr. J.* 127, 279–302.
- [3] Oke, T. R. (1973). City Size and the Urban Heat Island. *Atmos. Environ.* 7, 769–779.
- [4] Ackerman, B (1985). Temporal March of the Chicago Heat Island: *J. Clim. Appl. Meteorol.* 24, 547–554.
- [5] Moreno Garc'ia, M. C. (1994). Intensity and Form of the Urban Heat Island in Barcelona: *Int. J.Clim.* 14, 705–710.
- [6] Yamashita, S. (1990). The urban climate of Tokyo: *Geogr. Rev. Japan*, 63B, 98–107.
- [7] Smith, T.M., Reynolds R.W., and Lawrimore, J. (2008). Improvements to NOAA's historical merged land-ocean surface temperature analysis (1880-2006). *Clim.*, 21, 2283-2296.

- [9] Taha, H. (1988). An urban microclimate model for site-specific building. University of California, Berkeley, C.A.
- [10] Susca, T., Gaffin, S.R. and Dell'Osso, G.R. (2011). Positive effects of vegetation: Urban heat island and green roofs. *Environmental Pollution* 159: 2119-2126.
- [11] Science for Environment Policy (2011). European Commission DG Environment News Alert Service, edited by SCU, The University of the West of England, Bristol, LBL Report No. 26105.
- [12] Frumkin, H., Frank, L., Jackson, R.J. (2004). *Urban Sprawl and Public Health. Designing, Planning, and Building for Healthy Communities.* Island Press, Washington D.C.
- [13] Tratalos, J., R.A. Fuller, P.H. Warren, R.G. Davies, and K.J. Gaston. 2007. Urban form, biodiversity potential and ecosystem services. *Landscape and Urban Planning* 83(4):308-317.
- [14] Mallick, J., Kant, Y., and Bharath, B.D.(2008). Estimation of Land Surface Temperature over Delhi using Landsat-7 ETM+. *J. Ind. Geophys. Union*, 12(3), 131-140.
- [15] Joel, B., Smitha, L., Stephen, H., Schneider, B.C.L., Michael, O., Gary, W., Yohe, E., William, H. F., Michael, D., Mastrandrea, A. P., Ian, B. H., Jan, C.M., Chris, H. D., Magadzaj, H.M.
- [16] F., Barrie, P. A., Atiq, R. A. S., and Jean, P. V. Y. (2009). Assessing dangerous climate change through an update of the Intergovernmental Panel on Climate Change (IPCC) reasons for concern. Contributed by Stephen H. Schneider, December 9, 2008 (sent for review March 18, 2008)
- [17] Smith, M., Sanchirico, J., and Wilen, J. (2009). The Economics of Spatial-Dynamic Processes. Applications to Renewable Resources. *Journal of Environmental Economics and Management*, 57, 104–121.
- [18] Oke, T. R., (1997). Urban climates and global environmental change. *Applied Climatology* R.D.Thompson and A. Perry, Eds., Routledge, 273-287.
- [19] Health and the Environment in the Southeastern United States (2002). *Rebuilding Unity: Workshop Summary by Roundtable on Environmental Health Sciences, Research, and Medicine*, Board on Health Sciences Policy, Institute of Medicine and Howard Frumkin (Nov 1, 2002)
- [20] Environmental health in emergencies (2008). Geneva, World Health Organization. ([http://www.who.int/environmental health emergencies/en/](http://www.who.int/environmental_health_emergencies/en/))
- [21] Taha, H., Akbari, H., Rosenfeld, A., and Huang, J. (1988). Residential cooling loads and the urban heat island the effects of albedo. *Building and Environment* 23(4):271–283.
- [22] Saito, L., Ishihara, O., and Katayama, T. (1991). Study of the effect of green areas on the thermal environment in an urban area. *Energy and Buildings*, 15-16:493-498.
- [23] Saito, I., Ishihara, O., and Katayama, T. (1990-91). Study of the effect of green areas on the thermal environment in an urban area. *Energy and Buildings* 15-16:493-498.
- [24] Jamie, T., Richard, A., Fuller, P. H., Warren, R. G., and Davies, K. J. G. (2007). Urban form, biodiversity potential and ecosystem services. *Landscape and Urban Planning* 83 (2007) 308–317
- [25] Whitford, V., Ennos, A.R., and Handley, J.F. (2001). City form and natural process indicators for the ecological performance of urban areas and their application to Merseyside, UK. *Landscape and Urban Planning*, 57(2), pp. 91–103.
- [26] Mitchell, P., Kleinzeller, A., and Kotyk, A. (160). *Membrane Transport and Metabolism*, Symposium. Academic Press, Inc., New York, in the press.
- [27] Edwards, S., and Hedberg (1989). *Flora of Ethiopia*, vol 3. Addis Ababa and Asmara, Ethiopia and Uppsala, Sweden.
- [28] Edwards, S., Mesfin Tadesse., Sebsebe Demissew and Hedberg, I. (2000). *Flora of Ethiop and Eritrea*, 1st edn, Volume 2, Part 1: Addis Ababa, Ethiopia and Uppsala, Sweden.
- [29] Hedberg, I. And Edwards, S. 1989. *Flora of ethiopia: volume 3.* Addis Ababa and Asmara, Ethiopia and Uppsala, Sweden.
- [30] Lillesand, T.M., and Kiefer, R.W. (1994). *Remote Sensing and Photo Interpretation*, 3rd . ed. John Wiley & Sons: New York. 750 p
- [31] Jauregui, E. (1990/1991). Influence of a large urban park on temperature and convective precipitation in a tropical city. *Energy and Buildings* 15(3–4): 457–463, DOI: 10.1016/0378- 7788(90)90021-A.
- [32] Kotzen, B. (2003). An investigation of shade under six different tree species of the Negev desert towards their potential use for enhancing micro-climatic conditions in landscape architectural development. *J Arid Environ* 55:231-74
- [33] Nakaohkubo, K., Hoyano, A., and Kumakura, E. (2013). Development of thermal and ultraviolet environmental simulation tool for urban living space Part 1, Calculation method of ultraviolet environment. *Summaries of Technical Paper of Annual meeting of Architectural Institute of Japan*, in printing
- [34] Hunter, B. A., Livesley, S.J., Williams, N.S.G. (2012). Responding to the urban heat island. A review of the potential of green infrastructure. Victorian Centre for Climate Change Adaptation Research

- [35] Berry R, Livesley, S.J., Ayeb, L. (2013). Tree canopy shade impacts on solar irradiance received by building walls and their surface temperature. *Build Environ* 69:91-100
- [36] Georgi, J.N., Dimitriou, D. (2010). The contribution of urban green spaces to the improvement of environment in cities. Case study of Chania, Greece. *Build Environ* 45:1401-1414
- [37] Emmanuel, M., (2005). *An Urban Approach to Climate-sensitive Design: Strategies for the Tropics*. Taylor and Francis.
- [38] Hodder, S.G., Parsons, K. (2007). The effects of solar radiation on thermal comfort. *Int J Biometeorol* 51: 233-250
- [39] Kántor, N., Lin, T.P., Matzarakis, A. (2013) Daytime relapse of the mean radiant temperature based on the six directional method under unobstructed solar radiation. *Int J Biometeorol* DOI 10.1007/s00484-013-0765-5
- [40] Thorsson, S., Lindberg, F., Eliasson, I., Holmer, B. (2007). Different methods for estimating the mean radiant temperature in an outdoor urban setting. *Int J Climatol* 27:1983-1993
- [41] Susca, T., Gaffin, S.R., Dell'Osso, G.R. (2011). Positive effects of vegetation. Urban heat island and green roofs. *Environmental Pollution*, 159 (8-9), 2119-2126.
- [42] Givoni, B. (1998). Impact of green areas on site and urban climates. *Climate considerations in building and urban design*. J. Wiley & Sons, New York. pp. 303-30
- [43] Honjo, T., Takakura, T. (1990/91). Simulation of Thermal Effects of Urban Green Areas on their Surrounding Areas. *Energy and Buildings*, 15-16, 443.
- [44] Potchter, O., Cohen, P., Bitan, A. (2006). Climatic behavior of various urban parks during hot and humid summer Mediterranean city of Tel Aviv. *Israel. International Journal of Climatology* 26: 1695 – 1711.
- [45] Yu, C., and Hien, W.N. (2006).. Thermal benefits of city parks. *Energy and Buildings* 38, 105-20
- [46] Zouli, I., Santaouris, M., and Dimoudi, A. (2009). Monitoring the effect of urban green areas on the heat island in Athens. *Environmental Monitoring and Assessment* 156, 275-92.
- [47] Shashua, B. L., Potchter, O., Bitan, A., Bolanski, D., Yaakov, Y. (2010). Micro-climate modeling of street tree species within the varied urban morphology in the Mediterranean city of Tel Aviv. *Int. J. Climatol.*, 30: 44-57.
- [48] Ali-Toudert, F., and Mayer, H. (2007). Effects of asymmetry, galleries, overhanging facades and vegetation on thermal comfort in urban street canyons. *Solar Energy*, 81:742-754.
- [49] Lee, W.L., Liou, K., and Hall, A. (2011). Parameterization of solar fluxes over mountain surfaces for application to climate models. *J. Geophys. Res.-Atmos.*, 116, D01101, doi:10.1029/2010JD014722, 2011.

6/22/2023