



Influence Of Rubber Effluent On The Microbial Population and Physicochemical properties Of Soils

¹Iyakndue, M. L., ¹Brooks, A. A., ¹Unimke, A. A. and ²Agbor R.B

¹Department of Microbiology, University of Calabar, PMB 1115, Calabar, Cross River State, Nigeria.

²Department of Genetics and Biotechnology, University of Calabar, Calabar

E-mail: agborreagan@yahoo.com

Tel: +234 8063430125

Abstract: The study was carried out by artificially polluting an agricultural soil in Calabar with varying concentrations (0ml, 250ml, 500ml, 1000ml and 2000ml) of rubber effluent, in which 0ml served as a control, with the aim of assessing their effect on soil microflora and fertility. The polluted soil was analysed in terms of the following parameters; microbial population, soil PH organic matter, total nitrogen, available phosphorus, electrical conductivity, calcium, magnesium, potassium, sodium, effective cation exchange capacity, exchangeable acidity and base saturation. In the polluted soils, the total heterotrophic bacteria, total heterotrophic fungi and total heterotrophic actinomycetes increased significantly ($p < 0.05$) with a decrease in the concentration of pollutants. The total heterotrophic bacteria and total heterotrophic actinomycetes showed significant reduction with an increase in the length of pollution while total heterotrophic fungi did not show difference ($p > 0.05$) over the duration of pollution. Microorganisms isolated from the polluted soil were *Pseudomonas sp.*, *Bacillus sp.*, *Staphylococcus sp.*, *Micrococcus sp.*, *Flavobacterium sp.*, *Mucor sp.*, *Fusarium sp.*, *Penicillium sp.*, *Aspergillus sp.*, *Rhizopus sp.*, and *Streptomyces sp.* In the polluted soil, pH, organic matter, total nitrogen, potassium, available phosphorus, magnesium and sodium, showed significant differences ($p < 0.05$) in their values with the control, while calcium, electrical conductivity, base saturation, effective cation exchange capacity, and exchangeable acidity did not show significant difference ($p > 0.05$) with that of the control. The results of this study revealed that light application rubber effluent could enhance microbial proliferation and thus, increases soil fertility, while a heavy application inhibits the same.

[Iyakndue, M. L., Brooks, A. A., Unimke, A. A. and Agbor R.B. **A Influence Of Rubber Effluent On The Microbial Population and Physicochemical properties Of Soils.** *Nat Sci* 2024,22(8):28-37]. ISSN 1545-0740 (print); ISSN 2375-7167 (online). <http://www.sciencepub.net/nature> 05. doi:[10.7537/marsnsj220824.05](https://doi.org/10.7537/marsnsj220824.05).

Key Words: Rubber, Effluent, Soil, Microbial, Population

Introduction

Various waste materials including solid and liquid wastes and volatile organic gases produced in these industries are being continuously released into the environment without or in few cases with partial treatment and affect the basic environmental components such as, soil, water and air (Hossan, 2004; Bhabani, 1977). Soil and environment are under tremendous pressure due to industrial expansion and discharge of effluents. The industrial effluents and water drainage from spoil and rubbish heaps either washes direct to nearby fields and enter the local streams, river and ultimately into the soil. Once pollutants enter and are incorporated into the soil, the concentration in soil continuously increases and accumulates, and become toxic to all forms of life like plants, microorganisms and human being (Dilip, 2006).

Soil is an efficient purifying medium with a great capacity to receive and decompose wastes and matter by its microflora and precipitate out nutrients (Nagaraju *et al.*, 2007). However, if the input of the pollutants exceeds the soil purifying limit, the

effectiveness of soil microorganism activity is reduced considerably. As a result, there occurs an adverse change in the soil physico-chemical properties which consequently affect the growth and development of the crop plants (Jolly, *et al.*, 2008). Industrial effluents as pollutants contain a large number of both known and unknown substances formed during the production process. There is a direct impact of pollutants on minerals, organic matter and microbial community of soil (Nagaraju *et al.*, 2007). The discharge of industrial effluents especially without treatment may have profound influence on physico-chemical and biological properties of soil related to soil fertility.

Rubber effluent is known to contain a large amount of non-rubber substances in addition to traces of various processing chemicals. The controlled applications of rubber effluent on land have been reported to cause changes in soil properties (Orhue *et al.*, 2007). Rubber latex processing for example, involves sequential immersion in various chemicals before the final products are ready for the market. This process leaves behind toxic and concentrated aqueous solution

with obnoxious odour (Webster *et al.*, 1989). The discharge of such mixture may give rise to various types of harmful effects or outright pollution in the receiving environment (Nyholm, 1992). Pollution action of rubber effluents is due to the presence of large amounts of dissolved organic and inorganic solids and a large number of viable indicative which create a high oxygen demand (Soyza, *et al.*, 1994).

Materials and methods

Experimental design

The soil used in this study was an agricultural soil of the University of Calabar. The pollution of the soil was performed artificially with four concentrations (0ml, 250ml, 500ml, 1000ml and 2000ml) of rubber effluent. And the duration of this study was sixteen weeks.

Collection of samples

Soil sample

Soil samples of 0-30cm depth were collected from five locations on an agricultural soil within university of calabar by excavation using spade. The collected samples from all locations were thoroughly mixed on the spot in order to obtain composite sample. Five kilograms of this soil was weighed into four different polythene bags.

Rubber Mill Effluents Samples

Rubber effluent was collected from Pamol (Nigeria) Limited Estate, Odukpani, in Cross River State. The samples were collected with clean plastic containers rinsed several times with the same sample and transported within 24 hours to the Department of soil science, University of Calabar for physicochemical analysis and pollution of the soil.

Pollution of the Soil Sample

Pollution of the soil was achieved by employing the method of Orhue *et al* (2007). 0ml, 250ml, 500ml, 1000ml and 2000ml of the samples was added to 4kg of already dried soil in each polyethylene and mixed thoroughly for even distribution. These polluted soils and the unpolluted soil (control) were left outside under normal environmental condition for the duration of 16weeks (Four months)

Microbial analysis

Microbial analysis was carried out before and after pollution. 10g of Soil sample was collected aseptically, labelled and store in ice packed plastic coolers and transported to the Microbiology Department Laboratory University of Calabar where microbial analysis was carried out within 24 hours of sampling so as to maintain the stability of the sample without significant alteration in the microbial population.

Dilution

Serial dilution was carried out by weighing 10g of soil in to 90ml of sterile sterile water contained in a stoppered 200ml volumetric flask and agitated to dislodge the microorganisms from the soil particles. From this initial dilution, a ten-fold serial dilution was prepared.

Enumeration of heterotrophic Bacteria

The counts of total heterotrophic bacteria in the soil samples was determined by pour plating 1ml of desired dilutions into nutrient agar (NA). The medium was incorporated with antifungal agent (50µg/ml Nystatin), in order to prevent the growth of fungal contaminants. Bacterial colonies were counted after 24 hours of incubation at room temperature and reported as a number of colony forming units (CFU) per gram of soil.

Enumeration of heterotrophic Fungi

The total heterotrophic fungi count was measured by pour plating 1ml of 10⁻³dilution into Sabouroud dextrose agar (SDA) supplemented with antibacterial agents (50µg/ml of streptomycin and 30µg/ml of penicillin) to inhibit the growth of bacterial contaminants. Fungal counts were reported after 72 hours of incubation.

Enumeration of heterotrophic Actinomycetes

Enumeration of total actinomycetes was achieved by pour plate technique. 1ml of 10⁻²dilution was plated unto sodium caseinate agar, 50µg/ml of nystatin and 30µg/ml of tetracycline was added to inhibit fungal and bacterial growth. An actinomycetes count was reported 7 days after incubation at room temperature.

Maintenance of pure isolates

Bacterial colonies were repeatedly transferred to freshly prepared nutrient agar plates by the streak-plate method and allow growing for 48 hours before stocking. Similarly, distinct fungal and actinomycetes colonies were subculture repeatedly on freshly prepared Sabouroud dextrose agar plates and sodium caseinate agar, respectively. Pure isolates of the microorganisms were maintained on agar slants as stock, which were preserved in the refrigerator for further use.

Characterization and Identification of isolates

Various methods in Etok, *et al.*, (2004) General Microbiology Practical Manual were used to characterize and identify the isolates. The test results for bacteria were evaluated using characteristics presented in *Bergey's Manual of Determinative Bacteriology* (Holt *et al.*, 1994).

Representative colonies of fungal isolates were characterized and identified based on their cultural and morphological features as described by Barnett and Hunter (1987). The characterizations were achieved

through staining techniques-using lactophenol in cotton blue.

Soil physico-chemical analysis

Physicochemical analysis of the pristine soil was carried out before pollution. And after pollution with varying concentrations of the effluent on the soils, physiochemical analysis of each soil sample polluted with different concentrations of each effluent was also carried out bimonthly (every 8weeks) .

Particle size and textural class analysis

In carrying out this test, the Bouyoucos-type hydrometer method described by Day (1986) was used.

Soil pH

Soil pH was determined in water 1:2 soils: water ratio using pH meter with glass electrode. 20g of air-dried soil was weighed into a 50ml beaker, and 20ml of distilled water was added and allowed it to stand for 30minutes. The electrode of the pH meter was inserted into the 1:2.5 soil /water partly settled suspension and measured the pH. The result was recorded as soil pH measure in water.

Electrical conductivity (EC)

In the same soil solution (1:2.5 soil /water solution) for pH determination, electrical conductivity electrode was inserted into the partly settled suspension and the EC was measured.

Organic matter

This was determined by the dichromate wet-oxidation method as described by Nelson and Sommers (1996).

Total nitrogen

Total nitrogen was determined by the micro-Kjeldahl method as described by Bremner (1996).

Available phosphorus

Available phosphorus was extracted with acid fluoride using Bray P-1 method described by Bray and Kurtz (1996).

Exchangeable cations

The bases were extracted with neutral NH_4OAC . Calcium and magnesium were determined in the extract by EDTA titration, and potassium and sodium by the use of flame photometer (Udo *et al.*, 2009).

Exchangeable acidity

A^+ and H^+ were obtained by leaching the soil with INKCl solution and the and the extract titrated with standard NaOH as described by I.I.T.A (1979).

Exchangeable acidity ($\text{Al} + \text{H}^+$) –
Exchangeable $\text{Al} = \text{Exchangeable H}$

Effective cation exchange capacity

This was determined by calculation. That is, total exchangeable bases ($\text{Ca} + \text{Mg} + \text{Na}$) + Exchangeable acidity (EA).

Percentage base saturation

This will be achieved by dividing the total exchangeable bases by exchangeable cation capacity and multiplied by 100 (Agbenin, 1995).

$\% \text{ base saturation} = \frac{\text{summation of exchangeable bases} \times 100}{\text{ECEC}}$

Results

Microbial analysis

Table 1 below shows the enumeration of total heterotrophic bacteria (THB), total heterotrophic fungi (THF) and total heterotrophic actinomycetes (THA) in the pristine soil. The counts obtained were as follows $1.90 \pm 1.41 \times 10^7$ Cfug/g, $1.29 \pm 1.25 \times 10^5$ Cfug/g and $9.2 \pm 1.25 \times 10^3$ cfu/g respectively.

TABLE 1

Total heterotrophic bacteria, total heterotrophic fungi and total actinomycetes counts of the soil before the commencement of the study from an agricultural soil from University of Calabar

THB (CFU/g)	$1.90 \pm 1.41 \times 10^7$
THF (CFU/g)	$1.29 \pm 1.25 \times 10^5$
THA (CFU/g)	$9.2 \pm 1.25 \times 10^3$

Key: THB = Total heterotrophic bacteria, THF= Total heterotrophic fungi, THA= Total heterotrophic actinomycetes, CFU/g = Colony forming unit/gram

Table 2 below shows the effect of different concentrations of rubber effluent on microbial population. The results obtained for THB in 250ml, 500ml, 1000ml and 2000ml were $1.4 \pm 1.95 \times 10^7$ cfu/g , $1.23 \pm 1.26 \times 10^7$ cfu/g, $7.8 \pm 1.48 \times 10^6$ cfu/g and $5.3 \pm 1.31 \times 10^6$ cfu/g respectively. The results showed that there was no significant difference ($p < 0.05$) in the mean counts of THB obtained in 250ml and 500ml of the polluted soils and no significance difference ($p < 0.05$) was also observed in THB counts of 1000ml 2000ml of the effluent polluted soils (Table 2).

The results for the counts of THF in different concentrations of the effluent polluted soils showed a significant increase in THF counts of 250ml and 500ml with mean count of $2.58 \pm 3.80 \times 10^5$ cfu/g, $1.93 \pm 2.84 \times 10^5$ cfu/g respectively over the control while there were no significant difference in THF counts of 1000ml and 2000ml with mean counts of $1.05 \pm 2.26 \times 10^5$ cfu/g and $9.4 \pm 2.04 \times 10^4$ cfu/g respectively over the control (Table 2).

The total heterotrophic actinomycetes (THA) counts in different concentrations of polluted soils showed that, there were significant differences ($P < 0.05$) in THA counts among concentrations. There was

a significant increase ($p < 0.05$) in 250ml with a mean count of $1.25 \pm 2.34 \times 10^4$ cfu/g while there were no significant differences ($p > 0.05$) in THA in 500ml, 1000ml and 2000ml of the effluent polluted soil with

mean counts of $9.9 \pm 1.64 \times 10^3$ cfu/g, $6.9 \pm 1.34 \times 10^3$ cfu/g and $5.9 \pm 1.20 \times 10^3$ cfu/g respectively over the control (Table 2).

TABLE 2
Effect of concentrations of pollution on microbial population in rubber effluent polluted soil

	Rubber effluent				
	0ml	250ml	500ml	1000ml	2000ml
THB (cfug ⁻¹)	$1.87^a \pm 2.08 \times 10^7$	$1.47^a \pm 1.95 \times 10^7$	$1.23^a \pm 1.26 \times 10^7$	$7.8^b \pm 1.48 \times 10^6$	$5.3^b \pm 1.31 \times 10^6$
THF (cfug ⁻¹)	$1.28^c \pm 3.06 \times 10^5$	$2.58^a \pm 3.80 \times 10^5$	$1.93^b \pm 2.84 \times 10^5$	$1.05^c \pm 2.26 \times 10^5$	$9.4^c \pm 2.04 \times 10^4$
THA (cfug ⁻¹)	$9.3^b \pm 2.04 \times 10^3$	$1.25^a \pm 2.34 \times 10^4$	$9.9^b \pm 1.64 \times 10^3$	$6.9^b \pm 1.34 \times 10^3$	$5.9^b \pm 1.20 \times 10^3$

Key: means with the same letter along the horizontal arrays indicates significant difference ($P < 0.05$), and means with different letter along the column indicates no significance difference ($P > 0.05$)

THB = Total heterotrophic bacteria, THF= Total heterotrophic fungi, THA= Total heterotrophic actinomycetes, CFU/g = Colony forming unit/gram

Table 3 below shows the effect rubber effluent and duration of pollution on microbial populations. There was a significant reduction ($P < 0.05$) in THA counts obtained after 16 weeks of pollution as compared to 8 weeks of pollution.

TABLE 3
Effect of duration of pollution on microbial population in rubber effluent polluted soils

	8 weeks	16 weeks	LSD
THB(CFU g ⁻¹)	$1.64 \pm 2.42 \times 10^7$ a	$7.11 \pm 1.04 \times 10^6$ b	2.69
THF (CFU g ⁻¹)	$1.67 \pm 3.02 \times 10^5$ a	$1.44 \pm 2.86 \times 10^5$ a	3.38
THA (CFU g ⁻¹)	$1.09 \pm 2.73 \times 10^4$ a	$6.9 \pm 0.86 \times 10^3$ b	4.04

Key: means with the same letter along the horizontal arrays indicates significant difference ($P < 0.05$), and means with different letter along the column indicates no significance difference ($P > 0.05$)

THB = Total heterotrophic bacteria, THF= Total heterotrophic fungi, THA= Total heterotrophic actinomycetes, CFU/g = Colony forming unit/gram, LSD= Least significant difference

Characterization and

Identification of Microbial Isolates

Microbial isolates were characterized and identified based on their morphological and biochemical characteristics. The following microorganisms were isolated from rubber effluent polluted soils were *Staphylococcus* spp., *Bacillus* spp., *Pseudomonas* spp., *Flavobacterium* spp., *Micrococcus* spp., *Mucor* spp., *Fusarium* spp., *Penicillium* spp., *Aspergillus* spp., *Rhizopus* spp., and *Streptomyces* spp.

Physicochemical Properties of the Soil

In this study, the physico- chemical properties of the soil that were considered include, particle size distribution, soil pH organic matter, available phosphorus, total nitrogen, exchangeable bases, exchangeable acids, effective cation, exchange capacity, base saturation and electrical conductivity.

The physico-chemical properties of the soil sample before the commencement of the study are presented in Table 4 while that of the soil treated with varying concentration (0ml, 250ml, 500ml 1000ml and 2000ml) of effluent are presented in Table 5.

Soil pH

The pH values of the control (0ml) soils were in the range of 5.66 ± 0.50 to 5.7 ± 0.21 . The pH values in rubber effluent treated soils ranged from 6.88 ± 0.08 to 7.03 ± 0.09 .




Figure 1: Effect of concentrations of rubber effluent on soil pH

Organic Matter Content

The values of organic matter in control (0ml) soils were in the range of $2.41 \pm 0.23\%$ and $2.41 \pm 0.09\%$. The organic matter content of the soils treated with varying concentrations of rubber effluent treated soils ranged from $2.68 \pm 0.18\%$ to $4.24 \pm 0.26\%$ (Figure 2).




Figure 2: Effect of concentrations of rubber effluent on soil organic matter

Total Nitrogen Content

The control soil (0ml) had the total nitrogen content in the range of $0.06 \pm 0.2\%$ and $0.06 \pm 0.01\%$. In the soil treated with varying concentrations of rubber mill effluent, nitrogen content ranged from $0.09 \pm 0.02\%$ to $0.15 \pm 0.04\%$. Significant difference ($p < 0.05$) was not observed among concentrations of pollution (Table 5).




Figure 3: Effect of concentrations of rubber effluent on total nitrogen content of the polluted soils

Effective Cation Exchange Capacity (ECEC)

The values of ECEC in the control (0ml) soil were in the range of 9.45 ± 0.43 cmol/kg and 9.7 ± 0.06 cmol/kg (Table 5). The value of ECEC in rubber effluent polluted soil ranged from 9.43 ± 0.11 cmol/kg to 9.86 ± 0.20 cmol/kg.

TABLE 4
Physico-chemical properties of the soil before the commencement of the study from an agricultural soil in University of Calabar

Parameters	Values
pH	5.60
Electrical conductivity (ds/m)	0.055
Organic matter (%)	2.41
Total nitrogen (%)	0.06
Available Phosphorus (mg/kg)	31.54
THC (mg/kg)	BDL
Ca (cmol/kg)	4.00
Mg (cmol/kg)	2.10
Na (cmol/kg)	0.06
K (cmol/kg)	0.21
EC (cmol/kg)	1.11
ECEC (cmol/kg)	8.82
Base saturation (%)	85.16
Sand (%)	75.80
Silt (%)	12.60
Clay (%)	11.60
Lead (mg/kg)	6.20
Nickel (mg/kg)	9.44
Cadmium (mg/kg)	4.32
Chromium (mg/kg)	5.60
Iron (mg/kg)	919.66

TABLE 5
Effect of duration of pollution on physico- chemical properties of the soil polluted with rubber mill effluent

	8weeks	16weeks	LSD
pH	6.96 ^a ±0.03	6.45 ^c ±0.05	0.13
Organic matter	3.31 ^c ±0.11	3.03 ^d ±0.06	0.11
Avail. P	19.28 ^a ±0.18	17.93 ^b ±0.10	0.54
Total N	0.11 ^a ±0.03	0.11 ^a ±0.01	0.01
Mg	2.66 ^b ±0.07	1.85 ^d ±0.08	0.06
Na	0.10 ^c ±0.03	0.23 ^a ±0.02	0.02
Ca	5.46 ^a ±0.08	4.24 ^b ±0.10	0.38
Ec	0.08 ^b ±0.01	0.09 ^b ±0.02	0.01
Bs	78.80 ^a ±1.83	73.78 ^b ±0.98	0.86
K	0.22 ^b ±1.04	0.18.18 ^c ±0.06	0.01
ECEC	10.71 ^a ±0.16	8.81 ^c ±0.18	0.44
EA	2.27 ^a ±0.09	2.31 ^a ±0.08	0.13

Key: means with the same letter along the horizontal arrays indicates significant difference ($P < 0.05$), and means with different letter along the column indicates no significance difference ($P > 0.05$).

TABLE 6
Effect of concentrations of pollution on the physiochemical properties of the soil polluted with rubber mill effluent

	0ml	250ml	500ml	1000ml	2000ml	LSD
PH	5.75 ^c ±0.21	6.93 ^a ±0.14	7.03 ^a ±0.09	6.92 ^a ±0.10	6.88 ^a ±0.08	0.20
Organic matter	2.41 ^e ±0.09	2.83 ^d ±0.20	2.68 ^d ±0.18	3.68 ^c ±0.22	4.24 ^b ±0.26	0.18
Avail. P	31.54 ^a ±2.50	13.53 ^e ±1.09	14.73 ^d ±1.18	17.09 ^c ±0.98	16.16 ^d ±0.5	0.85
Total N	0.06 ^c ±0.01	0.12 ^b ±0.03	0.09 ^b ±0.02	0.13 ^b ±0.01	0.15 ^b ±0.04	0.02
Mg	2.40 ^d ±0.06	2.15 ^e ±0.10	2.08 ^e ±0.07	2.31 ^d ±0.12	2.33 ^d ±0.08	0.09
Sodium (Na)	0.09 ^b ±0.01	0.10 ^b ±0.02	0.11 ^b ±0.01	0.45 ^a ±0.03	0.10 ^b ±0.01	0.04
Calcium (Ca)	4.93 ^a ±0.04	4.73 ^a ±0.03	4.75 ^a ±0.03	4.80 ^a ±0.08	4.89 ^a ±0.04	0.61
Ec	0.05 ^b ±0.01	0.10 ^b ±0.02	0.07 ^b ±0.01	0.9 ^b ±0.01	0.09 ^b ±0.01	0.02
Base saturation	78.66 ^b ±0.58	75.61 ^b ±1.03	74.53 ^b ±1.11	76.28 ^b ±0.98	77.18 ^b ±0.84	1.35
K	0.21 ^b ±0.06	0.15 ^b ±0.04	0.19 ^b ±0.03	0.19 ^b ±0.02	0.29 ^a ±0.20	0.02
EFEC	9.7 ^b ±0.06	9.43 ^b ±0.11	9.54 ^b ±0.18	10.16 ^b ±0.20	9.86 ^a ±0.26	0.69
EA	2.07 ^a ±0.05	2.30 ^a ±0.08	2.43 ^a ±0.01	2.41 ^a ±0.06	2.25 ^a ±0.04	0.20

Key: means with the same letter along the horizontal arrays indicates significant difference ($P < 0.05$), and means with different letter along the column indicates no significance difference ($P > 0.05$).

Discussion

The pollutant level of rubber effluent varies with the quality of the raw material and production process used to manufacture the rubber from the latex. In this study, following pollution of soil with varying concentrations of rubber effluent, the soil showed profound changes in the microbial populations and physico-chemical properties of the soil.

In assessing the effect of different concentrations of rubber effluents on microbial population, the result showed no significant difference ($p > 0.05$) in THB mean count in 250ml and 500ml of effluent polluted soils over the control. This implies that the THB mean counts in 250ml and 500ml of polluted soils fall in the same range with that of the control. There was a significant reduction ($p \leq 0.05$) in THB in 1000ml and 2000ml over the control. Also, the result showed that

there was a significant increase ($p \leq 0.05$) in THF and THA in 250ml and 500ml of polluted soil (Table 2). Significant increase ($p \leq 0.05$) in THA count was also observed in 250ml and 500ml of polluted soil which had THA count higher than the control. This may be attributed to the acidic nature of the effluent and the soil. Soil actinomycetes can tolerate a pH level of up to 6.80, and this was the pH level in most of the treated soils in this study. In terms of concentrations of pollution, 250ml of effluents was found to be most favourable concentration for the proliferation of microorganism, followed by 500ml in this study. Similar result had earlier been reported by Nguago *et al.*, (2008).

Also in assessing the effect of duration of pollution and different concentrations of pollutant on microbial population, THB showed significant increase ($p < 0.01$) with decrease in concentrations of pollutant and significant reduction with increase in the duration of pollution (Table 3). This result implies that there was a steady reduction in THB with increase in concentrations and length of pollution (Table 3). Also, the THF mean counts at 250ml of rubber effluent polluted soil showed significant increase ($p < 0.05$) after 8 weeks of pollution and a significant reduction was observed in the same concentration of pollution after 16 weeks of pollution and while there was no significant difference in THF count at 500ml, 1000ml after 8 weeks and 16 weeks of pollution (Table 3). Similarly, there was significant increase ($p < 0.05$) in THA at 250ml, 500ml, after 8 weeks of pollution and 250ml after 16 weeks of pollution, and no significant difference was also observed in THA in 1000ml, 2000ml after 8 weeks of pollution and 500ml, 1000ml and 2000ml after 16 weeks of pollution. This results implies that, there was a significant increase ($p < 0.05$) in THB, THF and THA in all the samples treated with rubber mill effluents after 8 weeks of pollution, followed by a reduction in THB and THF after 16 weeks of pollution except THA which did not show reduction after 16 weeks of pollution (Table 3).

There were significant changes in physico-chemical properties. These changes have earlier been reported by Russell *et al.*, (1988) that continual applications of effluents on the soil can change soil properties, e.g. pH and nutrient concentrations.

References

- [1]. Bray, R. H. and Kurtz, L. T. (1945). Determination of Total, Organic and Available forms of Phosphorus in Soils. *Journal of Soil Science*. (5) : 35-45.
- [2]. Bremner, J. M. (1996). Total Nitrogen. In: D. L. Sparks (eds). *Methods of Soil analysis. Part 3-chemical methods*, SSSA Book Series 5, Madison, Wisconsin, Pp 1085-1122
- [3]. Carter, M. R. (2002). Soil quality for sustainable land management: organic matter and aggregation interactions that maintain soil functions. *Agronomy Journal*. 94: 38-47.
- [4]. Day, P. R. (1965). Particle Fractionation and particle size analysis. In: C. A. Black (eds). *determination of phosphorus in Natural waters. Analytical Chemistry ACTA* 27:31-38.
- [5]. Dilip K. D. (2006). *Introductory Soil Science*, Kalyani Publishers, Delhi.
- [6]. Dunphy, P. J., Whittle, K. J., Pencoek, J. F. and Morton, R.A. (1965). Identification and Estimation of tocotrienols in Heave latex. *Nature*, 207:521
- [7]. Holt, J. G., Krieg, N. R., Senath, P. H., Stanley, L. T. and Williams, S. T. (194). *Bergey's Manual of Determinative Bacteriology*. (9th Ed). Baltimore, Willams and Wilkins.
- [8]. Jolly, Y. N., Islam, A. and Quraishi, B. (2008). Effects of Paint Industry Effluent on Soil Productivity. *Journal of Bangladesh Academy of Sciences*, Vol. 32, No. 1, 41-53,
- [9]. Kulkarni, P. R. (1972). Recovery of algae grown on rubber effluent. In: *Waste Recovery by Micro-organisms*. Stanton WR (Ed). Ministry of Education, Kuala Lumpur, Malaysia, pp. 118 - 128.
- [10]. Kulkarni, P. R., Ho, P., Ratnasabapathy, M., and Stanton, W. R. (1973a). Utilisation of rubber effluent (1). *Planter* 49: 307 - 313.
- [11]. Láng, I. (1993). Environmental Management and Sustainable Development. In Nath, B., Hens, L. and Devuyt, D (Eds.). *Environmental Management, Vol. 3. Instruments for Implementation Brussels*: VUB University Press, VUB University.
- [12]. Nagaraju, M., Narasimha, G. and Rangaswami, V. (2009). Influence of sugar industry effluents on soil enzyme activities. *Ecol. Environ. Cons.*, 15:89-94.
- [13]. Narasimha, G., Babu, A. K. and Rajasekhar, R. B. (1999). Physico-chemical and biological properties of soil samples collected from soil contaminated with effluents of cotton ginning industry. *J. Environ. Biol*, 20: 235-239.
- [14]. Nelson, D. W., Sommers, L. E. (1982). Total carbon, organic carbon and matter. In: Page, A. L., Miller, R. H. and Keeney, D. R. (eds.) *Methods of Soil Analysis, Part 2 Chemical and Microbiological Properties*. Madison, Wisconsin pp. 539-579.
- [15]. Nilima, W. and Madhuri, S. (2005). Fungal diversity of paper industry waste and soils irrigated by the effluents. *Nat. Environ. Pollut. Tech.*, 4: 44-52.
- [16]. Nyholm, N. (1992). Environmental Impact Assessment and Control of Marine Industrial of

- Soil Analysis, Part 1- Physical and Mineralogical Methods*, SSSA Book Series 5,
- [17]. Okwute, O. L. and Isu, R. N. (2007). Impact Analysis of palm oil mill effluent on the Aerobic Bacterial density and ammonium oxidizers in a dumpsite in Anyigba, Kogi State. *Afr. J. Biotech* 6(2):116-119.
- [18]. Orhue, E. R., Uzu, O. F. and Osaigbovo, A, U. (2007). Effect of Combining Rubber Effluent with Rock Phosphate on Some Soil Chemical Properties and Early Growth of Maize (*Zea mays*. L). *International Journal of Soil Science*, 2: 82-95.
- [19]. Prescott, L.M., Klein, D, A. and Harley, J. P. (2001). Microbiology 5th edition. McGraw Hill, Education, United States.
- [20]. Russell, J. M., Cooper, R. N. And Donnison A. M. (1988). Irrigation of high-nitrogen containing wastes to pasture. In: Bhamidimarri, R. (Ed.), *Alternative Waste Treatment Systems*. Elsevier Applied Science, London. Pp.45-54.
- [21]. Savin, M. C., Görres, J. H. and Amador, J. A.(2004). Microbial and microfaunal community dynamics in artificial and *Lumbricus terrestris* (L.) burrows. *Soil Sci. Soc. Am.* 68:116-124.
- [22]. Schachtschabel, P., Blume, H.-P., Hartge, K.H., Schwertman, U., Brummer, G. and Renger, M. (1984). *Lehrbuch der Bodenkunde*, 11th edn. Ferdinand Enke Verlag, Stuttgart.
- [23]. Schipper, L.A., Williamson, J. C. Kettles, H. A., and Speir, T. W. (1996). Impact of land-applied tertiary-treated effluent on soil biochemical properties. *J. Environ. Qual.*, 27: 1073-1077.
- [24]. Scholes, R. J., Dala, I. R. And Singer, S. (1994). Soil Physics and Fertility: The Effects of Water, Temperature and Texture. In: Wooner P. L. And Swift M.J.(eds), *The Biological Management of Soil Fertility*. John Willey and Sons, Chichester, pp. 117- 136.
- [25]. Soyzo, Y. N. And Yapa P. A. (1994). Use of Rubber Factory effluent As A Fertilizer for Young Heavea Plants. *Jl. Rubb. Res. Inst Sri Lakan* 74: 1-9
- [26]. Udo, E. J., Ibia, T. O., Ogunwale, J. A., Ano, A. O. and Esu, I. E. (2009). *Manual of Soil , Plant and Water Analysis*. Sibon Books Limited, Lagos.
- [27]. Unimke, A.A., Antai, S.P. & Agbor, R.B. (2014). Influence of Seasonal Variation on the Microbiological and Physicochemical Parameters of Imo River Estuary of the Niger Delta Mangrove Ecosystem. *American International Journal of Biology*. 2 (1): 61 – 74.
- [28]. Unimke AA, Antai SP, Agbor RB, Nseabasi NO & Agbo BE. (2014). Evaluation of Seasonal Variation in the Microbial and Heavy Metal Concentrations of Imo River Estuary of the Niger Delta Mangrove Ecosystem. *Advance Research in Agriculture and Veterinary Science*. 1 (2): 88 – 94.
- [29]. Webster, C. C. and Kwill, B. W. J. (1989). Rubber. *Tropical Agricultural Series*.
- [30]. with dairy waste water. *Asian J. Microbiol. Biotech. Environ. Sc.*,12, (2010).
- [31]. with dairy waste water. *Asian J. Microbiol. Biotech. Environ. Sc.*, 12,

8/1/2024