

Review on Bovine Fasciolosis and its Economic Significance

Marie Teshager

Banja Woreda livestock office, Injibara, Ethiopia

teshagermarie@gmail.com

ABSTRACT: Bovine fasciolosis is an economically important parasitic disease of cattle caused by Fasciolidaetrematodes of the genus Fasciola. The two most important species of this genus, *Faciola hepatica* and *Faciolagigantica*, are commonly known as liver flukes. The distribution of fasciolosis is worldwide, however, the distribution of *Faciola hepatica*, is limited to temperate areas and highlands of tropical and sub-tropical regions while *Faciolagigantica*, which predominates in tropical area. The definitive hosts for *Faciola hepatica* are most mammals among which sheep and cattle are the most important once. The geographic distribution of trematode species is dependent on the distribution of suitable species of snails. Ethiopia has a high livestock population but productivity is low as a result of diseases, malnutrition and other management problems. Both *Faciola hepatica* and *Faciolagigantica* species of genus *Faciola* are found in Ethiopia and are transmitted by *Lymnaea truncatula* and *Lymnaea natalensis*, respectively and various reports indicated that it is a serious problem of livestock production in Ethiopia causing considerable economic losses. Diagnosis based on clinical signs, seasonal occurrence and a previous history of fasciolosis. Fasciolosis causes a substantial economic loss which includes death, loss in carcass weight, reduction in milk yield, condemnation of affected liver, decline production and productive performances, exposure of animals to other diseases due to secondary complications.

Marie Teshager. **Review on Bovine Fasciolosis and its Economic Significance.** *Nat Sci* 2024,22(10):21-26]. ISSN 1545-0740 (print); ISSN 2375-7167 (online). <http://www.sciencepub.net/nature> 04. doi:[10.7537/marsnsj221024.04](https://doi.org/10.7537/marsnsj221024.04).

Key words Fasciolosis,Bovine,

Introduction

Among many parasitic problems of farm animals, fasciolosis is a major disease, which imposes direct and indirect economic impact on livestock production, particularly of sheep and cattle (Menkir *et al.*, 2007). *Faciola hepatica* and *Faciolagigantica* are the two liver flukes commonly reported to cause fasciolosis in ruminants. The life cycle of these trematodes involves snail as an intermediate host (Walker *et al.*, 2008).

Bovine fasciolosis is an economically important parasitic disease of cattle caused by Fasciolidaetrematodes of the genus *Faciola*. The two most important species of this genus, *F. hepatica* and *F. gigantica*, are commonly known as liver flukes. The distribution of fasciolosis is worldwide, however, the distribution of *F. hepatica*, is limited to temperate areas and highlands of tropical and sub-tropical regions (Soulsby, 1986) and *Faciolais* commonly recognized as liver flukes and are responsible for wide spread morbidity and mortality in cattle characterized by weight loss, anemia and hypoproteinemia. The two most important species are *Faciola hepatica* found in temperate area and in cooler areas of high altitude in the tropics and subtropics and *Faciolagigantica*, which predominates in tropical area (Troncy, 1989).

The definitive hosts for *F. hepatica* are most mammals among which sheep and cattle are the most important once. The geographic distribution of trematode species is dependent on the distribution of suitable species of snails. The genus *Lymnaea* in general and *L. truncatula* in particular is the most common intermediate hosts for *F. hepatica*. This species of snail was reported to have a worldwide distribution (Urquhart *et al.*, 1996). Ethiopia has a high livestock population but productivity is low as a result of diseases, malnutrition and other management problems. Both *F. hepatica* and *F. gigantica* are found in Ethiopia and are transmitted by *Lymnaea truncatula* and *L. natalensis*, respectively and various reports indicated that it is a serious problem of livestock production in Ethiopia causing considerable economic losses. Recently, small-scale traditional irrigation scheme is expanding in many parts of the country. Hence, it is anticipated that implementation of irrigated agriculture is creating favorable habitat for intermediate snail vector thereby influence the occurrence of fasciolosis (Michael *et al.*, 2004).

Fasciolosis is a wide spread ruminant health problem and causes significant economic losses to the livestock industry in Ethiopia (Regassa *et al.*, 2012). In Ethiopia, *F. hepatica* and *F. gigantica* infections occur in areas above 1800 m and below 1200 m above sea level,

respectively which has been attributed to variations in the climatic and ecological conditions such as rainfall, altitude, and temperature and livestock management system. In between these altitude limits, both species coexists where ecology is conducive for both snail hosts, and mixed infections prevail (Yilma and Malones, 1998). Fasciolosis caused by *F. hepatica* and *F. gigantica*, is one of the most prevalent helminthes infections of ruminants in different parts of the world including Ethiopia (WHO, 1995; Okewole *et al.*, 2000). Fasciolosis causes a substantial economic loss which includes death, loss in carcass weight, reduction in milk yield, condemnation of affected liver, decline production and productive performances, exposure of animals to other diseases due to secondary complications and cost of treatment expenses. Both *F. hepatica* (high land) and *F. gigantica* (low land) type of liver flukes cause severe losses in Ethiopia where suitable ecological conditions for the growth and multiplication of intermediate host snails are available (Anne and Gary, 2006).

In Ethiopia, bovine fasciolosis is caused by the species *F. hepatica* and *F. gigantica* has been reported. They also a significant economic loss due to liver condemnation which affected by fasciola and moreover, the studies also showed that fasciolosis has higher economic significance on animal production and productivity. The economic losses due to fasciolosis throughout the world are enormous and these losses are associated with mortality, morbidity, reduced growth rate, condemnation of fluky, liver, increased susceptibility to secondary infections and expense due to control measures (Malone *et al* 1998). Therefore; the objective of this paper is collect a recent data and update information to review bovine fasciolosis in Ethiopia, and its current status and economic significance.

2. BOVINE FACIOLOSIS IN ETHIOPIA

2.1 Etiology

Bovine fasciolosis is caused by the two most important species of *F. hepatica* and *F. gigantica* which are commonly known as liver flukes. They are categorized in the phylum platyhelminthes, class trematoda, family fasciolidae and genus fasciola. *F. hepatica* mostly found predominantly in temperate areas & highlands that are in the cooler areas of high altitude in tropics & subtropics while *F. gigantica* predominates in tropical areas (Soulsby, 1982).

2.2 Epidemiology

2.2.1 occurrence, distribution and mode of transmission.

In general the distribution of Fasciolosis is worldwide. However, the distribution of *F. hepatica* is limited to temperate areas and high lands of tropical and subtropical regions (Soulsby, 1986). The distribution of *F. hepatica* is mostly encountered in temperate areas, and in cooler areas of high altitude in the tropics and subtropics, while *F. Gigantica* predominates in tropical areas. Snails are their intermediate hosts and amphibious snails of the genus *Lymnaea* species are widely distributed throughout the world and *L. trunculata* is the most common of them all (Soulsby, 1982 and Taylor *et al.*, 2007). The presence of snail intermediate host and therefore the distribution of the parasite are limited to geographic areas where appropriate snail species is present and is dependent on season (Mihreteab *et al.*, 2010).

Large numbers of metacercariae will usually be produced when there is optimal availability of suitable snail habitats, optimum temperatures and optimum moisture is present. For e.g. in Ethiopia the occurrence is closely associated to presence of suitable environmental conditions for the development of snails. Suitable snail habitats will include all areas where snails may survive in clear water or mud such as the edges of streams, ponds, rivers and veils (permanent natural habitats); or temporary man-made depressions filled with water (tractor tracks etc.) (Biniam *et al.*, 2012). A slightly acid environment may be more optimal. Temperature requirements are mean day/night temperatures of 10° C at which both the snails and the flukes will propagate. Below 5° C all activity will stop and above 15° C significant increase in both snails and fluke larval stages may be seen, with the optimum being 22 -26° C. Moisture levels are described as optimal when rainfall exceeds transpiration and when field saturation is achieved. Transmission occurs through ingestion of metacercariae during grazing of contaminated marshy areas (Urquhart *et al.*, 1996)

2.4 Pathogenesis

Pathogenesis varies according to the number of metacercariae ingested, the phase of parasitic development in the liver and the species of host involved. Acute hepatic fasciolosis is caused by the passage of young *F. hepatica* through the liver parenchyma. Clinical signs occur 5-6 weeks after the ingestion of large numbers of metacercariae. By this time, the migrating flukes are large enough to do substantial mechanical damage to the liver. Acute hepatic insufficiency and hemorrhage result quiescent spores of *Clostridium novyi* may become activated by

the anaerobic necrotic conditions created in the liver parenchyma by migrating *F. hepatica*, causing infectious necrotic hepatitis ('black disease') in sheep and cattle. This migration has also been thought to stimulate the development of occasional cases of bacillary hemoglobinuria in cattle (Radostits *et al.*, 2007).

Chronic hepatic fasciolosis develops only after the adult flukes establish in the bile ducts. Here results in calcification of the bile ducts and enlargement of the gallbladder. The calcified bile ducts often protrude from the liver surface cause cholangitis, biliary obstruction, fibrosis, and a leakage of plasma protein across the epithelium. Although this protein can be re-absorbed in the intestine, there is poor utilization and retention of nitrogen leading to hypoalbuminemia. There is also a loss of whole blood due to the feeding activities of the flukes. This exacerbates the hypoalbuminemia and eventually gives rise to anemia. Chronic infection may limit growth rate and feed conversion in growing heifers and growth rate in beef cattle. *F. hepatica* infection has been reported to increase the susceptibility of cattle to Salmonella Dublin and predispose to prolonged infection and fecal excretion. Aberrant migration of the flukes is more common in cattle and encapsulated parasites are often seen in the lungs (Taylor *et al.*, 2007).

2.5 Clinical signs

Acute fasciolosis not mostly common cattle however if the disease is observed clinically; it is manifested by; dullness, weakness, lack of appetite, pallor and edema of mucosa and conjunctivae, pain when pressure is exerted over the area of the liver. In heavy infections, where anemia and hypoalbuminaemia are severe, submandibular edema frequently occurs. With smaller fluke burdens, the clinical effect is minimal and the loss of productivity is difficult to differentiate from inadequate nutrition. It must be emphasized that diarrhoea is not a feature of bovine fasciolosis unless it is complicated by the presence of *Ostertagia* species (Radostits *et al.*, 2007). The chronic form is most important in cattle and seen in late winter that is early spring; and occurs as a result of ingesting moderate numbers (200–500) of metacercariae over longer periods of time; signs include anemia, unthriftiness, submandibular edema, and reduced milk production, but even heavily infected cattle may show no clinical signs but clinically, these are difficult to detect since the fluke burdens are usually low and anemia is not apparent. The main effects are a reduction in milk

yield and quality, particularly of the solids-non-fat component (Merck, 2005).

2.6 Diagnosis

This is based primarily on clinical signs, seasonal occurrence, prevailing weather patterns, and a previous history of fasciolosis on the farm or the identification of snail habitat. It can also be diagnosed by examination of faeces for fluke eggs (note: eggs of *Fasciola* are brownish yellow and eggs of Paramphistomidae are colorless) (Urquhart *et al.*, 1996).

2.7 Treatment

Triclabendazole is a compound specifically for use against *F. hepatica* in sheep (10 mg/kg) and cattle (12 mg/kg). Higher doses are required for the control of *F. gigantica* in buffalo.²⁶ It is highly effective against all stages of fluke from 2 days old in sheep and 2 weeks in cattle²⁷ and is the drug of choice in outbreaks of acute fluke disease. Albendazole is a broad-spectrum compound also active against nematodes and cestodes. It is effective against adult *F. hepatica* at a dose-rate of 7.5 mg/kg in sheep and 10 mg/kg in cattle. It is ovicidal and will kill any *F. hepatica* eggs present in bile ducts or the alimentary tract at the time of treatment. Netobimin (20 mg/kg) is metabolized to albendazole in the body and has similar activity against *F. hepatica* (Radostitis *et al.*, 2007).

At present there is only one drug, triclabendazole, which will remove the early immature (around 2 weeks of age in cattle) parenchyma stages. Apart from triclabendazole, the two drugs most commonly used for sub acute or chronic fasciolosis are nitroxylnil and oxcyclozanide also used to treat fasciolosis. Albendazole, ricobendazole and netobimin are also effective against adult fluke at an increased dosage rate. In lactating cows, where the milk is used for human consumption, the above drugs are either banned or have extended withdrawal periods. An exception is oxcyclozanide, which is licensed for use in lactating animals in many countries and has a milk-withholding time of up to 3 days. Resistance to flukicides is not a problem in cattle (Taylor *et al.*, 2007).

2.8 Control

Preventive measures are required in endemic areas as fasciolosis can cause death without warning or significant production losses. An integrated strategic approach is more cost beneficial than reliance on

routine dosing and is less likely to induce anthelmintics resistance, but requires detailed knowledge of the local epidemiological cycle. Reduction of pasture contamination with metacercariae will reduce future risk. This can be done by preventing the snails from becoming infected with *F. hepatica* or by diminishing the size of the snail population. To achieve the first objective, adult flukes should be eliminated from the bile ducts of all grazing stock in spring and early summer. This prevents egg excretion and minimizes the numbers of snail-seeking miracidia at this crucial stage in the epidemiological cycle. There may however be wildlife sources of *F. hepatica* eggs which cannot be controlled in this way. Snail numbers can be reduced by restricting the size of their habitat. This can be done, where feasible, by draining boggy areas and by making sure that ditches, land drains; water troughs etc. are well maintained (Radiostis *et al.*, 2007).

3.STATUS OF BOVINE FACIOLOSIS IN ETHIOPIA AND ITS ECONOMIC SIGNIFICANCE

3.1 Status of bovine faciolosis in Ethiopia

In Ethiopia, both *F. hepatica* and *F. gigantica* have been reported to exist in many parts of the country. The prevalence of bovine faciolosis based on coproscopy result varies from 11.5% in buno province to 87% in Debre Berhan. Abattor studies have also reported up to 88.5% prevalence in Debre Berhan. Faciolosis in sheep and cattle in animal's results that shows low productivity (low weight gain, low milk production etc.). Also in many countries including Ethiopia, livers from animals infected with *F. hepatica* and *F. gigantica* are condemned as unsuitable for human consumption (Zerihun, 2006). The prevalence of bovine faciolosis in Ethiopia has been reported by different researchers at different areas. Since those areas are suitable to the snail intermediate host (Yosef *et al.*, 2014). The presence of fasciolosis due to *F. hepatica* and *F. gigantica* at abattoir surveys in some parts of the country has long been known and its prevalence and economic significance have been reported by several workers (Rahmatho *et al.*, 2010; Biniam *et al.*, 2012; Yosef *et al.*, 2014; Regassa *et al.*, 2012; Mihreteab *et al.*, 2010; Alula *et al.*, 2013; Ibrahim *et al.*, 2009)

Table 1 the prevalence of bovine faciolosis in different areas of Ethiopia

Study area	Prevalence (%)	<i>F. hepatica</i> (%)	<i>F. gigantica</i> (%)	Mixed infection (%)	References
Hawassa Municipal abattoir, southern Ethiopia	28.6	58.9	10.6	14.7	(Rahmatho <i>et al.</i> , 2010)
in and around Woreta, Northwestern Ethiopia	41.41	NR	NR	NR	(Biniam <i>et al.</i> , 2012)
in and Around Bedelle District, Ethiopia	20.8	NR	NR	NR	(Yosef <i>et al.</i> , 2014)
Bishoofu Municipal Abattoir, Central Ethiopia	21.6	18.3	6.4	3.1	(Regassa <i>et al.</i> , 2012)
at Adwa Municipal Abattoir, North Ethiopia	32.3	13.9	7.7	6	(Mihreteab <i>et al.</i> , 2010)
in Nekemte Municipal abattoir	21.9	14.1	5.2	2.6	(Alula <i>et al.</i> , 2013)
At Kombolcha Industrial Abattoir, Ethiopia	39.6	24.3	7.14	5	(Ibrahim <i>et al.</i> , 2009)

NR: not recorded and the percentage of *F. hepatica* *F. gigantica* are among positive animals

3.2 Economic significance

Bovine fasciolosis is an important economical disease in Ethiopia. In line to the economic importance of bovine fasciolosis in Ethiopia is associated with loss of productivity (indirect loss) and due to condemnation of infected liver (direct loss). This reported by different researchers in different places and at different time (Tolosa and Tigre, 2006; Shiferaw *et al.*, 2011; Zeleke *et al.*, 2014; Abunna *et al.*, 2010; Yohannes *et al.*, 2012).

Table 2 direct and indirect economic loss of fasciolosis

Study area	Direct loss(due to liver condemnation)/annum	Indirect loss (due to loss of productivity)/ETB/ annum	Total/ETB	Reference
Jimma municipal abattoir	54,063.34 ETB	NR	NR	(Tolosa and Tigre, 2006)
In and Around Assela	37, 767.6 ETB	660, 933 ETB	698,700.6 ETB	(Shiferaw <i>et al.</i> , 2011)
Mettu Municipal Abattoir	47,570.00 ETB	466,150.00 ETB	513,720ETB	(Zeleke <i>et al.</i> , 2014)
Soddo municipal abattoir	4000 USD	NR	NR	(Abunna <i>et al.</i> , 2010)
Mekelle municipal abattoir	2245 USD	NR	NR	(Yohannes <i>et al.</i> , 2012)

NR, not recorded; USD, United States dollar; ETB, Ethiopian birr

4. CONCLUSION AND RECOMMENDATIONS

In general it could be concluded that Fasciolosis was one of major problem for livestock development in Ethiopia and closely linked to the presence of biotypes suitable to the development of snail intermediate host. Both *L. truncatula* and *L. natalensis* which are the intermediate hosts of *F. hepatica* and *F. gigantica* respectively which causes bovine fasciolosis are found in Ethiopia. The species of *F. hepatica* is predominantly occurring than *F. gigantica* and sometimes mixed infection also occur. There is highly significant economic loss that accounts in thousands birr due to direct and indirect effect of fasciolosis in Ethiopia. Therefore; based on the above conclusions the following recommendations are provided:

- ❖ Strategic anthelmintic treatment with appropriate fluckicide drug should be practiced
- ❖ A combination of control measures included drainage, fencing and snails had to be used to ensure a satisfactory and degree of control in the long run.
- ❖ Seasonal control applied.

5. REFERENCES

1. Abunna F., Asfaw L., Megersa B., Regassa A. (2010): *Bovine fasciolosis: coprological, abattoir survey and its economic impact due to liver condemnation at, Southern Ethiopia, Trop. Anim. Health Prod.* **42**(2)
2. Alula P., Addisu K., Amanuel W. (2013): Prevalence and economic significance of bovine fasciolosis in Nekemte Municipal abattoir, *J. Vet. Med, and Ani. Health* vol.5
3. Anne MZ, Gray MC. (2006). *Veterinary clinical Parasitology*, 7thed. American Association of the Proctolohist. pp. 185-210.
4. Biniam T., Hanna A. and Sisay G., (2012): Study on coprological prevalence of bovine fasciolosis in and around Woreta, Northwestern Ethiopia: *J. Vet. Med. and Ani. H.* **4**(7), 89-92.
5. Ibrahim N., Wasihun P., Tolosa T. (2009): Prevalence Of Bovine Fasciolosis And Economic Importance Due To Liver Condemnation At Kombolcha Industrial Abattoir, Ethiopia. *Int. J. Vet. Med.* **8** (2).

6. Kahn, M.C. (2005): The Merck Veterinary Manual. 9th.ed.USA: MERCK&CO; INC
7. Malone J B., Gommers R., Hansen J., Yilma J M., Slingenberg J., Snijders F., Nachtet O F and Ataman E.(1998): A Geographic Information System on the potential Distribution and abundance of *Fasciola hepatica* and *F. gigantica* in East Africa based on food and agriculture organization databases, Elev. Vet Parasitology, 78 87-101.
8. Menkir, MS., Ugglä, A.and Waller, PJ. (2007): Prevalence and seasonal incidence of nematode parasites and fluke infections of sheep and goats in eastern Ethiopia. *Trop. Ani. Heal. andProd.* **39(7)**:pp521-531.
9. Michael, A., P. Beyene, J. Yilma, P. Don, S. Yoseph, T. Girma and M. Mulugeta, 2004. Infection prevalence of ovine Fasciolosis in small scale irrigation schemes along the upper Awash River Basin. *Eth. Vet. J.*, 9: 19-26.
10. Okewole EA, Ogundipe GAT, Adejinmi JO, Olaniyan AO (2000). Clinical evaluation of three chemo prophylactic regimes against bovine helminthosis in a *Fasciola* endemic farm in Ibadan, Nigeria. *Israel. J. Vet. Med.* 56(1):15-28.
11. Radostits, O.M., C.C. Gay, K.W. Hinchcliff and P.D. Constable, 2007. *Veterinary Medicine: A Textbook of the Diseases of Cattle, Horses, Sheep, Pigs and Goats*, 10th ed. Elsevier Health Sciences, Philadelphia, PA, USA, pp1576-1579
12. Rahmeto A., Fufa A., Mulugeta B., Solomon M., Bekele M., Alemayehu R. (2010): Fasciolosis: Prevalence, financial losses due to liver condemnation and evaluation of a simple sedimentation diagnostic technique in cattle slaughtered at Hawassa Municipal abattoir, southern Ethiopia. *Eth. Vet. J.* **14** (1), 39-51
13. Regassa A., Woldemariam T., Demisie S., Moje N., Ayana D. and Abunna F., 2012: Bovine Fasciolosis: Coprological, Abattoir Survey and Financial Loss Due to Liver Condemnation in Bishooftu Municipal Abattoir, Central Ethiopia. *Eur. J. Bio. Sci.* 4 (3): 83-90,
14. Shiferaw M., Feyisa B., Ephrem T. (2011): Prevalence of Bovine Fasciolosis and its Economic Significance in and Around Assela, *Eth. Glo. J. Med. Res.*, **4** (XI)
15. Soulsby E. J. L. (1982): *Helminthes, Arthropods and Protozoa of Domesticated Animals*, Seventh Ed. Balliere Tindall, London, UK. Pp 40-52.
16. Taylor M. A., Coop R. L., Wall R. L., (2007): *Veterinary parasitology*, Third ed., Blackwell, UK. Pp 243-250.
17. Tolosa T., Tigre W. (2006): The Prevalence and Economic Significance of Bovine Fasciolosis at Jimma Abattoir, Ethiopia. *Int. J. Vet. Med.* **3**(2).
18. Troncy PM (1989). *Helminthes of livestock and poultry in Tropical Africa*, Manual of tropical veterinary parasitology. CAB int. UK. pp. 63-73.
19. Urquhart G M., Duncan J L., Armour J., Dunn A M., Jennings (1996): *Veterinary Parasitology*. Second Ed. Blackwell Science, UK. Pp 103-113.
20. Walker, SM, Makundi, AE, Namuba, FV, Kassuku, AA, Keyyu, J, Hoey, EM, Prodohl, P, Stothard, JR and Trudgett, A. (2008): The distribution of *Fasciola hepatica* and *Fasciola gigantica* within southern Tanzania-constraints associated with the intermediate host. *Parasitology* **135(4)**:495-503.
21. WHO (1995). *Control of food bore Trematodes infections*, Technical Report Series 849:61-63.
22. Yohannes E., Abebaw G., Mahendra P., Gebererufael G. (2012): Prevalence of bovine fasciolosis, amplitude of liver condemnation and its economic impact in municipal abattoir of Mekelle, Ethiopia, *Int. J. Live. Res.* **2**(2).
23. Yosef M., Yosef D. and Nuraddis I. (2014): Prevalence of Bovine Fasciolosis in and Around Bedelle District, Ethiopia: IDOSI Publications, *Acta Para. Glob.* **5** (2):
24. Zeleke M. A., Mengistu G., Tsegaye T. (2014): Economic Significance of Fasciolosis at Mettu Municipal Abattoir, Southwest Ethiopia, *J. Adv. Vet. Res.*, **4**(2)
25. Zerihun, A. (2006): Ruminant fasciolosis: Studies on the clinical occurrence, coprology, malcology, and abattoir survey in Debre Birhan and surrounding areas. Msc. Thesis, F.V.M., A.A.U., Debrezeit.