

# Phytochemical screening on calli of *Fagonia indica* and *Fagonia bruguieri* Dc.

EMAN, A. ALAM

Botany Department, National Research Centre, Dokki, Giza, Egypt  
[Eman200980@hotmail.com](mailto:Eman200980@hotmail.com)

**Abstract:** Tissue culture technique was used to produce calli from the two wild economic plants (*Fagonia indica* Burm f var. *indica* and *Fagonia bruguieri* Dc.). MS medium supplemented with 5 mg/l Kinetin + 1 mg/l NAA induced calli from *F. indica* stem segments explants. In this regard, *Fagonia bruguieri* leaf explants can induce calli on MS medium supplemented with 5 mg/l Kinetin + 1 mg/l NAA also, while terminal bud explants can induce calli on this medium and MS medium supplemented with 5 mg/l Kinetin + 1 mg/l 2,4-D. While MS medium supplemented with 6 mg/l Kinetin + 2 mg/l NAA was the most suitable medium for growth of these calli of the two plants. Phytochemical screening on both calli of *F. indica* and *Fagonia bruguieri* revealed a variation in the presence/ amount of carbohydrates and / or glycosides, saponins, tannins, unsaturated sterols and/or triterpenoids, alkaloids, cardiac glycosides, cyanogenic glycosides, flavonoids, coumarins, chlorides and sulphates.

[EMAN, A. ALAM. Phytochemical screening on calli of *Fagonia indica* and *Fagonia bruguieri* Dc. New York Science Journal 2010;3(12):158-164]. (ISSN: 1554-0200). <http://www.sciencepub.net/newyork>.

**Key words:** *Fagonia*, *F. indica*, *Fagonia bruguieri*, phytochemical screening, tissue culture, calli.

## 1. Introduction

Genus *Fagonia* is represented in Egypt by 18 species (Tackholm, 1974), but it was represented by 15 species in Boulos, 2000. *Fagonia* species were extensively studied by many workers regarding their medicinal uses, since these plants were antitumor, antioxidant, analgesic, astringent, febrifuge and prophylactic against small-pox agents, species of *Fagonia* were also used for the treatment of cancer in the indigenous system, fever, asthma, urinary discharges, toothache, stomach troubles and kidney diseases (Ahsan *et al.*, 2007 and Satpute *et al.*, 2009). Species of *Fagonia* have been found to contain saponins (Abdel- Khalik *et al.*, 2001), alkaloids (Sharawy and Alshammari, 2009), terpenoids (Perrone *et al.*, 2007), sterols (Shoeb *et al.*, 1994), flavonoids (Ibrahim *et al.*, 2008), proteins and amino acids (Sharma *et al.*, 2010), coumarins Zhang *et al.*, 2008), trace elements (Fatima *et al.*, 1999). Tissue culture technique was used to cultivate different genera of Zygophyllaceae for regeneration purposes such as *Zygophyllum xanthoxylon* (Bunge) (Sun, 2008), to produce more biologically active compounds such as ascorbic acid from callus of *Fagonia cretica* (Kapoor, 2002), alkaloids, saponins, flavonoids and phenolic compounds of antibacterial activity more than the intact plant (Eman *et al.*, 2010) diosgenin from callus of *Balanites aegyptiaca* (Gour and Kant, 2006) and beta -carboline and serotonin alkaloids and fatty acids from callus of *Peganum harmala* (Piacetini *et al.*, 2004).

The present study aims to produce calli from the two wild economic plants (*Fagonia indica* Burm f var. *indica* and *Fagonia bruguieri* Dc.) using tissue culture technique for the first time and to make a phytochemical screening on the chemical constituents of these calli.

## 2. Material and Methods

### Plant materials

Samples of *F. indica* Burm f. var. *indica* (= *F. parviflora* Boiss.) and *Fagonia bruguieri* Dc. were collected from Quatamia - Suez desert road (150 Km away from Suez City). All the samples were authenticated by comparison with voucher specimens in the herbarium of Botany Department, Faculty of Science, Ain Shams University, Cairo, Egypt, where voucher specimens were deposited.

### Methods

#### A- Tissue culture study

##### 1- Callus induction

This experiment was carried out to study the effect of different levels of 2,4-D, NAA and Kinetin on calli induction of different explants obtaining from *F. indica* Burm f. var. *indica* (= *F. parviflora* Boiss.) and *Fagonia bruguieri* Dc. (stem, leaf and terminal bud). Explants were surface sterilized by immersion in 70 % ethanol for 30-60 seconds, then

soaked in 50% of commercial Clorox for 15-20 minutes, then washed with sterile distilled water "3 times" (Hoda, 1994). The sterilized explants were aseptically transferred to sterilized MS medium (Murashige and Skoog, 1962); supplemented with

3% sucrose, 1% agar and either 5 mg/l Kinetin + 1 mg/l 2,4-D (1) or 5 mg/l Kinetin + 1 mg/l NAA (2). Calli cultures were incubated at  $25 \pm 2^\circ\text{C}$  for six weeks (Eman *et al.*, 2010).

## 2-Callus growth

Calli of *F. indica* (stem segments explants) and *Fagonia bruguieri* ( leaf and terminal bud explants) can be maintained by subculturing on MS medium supplemented with different concentrations of auxins and cytokinins (Eman *et al.*, 2010) as follows:

| Media | Hormones                         |
|-------|----------------------------------|
| a     | 6 mg/ l Kinetin+<br>2 mg/l NAA   |
| b     | 6 mg/ l Kinetin+<br>2 mg/l 2,4-D |
| c     | 5 mg/ l Kinetin +<br>1 mg/l NAA  |
| d     | 5 mg/ l Kinetin+<br>1 mg/l 2,4-D |
| e     | 5 mg/l Kinetin +<br>2 mg/l NAA   |
| f     | 5 mg/l Kinetin +<br>2 mg/l 2,4-D |

Calli cultures were incubated at  $25 \pm 2^\circ\text{C}$  for four weeks. Weight of calli was taken as an indicator of callus growth on each medium as follows:

|     |  |
|-----|--|
| -   | no growth of calli were<br>observed on this medium |
| +   | $\geq 1$ g   |
| ++  | $\geq 2$ g   |
| +++ | $\geq 5$ g   |

## 3. Results and Discussion

### A- Tissue culture study

#### 1- Calli induction

MS medium supplemented with 5 mg/l Kinetin + 1 mg/l NAA induced calli from *F. indica* stem segments explants. In this regard, *Fagonia bruguieri* leaf explants can induce calli on MS medium supplemented with 5 mg/l Kinetin + 1 mg/l NAA also, while terminal bud explants can induce calli on both this medium and MS medium supplemented with 5 mg/l Kinetin + 1 mg/l 2,4-D (Photos: 1- 4).



Photo (1): Callus induction of *F. indica* stem explants on MS+5mg/l Kinetin+1mg/l NAA.



Photo (2): Callus induction of leaf of *F. bruguieri* on MS+5mg/l Kinetin+1mg/l NAA.



Photo (3): Callus induction of terminal bud of *F. bruguieri* on MS+5mg/l Kinetin+1mg/l 2,4-D



Photo (4): Callus induction of terminal bud of *F. bruguieri* on MS+5mg/l Kinetin+1mg/l NAA.

## 2- Callus growth of *F. indica* (stem segments explants) and *Fagonia bruguieri* ( leaf and terminal bud explants)

Data in Table (1) and Photos (5-7) revealed that, callus of *F. indica* stem segments explants grew on media number a, d and e only, with special reference to that on medium number a. While *Fagonia bruguieri* leaf explants calli can grow on media number b only, terminal bud explants of *Fagonia bruguieri* can grow on media number a, c, d and f only, with special reference to that on medium number a.

### B-Preliminary phytochemical screening on both callus of *F. indica* hypocotyle explants and the intact hypocotyle:

Preliminary phytochemical screening (Table: 2) [on both calli of *F. indica* stem segments explants that grew on media number a, d and e ( callus 1a, 1d, 1e), *Fagonia bruguieri* leaf explants calli that grew on media number b ( callus 2b), terminal bud explants of *Fagonia bruguieri* that grew on media number a, c, d and f ( callus 3a, 3c, 3d, 3f)] revealed that, there were variations in the presence/ amount of saponins, flavonoids and chlorides within different calli of *F. indica* stem explants that grew on different media. Regarding *F.bruguieri* terminal bud explants there were variations also in these phytochemicals in addition to Sulphates, irodoids and cyanogenic glycosides within different calli that grew on different media, and there were also variations in these phytochemicals within calli of terminal bud and leaf explants of *Fagonia bruguieri*. So we can conclude that there were two factors controlled the *in vitro* production of phytochemicals; explants used and concentrations and types of plant growth regulators added to the medium (such as concentrations and types of auxins " NAA or 2, 4-D" in our study). So we can use these factors for directing the tissues to produce the needed phytochemicals.

**Table (1): Qualitative estimation of calli of *F. indica* (stem segments explants) and *Fagonia bruguieri* ( leaf and terminal bud explants) .**

| Media | Weight of callus of <i>F. indica</i> stem segments explants | Weight of callus of <i>F. bruguieri</i> leaf explants | Weight of callus of <i>F. bruguieri</i> terminal bud explants |
|-------|---|---|---|
| a     | +++   | -   | +++   |
| b     | -   | +   | -   |
| c     | -   | -   | ++  |
| d     | ++  | -   | ++  |
| e     | +   | -   | -   |
| f     | -   | -   | +   |




Photo (5): Callus growth of stem segments explants of *F. indica* on medium number (a).



Photo (6): Callus growth of leaf explants of *F. bruguieri*. on medium number (b) .



Photo (7): Callus growth of terminal bud explants of *F. bruguieri* on medium number (a).

**Table (2): Preliminary phytochemical screening on both calli of *F. indica* (stem segments explants) and *Fagonia bruguieri* (leaf and terminal bud explants) on different media.**

| Experiment                                    | Calli of <i>F. indica</i> stem explants |    |    | Calli of <i>Fagonia bruguieri</i> (leaf and terminal bud explants) |    |    |    |    |
|---|---|----|----|--|----|----|----|----|
|   | 1a                                      | 1d | 1e | 2b   | 3a | 3c | 3d | 3f |
| 1- Carbohydrates and / or Glycosides          | +                                       | +  | +  | +  | +  | +  | +  | +  |
| 2- Saponins                                   | -                                       | -  | +  | -  | +  | ++ | -  | -  |
| 3- Tannins                                    | +                                       | +  | +  | +  | ++ | +  | +  | +  |
| 4- Unsaturated sterols and / or Triterpenoids | +                                       | +  | +  | +  | +  | +  | +  | +  |
| 5- Alkaloids                                  | +                                       | +  | +  | +  | +  | +  | +  | +  |
| 6- Cardiac glycosides                         | +                                       | +  | +  | +  | +  | +  | +  | +  |
| 7- Cyanogenic glycosides                      | +                                       | +  | +  | +  | -  | +  | +  | +  |
| 8- a- Chlorides                               | -                                       | -  | +  | -  | +  | -  | -  | -  |
| 8- b- Sulphates                               | +                                       | +  | +  | +  | -  | +  | +  | +  |
| 9- Irodoids                                   | -                                       | -  | -  | -  | -  | -  | +  | +  |
| 10-Flavonoids                                 | -                                       | -  | +  | -  | +  | -  | -  | -  |
| 11- Coumarins                                 | +                                       | +  | +  | +  | +  | +  | +  | +  |
| 12- Anthraquinones                            | -                                       | -  | -  | -  | -  | -  | -  | -  |

**Corresponding author:****Eman, A. Alam**Botany Department, National Research Centre,  
Dokki, El- Behoos Street, Giza, Egypt.[Eman200980@hotmail.com](mailto:Eman200980@hotmail.com)**Mobile: 0179469341-0187180968 (Egypt).****References**

1. Tackholm, V. : Students flora of Egypt 1974. Cairo University Press, Cairo, Egypt: 302-307.
2. Boulos, L. : Flora of Egypt 2000. Alhadara Publishing, Egypt: 12-18.
3. Ahsan, H.; Muhammad, Z. and Bushra, M.: Cytotoxic and antitumor potential of *Fagonia cretica* L. Turkish Journal of Biology 2007; 31(1): 19-24.
4. 2. Satpute, R.M; Kashyap, R.S.; Deopujiari, J.Y.; Taori, G.M. and Dagainawala, H.F.: Protection of PC12 cells from chemical ischemia induced oxidative stress by *Fagonia arabica*. Food and chemical toxicology 2009; 47(11): 2689-2695.
5. Abdel-Khalik, S.M.; Miyase, T.; Melek, F.R. and El-Ashaal, H.A.: Further saponins from *Fagonia cretica*. Dipharmazie 2001; 56(3): 247-250.
6. Sharawy, S.M. and Alshammari, A.M.: Checklist of Poisonous Plants and Animals in Aja Mountain, Ha'il Region, Saudi Arabia. Australian Journal of Basic and Applied Sciences 2009; 3(3): 2217-2225.
7. Perroni, A.; Masullo, M.A.; Basarello, C.; Hamed, A.I.; Belisario, M.A.; Pizza, C. and Piacente, S.: Journal of natural products 2007; 70(4): 584-588.
8. Shoeb, H.A.; Sharada, M.M.; El-Sayed, L.A.R. and El-Wakeel, E.: Triterpenoid and Sterol glycosides from *Fagonia arabica* L.. Al-Azhar Journal of Pharmaceutical Sciences 1994;13: 41-48.
9. Ibrahim, L.F.; Kawashty, S.A.; El-Hagrassy, A.M.; Nassar, M.L. and Mabry, T.j.: Anew kaempferol triglycoside from *Fagonia taekholmiana*: cytotoxic activity of its extracts. Carbohydrate research 2008; 343(1): 155-158.
10. Sharrma, S; Gupta, V. and Sharma G.: Phytopharmacology of *Fagonia Indica* (L): A Review. Journal of Natura Conscientia 2010; 1(1): 143-147.
11. Zhang, W.; Krohn, K.; Draeger, S. and Schulz, B.: Bioactive isocoumarins isolated from the endophytic fungus *Microdochium bolleyi*. Journal of natural products 2008; 71(6): 1078-1081
12. Fatima, K.; Khaula, S.; Kalhor, M.A.; Muhammad, Q. and Yasmeen, B.: Trace

- elements in indigenous medicinal plants (*Rhazya stricta*, *Vinca rosea* and *Fagonia cretica*). *Phytochemistry* 1999; 42(4): 182-1831.
13. Sun .Rapid clonal propagation of *Zygophyllum xanthoxylon* (Bunge) Maxim., an endangered desert forage species.*In-Vitro-Cellular-and-Developmental-Biology-Plant* 2008;44(5):.396-400.
  14. Kapoor, B.B.S.. Production of endogenous ascorbic acid from tissue cultures of *Fagonia cretica* L. . *Journal of Phytological Research*2002; 15(1): 111-112.
  15. Eman,A. Alam; Gehan, H. Amin; Yassin, M . ElAyouty and Mohamed, S. Abdel-Hady. Chemical Composition and Antibacterial Activity Studies on Callus of *Fagonia arabica* L.. *Academia Arena* 2010;2(12):91-106.
  16. Gour, V.S. and Kant, T.. Optimized callus production in *Balanites aegyptiaca* (L.) Del. . *Journal of Phytological Research* 2006; 19(1):53-57.
  17. Piacetini, M.P.; Piatti, E. and Faternale. Phospholipase C-dependent phosphoinositide breakdown induced by ELF-EMF in *Peganum harmala* callus . *Biochemimie (Paris)* 2004; 86(4-5):.343-349.
  18. Hoda, A.M.. An attempt for producing callus from tomato plants resistant to salinity. M.Sc. Thesis 1994, Vegetables and Horticulture Department, Faculty of Agriculture, Zagazig University, Egypt: 36.
  19. Murashige, T. and Skoog, F.. A revised medium for rapid growth and bioassays with Tobacco tissue cultures. *Physiology of Plants* 1962; 15: 473-479.
  20. Mabry, T.T.; Markhan, K.R.and Thomas, M.B.: The systemic identification of flavonoids 1970. Springer, Verlag, New York: 46-54.
  21. Farnsworth ,N.R. ;Fong ,H.H. ;Blomster ,R.N. and Draus ,F.G.: Studies on *Vinca major* (Apocycaceae). *Journal of Pharmaceutical Science* . 1969; 51(3): 217-224.
  22. Trease, G.T. and Evans, W.C.: Text book of Pharmacognsy 1978 , Bailliere Tindall and Cox, London , 11<sup>th</sup> Ed.:536.
  23. Shellard, E.J.: Practical plant chemistry. Pitman 1957, Medicinal publishing Co., LTD, London: 53-54.
  24. Hungund, B.L. and Pathak, C.H.: USDA forest 1971, Service Research Paper, NE: 201.
  25. Stank, J.; Cerny, M.; Kocoursk, J. and Pacok, J.: The monosaccharides 1963, Publishing House of the Czechoslovak, Academy of Sciences, Prague: 22-100.
  26. Weifferring, J.H.: Aucubinartige glucoside und verwandte heteroside als systematische merkmale .*Phytochemistry* 1966; 5: 1053.
  27. Feigl , F.: Spot tests in organic analysis 1960. Elsevier Publishing Co. , New York ,6<sup>th</sup> ed.: 29-59.
  28. Islam, A.M.; Hassan, E.A. and Hannout, I.B.: Manual of Practical Chemistry 1993, Dar Al-Maaref, Egypt, 2<sup>nd</sup> ed.: 19-39.
  29. Claus , E.P.: Pharmacognosy 1967, Henery Krimpton , London , 5<sup>th</sup> ed.: 168.
  30. Schmidt, J.: Textbook of Organic Chemistry 1964. Olivar and Poyed ed., London: 673.
  31. Balbaa, S.I.; Sayed, H.H. and Ashgan, Y.Z.: Medicinal plant constituent 1981, General organization for university and school books,3<sup>rd</sup> ed :190-255.
  32. Afifi, M.: Phrmacological studies on some genera of Polygonaceae and cucurbitaceae grown in Egypt. Ph.D. Thesis 1972, Pharmacology Department, Faculty of Pharmacy, Cairo University, Egypt: 68.

11/22/2010