

Complex Hyperbolic Functions And Fermat's Last Theorem

Jiang Chunxuan

Institute for Basic Research, Palm Harbor, FL 34682-1577, USA

And: P. O. Box 3924, Beijing 100854, P. R. China

jiangchunxuan@sohu.com, cxjiang@mail.bcf.net.cn, jcxuan@sina.com, Jiangchunxuan@vip.sohu.com,
jcxxxx@163.com, liukxi@public3.bta.net.cn

Abstract: In 1637 Fermat wrote: "It is impossible to separate a cube into two cubes, or a biquadrate into two biquadrates, or in general any power higher than the second into powers of like degree: I have discovered a truly marvelous proof, which this margin is too small to contain."

This means: $x^n + y^n = z^n (n > 2)$ has no integer solutions, all different from 0 (i.e., it has only the trivial solution, where one of the integers is equal to 0). It has been called Fermat's last theorem (FLT). It suffices to prove FLT for exponent 4. and every prime exponent P . Fermat proved FLT for exponent 4. Euler proved FLT for exponent 3.

[Jiang Chunxuan. **Complex Hyperbolic Functions And Fermat's Last Theorem**. *Rep Opinion* 2017;9(4s):27-32]. ISSN 1553-9873 (print); ISSN 2375-7205 (online). <http://www.sciencepub.net/report>. 5. doi:[10.7537/marsroj0904s17.05](https://doi.org/10.7537/marsroj0904s17.05).

Key words: complex hyperbolic, Euler proof.

2000 mathematics subject classification 11D41.

In this paper using the complex hyperbolic functions we prove FLT for exponents $3P$ and P , where P is an odd prime. The proof of FLT must be direct. But indirect proof of FLT is disbelieving.

In 1974 Jiang found out Euler formula of the cyclotomic real numbers in the cyclotomic fields

$$\exp\left(\sum_{i=1}^{n-1} t_i J^i\right) = \sum_{i=1}^n S_i J^{i-1} \quad (1)$$

where J denotes a n th root of unity, $J^n = 1$, n is an odd number, t_i are the real numbers.

S_i is called the complex hyperbolic functions of order n with $n-1$ variables [1-7].

$$S_i = \frac{1}{n} \left[e^A + 2 \sum_{j=1}^{\frac{n-1}{2}} (-1)^{(i-1)j} e^{B_j} \cos\left(\theta_j + (-1)^j \frac{(i-1)j\pi}{n}\right) \right] \quad (2)$$

where $i=1,2,\dots,n$;

$$A = \sum_{\alpha=1}^{n-1} t_\alpha, \quad B_j = \sum_{\alpha=1}^{n-1} t_\alpha (-1)^{\alpha j} \cos \frac{\alpha j \pi}{n},$$

$$\theta_j = (-1)^{j+1} \sum_{\alpha=1}^{n-1} t_\alpha (-1)^{\alpha j} \sin \frac{\alpha j \pi}{n}, \quad A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j = 0 \quad (3)$$

(2) may be written in the matrix form

$$\begin{bmatrix} S_1 \\ S_2 \\ S_3 \\ \dots \\ S_n \end{bmatrix} = \frac{1}{n} \begin{bmatrix} 1 & 1 & 0 & \dots & 0 \\ 1 & -\cos \frac{\pi}{n} & -\sin \frac{\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{2n} \\ 1 & \cos \frac{2\pi}{n} & \sin \frac{2\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \cos \frac{(n-1)\pi}{n} & \sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \begin{bmatrix} e^A \\ 2e^{B_1} \cos \theta_1 \\ 2e^{B_1} \sin \theta_1 \\ \dots \\ 2 \exp B_{\frac{n-1}{2}} \sin \theta_{\frac{n-1}{2}} \end{bmatrix} \quad (4)$$

where $(n-1)/2$ is an even number.

From (4) we have its inverse transformation

$$\begin{bmatrix} e^A \\ e^{B_1} \cos \theta_1 \\ e^{B_1} \sin \theta_1 \\ \dots \\ \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & -\cos \frac{\pi}{n} & \cos \frac{2\pi}{n} & \dots & \cos \frac{(n-1)\pi}{n} \\ 0 & -\sin \frac{\pi}{n} & \sin \frac{2\pi}{n} & \dots & \sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & -\sin \frac{(n-1)\pi}{2n} & -\sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \begin{bmatrix} S_1 \\ S_2 \\ S_3 \\ \dots \\ S_n \end{bmatrix} \quad (5)$$

From (5) we have

$$\begin{aligned} e^A &= \sum_{i=1}^n S_i, & e^{B_j} \cos \theta_j &= S_1 + \sum_{i=1}^{n-1} S_{1+i} (-1)^{ij} \cos \frac{ij\pi}{n} \\ e^{B_j} \sin \theta_j &= (-1)^{j+1} \sum_{i=1}^{n-1} S_{1+i} (-1)^{ij} \sin \frac{ij\pi}{n} \end{aligned} \quad (6)$$

In (3) and (6) t_i and S_i have the same formulas. (4) and (5) are the most critical formulas of proofs for FLT. Using (4) and (5) in 1991 Jiang invented that every factor of exponent n has the Fermat equation and proved FLT [1-7] Substituting (4) into (5) we prove (5).

$$\begin{bmatrix} e^A \\ e^{B_1} \cos \theta_1 \\ e^{B_1} \sin \theta_1 \\ \dots \\ \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} = \frac{1}{n} \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & -\cos \frac{\pi}{n} & \cos \frac{2\pi}{n} & \dots & \cos \frac{(n-1)\pi}{n} \\ 0 & -\sin \frac{\pi}{n} & \sin \frac{2\pi}{n} & \dots & \sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & -\sin \frac{(n-1)\pi}{2n} & -\sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \times$$

$$\begin{aligned}
 & \begin{bmatrix} 1 & 1 & 0 & \dots & 0 \\ 1 & -\cos \frac{\pi}{n} & -\sin \frac{\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{2n} \\ 1 & \cos \frac{2\pi}{n} & \sin \frac{2\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \cos \frac{(n-1)\pi}{n} & \sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \begin{bmatrix} e^A \\ 2e^{B_1} \cos \theta_1 \\ 2e^{B_1} \sin \theta_1 \\ \dots \\ 2 \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} \\
 &= \frac{1}{n} \begin{bmatrix} n & 0 & 0 & \dots & 0 \\ 0 & \frac{n}{2} & 0 & \dots & 0 \\ 0 & 0 & \frac{n}{2} & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \frac{n}{2} \end{bmatrix} \begin{bmatrix} e^A \\ 2e^{B_1} \cos \theta_1 \\ 2e^{B_1} \sin \theta_1 \\ \dots \\ 2 \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} \\
 &= \begin{bmatrix} e^A \\ e^{B_1} \cos \theta_1 \\ e^{B_1} \sin \theta_1 \\ \dots \\ \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix}, \tag{7}
 \end{aligned}$$

where $1 + \sum_{j=1}^{n-1} (\cos \frac{j\pi}{n})^2 = \frac{n}{2}$, $\sum_{j=1}^{n-1} (\sin \frac{j\pi}{n})^2 = \frac{n}{2}$.
 From (3) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j) = 1 \tag{8}$$

From (6) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j) = \begin{vmatrix} S_1 & S_n & \dots & S_2 \\ S_2 & S_1 & \dots & S_3 \\ \dots & \dots & \dots & \dots \\ S_n & S_{n-1} & \dots & S_1 \end{vmatrix} = \begin{vmatrix} S_1 & (S_1)_1 & \dots & (S_1)_{n-1} \\ S_2 & (S_2)_1 & \dots & (S_2)_{n-1} \\ \dots & \dots & \dots & \dots \\ S_n & (S_n)_1 & \dots & (S_n)_{n-1} \end{vmatrix}, \tag{9}$$

where $(S_i)_j = \frac{\partial S_i}{\partial t_j}$ [7].
 From (8) and (9) we have the circulant determinant

$$\exp(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j) = \begin{vmatrix} S_1 & S_n & \cdots & S_2 \\ S_2 & S_1 & \cdots & S_3 \\ \cdots & \cdots & \cdots & \vdots \\ S_n & S_{n-1} & \cdots & S_1 \end{vmatrix} = 1 \tag{10}$$

If $S_i \neq 0$, where $i = 1, 2, \dots, n$, then (10) has infinitely many rational solutions.

Assume $S_1 \neq 0$, $S_2 \neq 0$, $S_i = 0$ where $i = 3, 4, \dots, n$. $S_i = 0$ are $n - 2$ indeterminate equations with $n - 1$ variables. From (6) we have

$$e^A = S_1 + S_2, \quad e^{2B_j} = S_1^2 + S_2^2 + 2S_1S_2(-1)^j \cos \frac{j\pi}{n} \tag{11}$$

From (10) and (11) we have the Fermat equation

$$\exp(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j) = (S_1 + S_2) \prod_{j=1}^{\frac{n-1}{2}} (S_1^2 + S_2^2 + 2S_1S_2(-1)^j \cos \frac{j\pi}{n}) = S_1^n + S_2^n = 1 \tag{12}$$

Example[1]. Let $n = 15$. From (3) we have

$$A = (t_1 + t_{14}) + (t_2 + t_{13}) + (t_3 + t_{12}) + (t_4 + t_{11}) + (t_5 + t_{10}) + (t_6 + t_9) + (t_7 + t_8)$$

$$B_1 = -(t_1 + t_{14}) \cos \frac{\pi}{15} + (t_2 + t_{13}) \cos \frac{2\pi}{15} - (t_3 + t_{12}) \cos \frac{3\pi}{15} + (t_4 + t_{11}) \cos \frac{4\pi}{15} \\ - (t_5 + t_{10}) \cos \frac{5\pi}{15} + (t_6 + t_9) \cos \frac{6\pi}{15} - (t_7 + t_8) \cos \frac{7\pi}{15},$$

$$B_2 = (t_1 + t_{14}) \cos \frac{2\pi}{15} + (t_2 + t_{13}) \cos \frac{4\pi}{15} + (t_3 + t_{12}) \cos \frac{6\pi}{15} + (t_4 + t_{11}) \cos \frac{8\pi}{15} \\ + (t_5 + t_{10}) \cos \frac{10\pi}{15} + (t_6 + t_9) \cos \frac{12\pi}{15} + (t_7 + t_8) \cos \frac{14\pi}{15},$$

$$B_3 = -(t_1 + t_{14}) \cos \frac{3\pi}{15} + (t_2 + t_{13}) \cos \frac{6\pi}{15} - (t_3 + t_{12}) \cos \frac{9\pi}{15} + (t_4 + t_{11}) \cos \frac{12\pi}{15} \\ - (t_5 + t_{10}) \cos \frac{15\pi}{15} + (t_6 + t_9) \cos \frac{18\pi}{15} - (t_7 + t_8) \cos \frac{21\pi}{15},$$

$$B_4 = (t_1 + t_{14}) \cos \frac{4\pi}{15} + (t_2 + t_{13}) \cos \frac{8\pi}{15} + (t_3 + t_{12}) \cos \frac{12\pi}{15} + (t_4 + t_{11}) \cos \frac{16\pi}{15} \\ + (t_5 + t_{10}) \cos \frac{20\pi}{15} + (t_6 + t_9) \cos \frac{24\pi}{15} + (t_7 + t_8) \cos \frac{28\pi}{15},$$

$$B_5 = -(t_1 + t_{14}) \cos \frac{5\pi}{15} + (t_2 + t_{13}) \cos \frac{10\pi}{15} - (t_3 + t_{12}) \cos \frac{15\pi}{15} + (t_4 + t_{11}) \cos \frac{20\pi}{15} \\ - (t_5 + t_{10}) \cos \frac{25\pi}{15} + (t_6 + t_9) \cos \frac{30\pi}{15} - (t_7 + t_8) \cos \frac{35\pi}{15},$$

$$B_6 = (t_1 + t_{14}) \cos \frac{6\pi}{15} + (t_2 + t_{13}) \cos \frac{12\pi}{15} + (t_3 + t_{12}) \cos \frac{18\pi}{15} + (t_4 + t_{11}) \cos \frac{24\pi}{15} \\ + (t_5 + t_{10}) \cos \frac{30\pi}{15} + (t_6 + t_9) \cos \frac{36\pi}{15} + (t_7 + t_8) \cos \frac{42\pi}{15},$$

$$\begin{aligned}
B_7 = & -(t_1 + t_{14}) \cos \frac{7\pi}{15} + (t_2 + t_{13}) \cos \frac{14\pi}{15} - (t_3 + t_{12}) \cos \frac{21\pi}{15} + (t_4 + t_{11}) \cos \frac{28\pi}{15} \\
& - (t_5 + t_{10}) \cos \frac{35\pi}{15} + (t_6 + t_9) \cos \frac{42\pi}{15} - (t_7 + t_8) \cos \frac{49\pi}{15}, \\
A + 2 \sum_{j=1}^7 B_j = & 0, \quad A + 2B_3 + 2B_6 = 5(t_5 + t_{10})
\end{aligned} \tag{13}$$

Form (12) we have the Fermat equation

$$\exp(A + 2 \sum_{j=1}^7 B_j) = S_1^{15} + S_2^{15} = (S_1^5)^3 + (S_2^5)^3 = 1 \tag{14}$$

From (13) we have

$$\exp(A + 2B_3 + 2B_6) = [\exp(t_5 + t_{10})]^5 \tag{15}$$

From (11) we have

$$\exp(A + 2B_3 + 2B_6) = S_1^5 + S_2^5 \tag{16}$$

From (15) and (16) we have the Fermat equation

$$\exp(A + 2B_3 + 2B_6) = S_1^5 + S_2^5 = [\exp(t_5 + t_{10})]^5 \tag{17}$$

Euler proved that (14) has no rational solutions for exponent 3[8]. Therefore we prove that (17) has no rational solutions for exponent 5[1].

Theorem 1. [1-7]. Let $n = 3P$, where $P > 3$ is odd prime. From (12) we have the Fermat's equation

$$\exp(A + 2 \sum_{j=1}^{\frac{3P-1}{2}} B_j) = S_1^{3P} + S_2^{3P} = (S_1^P)^3 + (S_2^P)^3 = 1 \tag{18}$$

From (3) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{P-1}{2}} B_{3j}) = [\exp(t_P + t_{2P})]^P \tag{19}$$

From (11) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{P-1}{2}} B_{3j}) = S_1^P + S_2^P \tag{20}$$

From (19) and (20) we have the Fermat equation

$$\exp(A + 2 \sum_{j=1}^{\frac{P-1}{2}} B_{3j}) = S_1^P + S_2^P = [\exp(t_P + t_{2P})]^P \tag{21}$$

Euler proved that (18) has no rational solutions for exponent 3[8]. Therefore we prove that (21) has no rational solutions for $P > 3$ [1, 3-7].

References

1. Jiang, C-X, Fermat last theorem had been proved, Potential Science (in Chinese), 2.17-20 (1992), Preprints (in English) December (1991). <http://www.wbabin.net/math/xuan47.pdf>.
2. Jiang, C-X, Fermat last theorem had been proved by Fermat more than 300 years ago, Potential Science (in Chinese), 6.18-20(1992).
3. Jiang, C-X, On the factorization theorem of circulant determinant, Algebras, Groups and Geometries, 11. 371-377(1994), MR. 96a: 11023, <http://www.wbabin.net/math/xuan45.pdf>.
4. Jiang, C-X, Fermat last theorem was proved in 1991, Preprints (1993). In: Fundamental open problems in

- science at the end of the millennium, T.Gill, K. Liu and E. Trelle (eds). Hadronic Press, 1999, P555-558. <http://www.wbabin.net/math/xuan46.pdf>.
5. Jiang, C-X, On the Fermat-Santilli theorem, Algebras, Groups and Geometries, 15. 319-349(1998).
 6. Jiang, C-X, Complex hyperbolic functions and Fermat's last theorem, Hadronic Journal Supplement, 15. 341-348(2000).
 7. Jiang, C-X, Foundations of Santilli Isonumber Theory with applications to new cryptograms, Fermat's theorem and Goldbach's Conjecture. Inter, Acad. Press. 2002. MR2004c:11001, <http://www.wbabin.net/math/xuan13.pdf>. <http://www.i-b-r.org/docs/jiang.pdf>.
 8. Ribenboim, P, Fermat last theorem for amateur, Springer-Verlag, (1999).

5/7/2017