



THE PREVALENCE OF BOVINE TRYPANOSOMOSIS AND ASSOCIATED RISK FACTORS IN BULLEN DISTRICT OF METEKEL ZONE OF BENISHAGUL GUMUZ REGIONAL STATE

Endalkacho Mekonen and Asmamaw Aki*

Assosa, Regional Veterinary Diagnostic, Surveillance, Monitoring and Study Laboratory, P.O. Box 326, Assosa, Ethiopia; asmamawaki@gmail.com, phone: 0902330029

ABSTRACT: A cross-sectional study was carried out in Bullen District of Benishangul Gumuz Regional State, western Ethiopia from December 2020 to January, 2021 to determine the prevalence of trypanosomosis in Bovine and the prevailing species of trypanosomes, associated risk factors and its vector density. Blood samples were collected from (n=384) randomly sampled cattle and examined using parasitological (buffy coat technique) and hematological (measurement of packed cell volume) procedures. An overall, 10.15% (39/384) prevalence was recorded. The infection was caused by *T. congolense* 32/48 (66.66%), *T. vivax* 9/48 (18.75%), *T. brucei* 3/48 (6.25%) and mixed infection was found to be 4/48 (8.33 %). The infection rate was found statistically significant ($P < 0.000$) among trypanosome species. Mean packed cell volume (PCV) value of infected animals was lower (19.06%) than non-infected animals (26.01%) and the variation was statistically significant ($P < 0.000$). Non-significant difference was recorded within study sites, sex and age categories of animals ($P > 0.05$), where as significant association was observed in body conditions. *Glossina tachnoides* were the tsetse fly caught and its mean apparent density measured as fly per trap density was 2.99. In addition, other mechanical vectors such as Stomoxys, tabanids and haematopota with f/t/d of 1.77, 0.26 and 0.25 were recorded respectively. In conclusion, the result of the current study showed the economical importance of trypanosomosis in the study area signaling for strategic control efforts.

[Endalkacho Mekonen and Asmamaw Aki. **THE PREVALENCE OF BOVINE TRYPANOSOMOSIS AND ASSOCIATED RISK FACTORS IN BULLEN DISTRICT OF METEKEL ZONE OF BENISHAGUL GUMUZ REGIONAL STATE.** *Researcher*2023;15(11):40-48]. ISSN 1553-9865 (print); ISSN 2163-8950 (online). <http://www.sciencepub.net/researcher>. 07.doi:[10.7537/marsrsj151123.07](https://doi.org/10.7537/marsrsj151123.07).

Key words: Blood, Packed cell volume, risk factor, Trypanosomosis, and Tsetse fly

1. Introduction

Trypanosomiasis is a devastating disease of livestock caused by protozoal parasites of the genus trypanosoma that inhabits blood and other tissues of vertebrates including animals, wildlife and human (Adam *et al.*, 2003; Gupta *et al.*, 2009; Bal *et al.*, 2014). It is a vector borne disease that is transmitted biologically by tsetse flies and mechanically by other biting flies (FAO, 2002; OIE, 2009). It is a major constraint contributing to direct and indirect economic losses to crop and livestock production (Abebe, 2005) and has a significant negative impact on economic growth in many parts of the world (Taylor *et al.*, 2007; Sharma *et al.*, 2013), particularly in sub-Saharan Africa (Cecchi *et al.*, 2008).

The Diagnosis of trypanosome infection is based on clinical signs; but the clinical signs of the African Animal trypanosomosis are indicative but are not sufficiently pathognomonic. Therefore, standard

methods have been developed and applied practically to diagnose the disease in animals. The methods include: direct microscopic examination of blood, either by the wet film method; but it is insensitive (Getachew, 2005). Stained thin and thick smear techniques permit detailed morphological studies and identification of different *Trypanosoma* species by light microscopy. Sensitivity can be improved through parasitological buffy coat techniques of concentration of the parasites by centrifugation and blood inoculating into susceptible laboratory animals (Getachew, 2005).

As revealed by different studies, tsetse transmitted animal trypanosomosis is a serious constraint to livestock production and agricultural development, exorcising farmers and livestock keepers out of areas having very high potential for growth, and forcing them to live on a highly degraded highlands of Ethiopia (Abebe, 2005). The problem caused by tsetse and trypanosomosis is not only limited to inflicting diseases

but also leading to significant negative impacts such as losses due to mortality and morbidity in domestic animals, cost of livestock treatment and tsetse control, and getting rid of draught animals from their infestation areas (Juyal *et al.*, 2005).

Five species of Glossina (*G. morsitans submorsitans*, *G. Pallidipes*, *G. tachnoides*, *G. f. fuscipes* and *G. longipennis*) have been registered in Ethiopia (Keno, 2005). The tsetse flies (vectors), *G. fusca*; the bush fly, *G. morsitans*, which inhibit principally savannah area and *G. palpalis*; a riverine species, effectively prevent the rearing of the cattle over the large area of the Africa (Blood *et al.*, 1989). The most important trypanosome species affecting livestock in Ethiopia are *Trypanosoma congolense*, *Trypanosoma vivax*, and *Trypanosoma brucei* in cattle, sheep and goats, *Trypanosoma evansi* in camels and *Trypanosoma equiperdium* in horses (Abebe G, 2005).

Benishangul Gumuz is one of the five regions of Ethiopia infested with more than one species of tsetse flies (Keno, 2005). And nearly 31,000 km² or 62% of the region's total land area is believed to be infested with tsetse fly (NTTICC, 1996). In addition to this, the major livestock diseases such as pasteurellosis, black leg and parasitism due to both internal and external parasites are identified in the region, Despite this fact, very scant information is available about the disease epidemiology and its vector with baseline data in the Bullen district. The aims of the present study was, therefore, to assess the epidemiology of trypanosomosis and its vector density in six kebeles of Bullen district.

Therefore, the objectives of the study were

- To determine the prevalence and assess associated risk factor of bovine trypanosomosis
- To determine apparent density of tsetse and other biting flies.

3. Materials and Methods

2.1 Study area

The study was conducted from December 2020 to January 2021 in Bullen district of Benishangul Gumuz Regional State, western part of Ethiopia. Area lies at latitude of 10° and 36'15.1 N and longitude of 036° and 04'52.1'' E at an altitude of 1465 meter above sea level. Annual average temperature of area is 29.5°C and its rainfall range is 900 to 1100 mm, and land size of the area is about 3252,397 km². The major Agricultural activity in the area is mixed farming system whereby crops are cultivated and different species of livestock are kept (NMSA, 2007). The district has 19 kebeles covering an area of 3252.397 km² with human population of 46,920. Extensive livestock husbandry, feeding and outdoor housing system was found, Back yard chicken Management system is practiced, where local breeds are allowed to

scavenge, what nature provides them, In the woreda animal movement was due to agriculture and trade purpose. Animal in the area was used for meat, milk, ploughing and income generation. Socio economy of the people in the area was mainly depending on mixed farming. Mixed agriculture is a common practice with livestock population of 47218 cattle, 16367 sheep, 6392 goats, 5211 equines, 51089 poultry and 1420 beehives (CSA, 2016 & Bullen woreda agriculture office, 2016).

2.2 Study Design and Study Animals

Cross- sectional study design was used. A local zebu cattle, which are usually kept under an extensive husbandry system grazing the communally owned pasture land throughout the year were randomly sampled. They grazed together during the day time and returned to their individual owner's farmstead each evening. The body condition of each of the study cattle was scored as good, medium and poor as designed by Nicholson M, and Butterworth M (1986). Concurrently, their age was categorized in years (< 2, 2-7, >7) based on De-Lahunta A, and Habel RE (1986).

2.3 Sampling Techniques and Sample Size Determination

This study was conducted in six peasant associations including Bullen town, mora, mata, Emange, Dobi and mati. The study sites were selected purposively as convenient, particularly based on accessibility, agro ecology, and presence of cattle. The animals were sampled randomly involving both sexes, all age groups, and all types of body conditions. The desired sampling size was calculated according to the formula given by (Thrusfield, 2005). The sample size was determined based on the expected prevalence of 50%; confidence level of 95%, and 5% desired absolute precision.

So, it was calculated by using the formula

$$n = \frac{1.96^2 * 50\% (1 - 50\%)}{(5\%)^2} = \frac{3.84 * 0.5(1-0.5)}{(0.05)^2} = 384$$

As result a total of 384 cattle were calculated, and these cattle were sampled at their communal grazing area using simple random sampling.

2.4 Study methods

2.4.1 Packed cell volume (PCV) determination

Blood samples were obtained by puncturing the marginal ear vein with a lancet and collected directly into a pair of heparinized capillary tubes. The tubes were then sealed at one end with crystal seal. PCV was measured in a micro-haematocrit centrifuge (Hermmlle Labortechnik, type Z, Germany). The capillary tubes were placed in microhematocrits centrifuge with sealed end outermost. Then, the tube was loaded symmetrically to ensure good balance. After screwing the rotary cover and closing the centrifuge lid, the

specimens were allowed to centrifuge at 12,000 rpm for 5 minutes. After centrifugation, the capillary tubes were placed in a haematocrit reader. The length of the packed red blood cells column is expressed as a percentage of the total volume of blood. Animals with PCV less than 24% were considered to be anemic (OIE, 2008).

2.4.2 Buffy coat technique

Heparinized microhaematocrit capillary tubes, containing blood samples were centrifuged for 5 minutes at 12,000 rpm. After the centrifugation, trypanosomes were usually found in or just above the buffy coat layer. The capillary tube was cut using a diamond tipped pen 1 mm below the buffy coat to include the upper most layers of the red blood cells and 3 mm above to include the plasma. The content of the capillary tube was pour onto a glass slide, and covered with cover slip. The slide was examined under x40 objective and x10 eye piece for movement of parasite (Paris, 1982). Trypanosome species were identified according to their morphological descriptions as well as movement in wet film preparations (OIE, 2008).

2.4.3 Entomological survey

A total of 56 odour baited traps of monoconical, biconical and mono pyramidal were deployed at 200-250 m intervals to assess the density and species of tsetse flies during the study. Each and every trap was odour baited with acetone and cow urine. The underneath of each trap pole was smeared with grease in order to prevent ants climbing up the pole towards the collecting cage that could damage the tsetse flies. The trap deployment time was 48 hours. After capturing the flies in the collecting cage, they were then sorted by sex and species and recorded. The species of tsetse was identified based on the characteristic morphology. Other biting flies were also separated according to their morphological characteristics such as size, color, proboscis and wing venation structures at the genus level (Fischer and Say, 1989). Sexing was done for tsetse fly just by observing the posterior end of the ventral aspect of abdomen by hand lens, accordingly, male flies were easily identified by enlarged hypophageum. The apparent density of tsetse flies was determined based on the daily mean number of flies captured in odour-baited traps and recorded as fly per trap per day (F/T/D) (Leak *et al.*, 2009).

2.5 Data management and Analysis

All the collected raw data and the results of parasitological and hematological examination data were entered into a Microsoft excel spread sheets program and then was transferred to Intercool STATA version 10.0 for analysis. The prevalence of

trypanosome infection was calculated as the number of positive animals as examined by buffy coat method divided by the total number of animals examined at the particular time. Data collected on PCV values were analyzed by ANOVA one to compare the mean PCV values of infected animals against that of uninfected animals. Pearson's chi-square (χ^2) was used to evaluate the association of different variables with the prevalence of trypanosome infection. In all of the statistical analysis, a confidence level of 95% is used and P-value of less than 0.05 (at 5% level of significance) was considered as statistically significant.

4. RESULT

4.1 Parasitological survey result

Out of the total animals examined (n=384), 39/384(10.15 %) were found to be infected with trypanosomes (Table-5). The prevalence in terms of trypanosome species was 8.33% for *T. congolense*, 1.56 % for *T. vivax*, 0.52% for *T. brucei* and 0.78 % was found to be mixed infection. The proportion of trypanosome species was 32/43(74.41%) for *T. congolense*, 6/ 43(13.95%) for *T. vivax*, 2/43(4.65%) for *T. brucei* and 3/43 (6.97%) for mixed infection and the infection rate was found to be statistically significant (P<0.000) among trypanosome species (Table 4).

Table 1. Species based prevalence of bovine trypanosomosis in Bullen district

Trypanosomes	No. positive	Positive (%)	X ² (p-value)
<i>T. congolense</i>	32	74.41	308.81 (P<0.000)
<i>T. vivax</i>	6	13.95	
<i>T. brucei</i>	2	4.65	
Mixed(<i>T. congolense</i> & <i>T. vivax</i>)	3	6.97	
Total	43	100	

4.2 Haematological survey result

The mean PCV value for all examined animals was 23.02. However, the mean PCV value for non infected and infected animals was 26.01 and 19.06 respectively. The mean PCV values of cattle were significantly ($P < 0.000$) influenced by trypanosome infection as 19.06 % and 26.01 % PCV values in trypanosome positive and negative animals were registered, respectively (Table 7).

4.3 Risk factors

The highest prevalence (11.80%) of trypanosomosis was recorded in animals >7 years old whilst the lowest prevalence (8.63 %) was recorded in animals 2-7 years

of old and the association was not found statistically significant among the age groups (Table 6). Slightly higher prevalence was registered in female animals (11.16 %) than in male animals (8.98 %), which was not found to be statistically significant ($p > 0.05$) (Table 6). Trypanosomosis was recorded across the study sites with the highest and lowest prevalence of (18.29%) and (4.54 %) in mati and matha respectively and prevalence of trypanosomosis was not significant across the study

sites (Table 5). The highest prevalence of trypanosomosis (16.82%) was found in animals with poor body condition while the lowest (6.89 %) was recorded in animals with medium body conditions respectively, and the difference was significant ($p < 0.000$). The effect of age, sex, sites and body condition on prevalence of trypanosomosis is summarized in Table 5 and Table 6.

Table 2: Origin based Prevalence of bovine trypanosomosis in Bullen District

Sites(PA)	No. examined	No. positive	(%) positive	χ^2 (p-value)
Bullen town	72	7	9.72	9.40(P>0.09)
Mora	64	4	6.25	
Matha	44	2	4.54	
Emange	81	6	7.40	
Dobi	41	5	12.19	
Mati	82	15	18.29	
Total	384	39	10.15	

Pa: peasant association

Table 6: Prevalence of trypanosomosis with different potential risk factors

Risk factors	No. examined	No. positive	(%) positive	χ^2 (p-value)
Sex				0.49(P>0.05)
Male	178	16	8.98	
Female	206	23	11.16	
Total	384	39	10.15	
Age(years)				0.78(P>0.05)
< 2	101	10	9.90	
2 – 7	139	12	8.63	
> 7	144	17	11.80	
Total	384	39	10.15	
Body conditions				7.32(P<0.02)
Good	161	13	8.07	
Medium	116	8	6.89	
Poor	107	18	16.82	
Total	384	39	10.15	

Table 7: Mean PCV comparison of parasitaemic and aparasitaemic animals

Status	Frequency	Mean PCV (%)	SE	X ² (p-value)
Infected	179	19.06	02.9	25.18(p<0.000)
Non-infected	205	26.01	01.18	
Total	384	23.02	01.54	

PCV: packed cell volume; SE: standard Error

4.4 Entomological survey result

The present survey of tsetse flies depicted that *G. tachinoide* is the only species of tsetse fly responsible for cyclical transmission of trypanosomosis in the study area. Tsetse fly survey was carried out in six kebele of the study district by deploying a total of 56 geo-referenced traps (16 mono-conical, 20 mono-pyramidal and 20 biconical traps) in the river border, open wood land (savanna grass land) and on grazing fields of cattle, the number of tsetse flies captured in each study site after 48 hour is 109, 79, 68, 92, 122,

142 for Bullen town, mora, matha, Emange, Dobi, mati respectively. A total of 612 tsetse and biting flies were caught from different sites during the study period. The mean apparent density of *G. tachinoide* in the survey sites was investigated as 2.99 f/t/d while the mean apparent density of mechanical vectors such as stomoxys (1.77 f/t/d), tabanids (0.26 f/t/d) and haematopota (0.25 f/t/d) were recorded. The highest fly density was observed in mati peasant association (7.88 f/t/d) and the lowest was recorded in mata (3.4 f/t/d) (Table 8).

Table 8: Flies caught in different areas of survey sites at Bullen district

Sites	Total flies caught	No. of traps	Tsetse flies caught					Biting flies		
			No.	species	M	F	*F/T/D	Stomoxys	Tabanid	Haematopota
Bullen town	109	10	52	GT	22	30	2.6	30	4	3
Mora	79	9	44		15	29	2.4	26	5	4
Matha	68	10	30		11	19	1.5	28	7	3
Emange	92	9	61		22	39	3.38	20	5	6
Dobi	122	9	71		26	45	3.94	43	3	5
Mati	142	9	71		28	49	4.27	52	6	7
Total	612	56	335		124	211	2.99	199(1.77)	30(0.26)	28(0.25)

F/T/D=fly per trap per day, GT= Glossina tachinoides, M=male, F=female

5. DISCUSSION

10.15 % of Cattle trypanosomosis prevalence were reported in the study area. This finding was agree with the study conducted by (Aki A *et al.*, 2015) Who reported 8.96% bovine trypanosomosis prevalence in Kameshi District, Benishagul Gumuz region, western Ethiopia. In contrast, 22.38 % bovine trypanosomosis prevalence was reported by Bayisa *et al.* (2015) in Asossa district, which was high as compared to the current study. Similarly, 26.30% cattle trypanosomosis prevalence was reported by Aki A *et al.* (2017) in Mandura district which were higher than the present findings.

This research showed that the infection was predominantly caused by *T. congolense* 32/43

(74.41%), *T.vivax* 6/43(13.95%), and *T. brucei* 2/43(4.65%) and mixed infection 3/43(6.97%). This result is in line with the reported proportions of *T.congolense* (75.5 %) followed by *T.vivax* (14.28%) from Metekel and Awi zones (Mekuria *et al.*, 2011). This result was also in consistent with prior reports of (Mulaw *et al.*,2011) who studied on prevalence of major trypanosomes affecting cattle in Assosa district of Benishangul Gumuz Regional State, western Ethiopia and who found proportional prevalence of *T. congolense* to be 66.7% ; (Abraham *et al.*, 2012) conducted their study on prevalence of bovine trypanosomosis in selected sites of Arba Minch district, Southern Ethiopia whose result showed proportional prevalence of *T. congolense* to be 61.4%; (Biyazen *et*

al., 2014) reported proportional prevalence of *T. congolense* to be 63.64% during their work on trypanosomosis and anemia in cattle population of Dale Wabera district of Kellem Wollega Zone, western Ethiopia.

The high proportional infection rate of *T. congolense* in cattle might be attributable to the high number of serodemes of *T. congolense* relative to other species of trypanosomes. It could also be due to the possible development of better immune response to *T. vivax* by the infected animals as demonstrated by (Leak *et al.*, 1993). Further, it might be attributed to the efficient transmission of *T. congolense* by cyclical vectors than *T. vivax* in tsetse-infested areas. Previous reports indicated that *T. congolense* and *T. vivax* are the most prevalent trypanosomes that infect cattle in tsetse infested and tsetse free areas of Ethiopia respectively (Leak *et al.*, 1999). Different studies (Leak *et al.*, 1993; Rowland *et al.*, 1995) have indicated that *T. vivax* is highly susceptible to treatment while the problems of drug resistance are higher in *T. congolense*.

The effect of different risk factors such as sex, age categories, study sites, PCV status, trypanosome species, and body conditions on prevalence of cattle trypanosomosis was studied and, statistically significant associations were observed in, PCV status, body conditions and trypanosomes species ($p < 0.05$) while sex groups, age categories and study sites were not found to be statistically significant ($p > 0.05$). This result is in agreement with previous reports of (Lelisa *et al.*, 2015 and Bayisa *et al.*, 2015).

The overall mean PCV value for examined animals was 23.02. The mean PCV value of infected animals was significantly lower (19.06) than that of non - infected animals (26.01). This result is in alignment with previous works of (Ali *et al.*, 2011; Mulaw, 2011).

Entomological survey result indicated that, *G. tachnoide* was the only tsetse fly caught and its mean apparent density measured as f/t/d was found to be 2.99. It accounts for 54.73 % (335/612) out of the total flies caught. The highest fly density was observed in mati peasant association (7.88 f/t/d) and the lowest was recorded in mata (3.4 f/t/d). 2.99 (f/t/d) Tsetse fly, 1.77(f/t/d) stomoxy, 0.26 (f/t/d) tabanidae, 0.25 (f/t/d) hematopota were investigated in this research. The current finding is concord with the previous findings of (NTTICC, 2012-2014) at neighboring mandura district of western Ethiopia which was reported to be 3.59 f/t/d, 1.38 f/t/d, 0.33 f/t/d and 0.014 f/t/d, for tsetse fly, *stomoxys*, *haematopota*, and *tabanus* respectively. Similarly, this finding was inline with the previous findings of (Aki A *et al.*, 2015) in Kamshi Distirict, which was reported to be 2.54, 2.84, 1.54, and 0.92 for

G. tachnoides, stomoxys, tabanids and haematopota respectively.

6. CONCLUSION AND RECOMMENDATIONS

The high prevalence of Cattle trypanosomosis (10.15 %) were remains a major problem that hinders livestock production and productivity in the district. The most widely distributed and dominant species of trypanosomes in the study sites are *T. congolense* (74.41%) followed by *T. vivax* (13.95%), and to some extent *T. brucei* (4.65 %) which was mainly transmitted by *Glossina tachnoide* and other biting flies with f/t/d/ of 2.99, 1.77, 0.26 and 0.25 for *G. tachnoides*, *stomoxys*, *tabanids* and *haematopota* respectively. Parameters of study animals such as sex , sites, and age were not found to be a risk factor for trypanosomosis whereas body conditions and trypanosome species were risk factors. This study showed that trypanosome infection and other factors such as (nutritional, seasonal, concurrent disease) was found to negatively affect the PCV values of animals. Based on the current findings, the following recommendations are forwarded:-

- Particular attentions towards the identified trypanosome species are essential to control the impact of the disease on cattle that are potential reservoir of the infections.
- Development of control options that could minimize the tsetse fly and biting flies in the study area should be introduced in a wholistic approach.
- Proper and strict follow up of trypanocidal drug distribution, therapeutic strategies and alternative control measures should be implemented by concerned stake holders.
- The farmer in the area should be trained how to control the vector of the disease and provided with materials
- Further study on the trypanosomosis, tsetse fly investigation and also on possible factors should be carried out by laboratory experts to give the best strategic control and prevention measures in the study area.

Corresponding Author:

Dr. Asmamaw Aki
Regional Veterinary Diagnostic and Research Laboratory
Telephone: +251 0902330029;
Email: asmamawaki@gmail.com

7. REFERENCE

- [1]. Abebe G., (2005): Current situation of Trypanosomosis. In: review article on:

- Trypanosomosis in Ethiopia. *Ethiop. J. Biol.med. Sci.* **4**(1): 75-121
- [2]. Adam K., Paul J., Zaman V., (2003): Medical and Veterinary Protozoology. Churchill living stone Edinburgh and London.
- [3]. Abraham Z., and Zeryehun T., (2012): Prevalence of Bovine Trypanosomosis in Selected District of Arba Minch, Snnpr, Southern Ethiopia, *Glob. Vet.* **8**(2): 168-173.
- [4]. Aki A., (2016): Epidemiology of bovine Trypanosomosis in Kamashi District of Benishangul Gumuz Regional State, Western Ethiopia: Prevalence, Vector density and associated risks. *J. Vet. Adv.*, **2**(4): 48-54.
- [5]. Ali D., and Bitew M.,(2011): Epidemiological study on bovine trypanosomiasis in Maokomo special district Benishangul-Gumuz regional State, Western Ethiopia. *Glob. Vet.* **6**: 402-408.
- [6]. Awoke K., (2000): Study of Trypanosomosis and its Vector in Humba and Merab Woreda of Eastern Ethiopia: *J. Ethio. Vet. Assoc.*, **9**:81-83.
- [7]. Bal M., Sharma A., Ashuma B., Kaur P., and Singla L., (2014): Detection and management of latent infection of *Trypanosoma evansi* in a cattle herd. *Ind. J. Anim. Res.* **48**(1): 31-37.
- [8]. Biyazen H., Duguma R., and Asaye M., (2014): Trypanosomosis, Its Risk Factors, and Anaemia in Cattle Population of Dale Wabera District of Kellem Wollega Zone, Western Ethiopia, *J. Vet.Med.*
- [9]. Bayisa K., Getachew D., Tadele T., (2015): Bovine Trypanosomosis in Asossa District, Benishangul Gumuz Regional State, Western Ethiopia: Prevalence and Associated Risk Factors, *Euro. J. App. Sc.* **7**(4): 171-175, 2015.
- [10]. Blood D., Radostits O., and Henderson J., (1989): Diseases caused by Protozoa. A textbook of the Diseases of cattle, pigs, goats and horses. *7th ed.* Oxford: ELBS.PP.1012-015.
- [11]. Brown C., Hunter D., and Lukins A., (1990): Disease caused by Protozoa. Hand book on Animals Disease in Tropics (Swell, and Brocklesby), *4th Ed.* Bailer. Tindall, and London.
- [12]. Brown C., (1991): Diseases caused by protozoa. In: Handbook on animal diseases in the tropics. (Sewell, M. M. H. and Brocklesby, D. W. eds.) 4th edition. British Veterinary Association.
- [13]. Cecchi G., Mattioli R., Slingenbergh J., de la Rocque S., (2008): Land cover and tsetse fly distributions in sub-Saharan Africa. *Med. Vet. Entom.* **22**: 364-373.
- [14]. Cherenet T., Sani R., Panandam J., Nadzr S., Speybroeck N., Van Den Bossche P., (2004): Seasonal prevalence of bovine trypanosomiasis in a tsetse-infested zone and a tsetse-free zone of [15]. the Amhara region, north-west Ethiopia. Onderstepoort *J. Vet. Res.* **71**(4): 307-312.
- [16]. CSA (Central Statistical Authority), (2015): Agricultural Sample Survey, Statistical Bulletin, Ethiopia, Addis Ababa, pp: 39-47.14. Nicholson M, and Butterworth M.
- [17]. De-Lahunta A., and Habel R., (1986): Teeth. Applied veterinary Anatomy. USA. W. B. Saunders. Company, pp: 4-16.
- [18]. Getachew A., (2005): Trypanosomosis in Ethiopia, A.A.U, faculty of veterinary medicine. Debre zeit, pp 18-20.
- [19]. Gupta M., Kumar H., and Singla L., (2009): Trypanosomiasis concurrent to tuberculosis in black bucks. *Ind. Veter. J.* **86**: 727-728.
- [20]. Murray M., and Dexter T., (1988): Anemia in Bovine in African Animal Trypanosomosis.
- [21]. *Acta. Top-45*: 389-432.
- [22]. Keno M., (2005): The current situation of tsetse and trypanosomosis in Ethiopia, Ministry of Agriculture and Rural Development, Veterinary service department, in proceeding of 28th meeting of International Scientific Council for Trypanosomosis Research and Control.
- [23]. Mulaw S., Addis M., and Fromsa A., (2011): Study on the Prevalence of Major Trypanosomes Affecting Bovine in Tsetse Infested Asossa District of Benishangul Gumuz Regional State, Western Ethiopia. *Glob. Vet.* **7** (4): 330- 336.
- [24]. Mekuria S, and Gadissa F (2011): Survey on bovine trypanosomosis and its vector in Metekel [25]. and Awi zones of North west Ethiopia. *Acta Tropica*, **117**: 146-151.
- [26]. Murray M., Murray P., and McIntyre W., (1977): An improved parasitological diagnostic technique for the diagnosis of African trypanosomiasis. *Transactions of the royal society of tropical medicine and hygiene.* *Vol.* **71** pp. 325-326
- [27]. Murray M., Barry J., Morrison W., Williams., Hirumi H., and Rovis L., (1979): A review of the prospects of vaccination in African trypanosomiasis. Part I. (32) pp. 9-13
- [28]. Murray M., and Gray A., (1984): The current situation on animal trypanosomiasis in Africa. *Prev. Vet.Med.* *Vol.* **2** pp. 23-30
- [29]. Murray M., Steer M., Trail J., D'letern., Agyemang K., and Dwinger R., (1991): Trypanosomiasis in cattle: Prospects for control. Breeding for disease resistance in farm animals. (Cowen, B. and Axford, R.F.E. eds.) CAB International, Wallingford, chapter12 pp. 203-223
- [30]. NTTICC (National Tsetse and Trypanosomosis Investigation and Control Centre), (1996):

- [31]. Annual Report on Tsetse and Trypanosomosis, Survey, Bedele, Ethiopia. Pp.11-15.
- [32]. NTTICC (2004): National Tsetse and Trypanosomosis Investigation and Control Center. Report for the period 7th June 2003-6th July 2004. Bedelle, p. 3.
- [33]. NMSA (National Meteorological Services Agency), (2014): Monthly report on temperature and Rainfall distribution for Kamashi Zone, Regional Metrological Office, Assosa, Ethiopia, pp: 17-19.
- [34]. Langridge W., Tsetse and Trypanosomosis Survey of Ethiopia. Ministry of Overseas Department UK, 1976; Pp.1-40.
- [35]. OIE (2009): Manual of standards for diagnostic tests and vaccines for terrestrial animals, 6th ed. Paris. pp: 813-2008
- [36]. OIE (Office of Internationale des Epizootics), (2008): "Standardized techniques for the diagnosis of tsetse transmitted trypanosomosis," in *OIE Terrestrial Manual*, p. 49, Rome, Italy.
- [37]. FAO (1998): A field guide for the diagnosis, treatment and prevention of African Trypanosomes: In Food and Agriculture organization of the United Nations Rome FAO. <http://www.Fao.org/DOCREP/006/X0413/X0413E00.HTM>. Pp.170.
- [38]. Nicholson M., and Butterworth M., (1986): A guide to condition scoring of zebu cattle. ICCA, Addis Ababa, Ethiopia.
- [39]. Jemere B., (2004): Control of Tsetse and Trypanosomiasis in the southern Rift valley (step area): Evaluation of Deltamethrin Applications, Msc thesis, faculty of vet. Med. AAU., Debrezeit, Ethiopia.
- [40]. Urquhart G., Armou J., Dunn A., and Jenings F., (1996): *Veterinary Parasitology 2nd ed.* Black Well publishing. Pp. 212-217.
- [41]. IAEA (2002): Campaign launched to eliminate Tsetse fly; which has turned much of Africa in to a Green Desert Press Reports on PATTEC.
- [42]. Thrusfield M., (2005): *Veterinary Epidemiology*, 3rd edition, Blackwell Science Ltd, Oxford, UK. Pp.233-250.
- [43]. Kassa B., (2005): Standard Veterinary Laboratory, Diagnostic manual, and volume 3 Parasitology. Minister of Agriculture and Rural Development, Animal Health Department, Addis Ababa, Ethiopia. Pp 32-37.
- [44]. Lelisa K., Damena D., Kedir M., and Feyera T., (2015): Prevalence of Bovine Trypanosomosis and Apparent Density of Tsetse and Other Biting Flies in Mandura District, Northwest Ethiopia. *J Vet. Sci. Technol* 6: 229.
- [45]. Holmes P., Mammo E., Thomson A., Knight P., Lucken R., Murray P., Murray M., Jennings F., and Urquhart M., (1974): Immunosuppression in bovine trypanosomiasis. *Vet. record vol. 95* pp.86-87.
- [46]. Paris J., Murray M., and Macodimba F., (1982): A comparative evaluation of parasitological techniques currently available for the diagnosis of African trypanosomiasis in cattle. *Acta tropica. Vol. 39* pp. 307-316.
- [47]. Singla L., Aulakh G., Juyal P., Singh J., (2004): Bovine trypanosomosis in Punjab, India. Proceeding of the 11th International Conference of the Association of Institutions for Tropical Veterinary Medicine and 16th Veterinary Association Malaysia Congress, 23-27 August 2004, Pet aling Jaya, Malaysia, pp: 283-285. 4.
- [48]. Jemere B., (2004): Control of Tsetse and Trypanosomiasis in the southern Rift valley (step area): Evaluation of Deltamethrin Applications, Msc thesis , Faculty of Vet. Med. AAU., Debrezeit, Ethiopia.
- [49]. Urquhart G., Armou J., Dunn A., and Jenings F., (1996): *Veterinary Parasitology 2nd ed.* Black Well publishing. Pp. 212-217.
- [50]. Kassa B., (2005): Standard Veterinary Laboratory, Diagnostic manual, and volume 3 Parasitology. Minister of Agriculture and Rrural Development,Animal Health Department. Addis Ababa, Ethiopia. Pp 32-37.
- [51]. Juyal P., Singla L., Kaur P., (2005): Management of surra due to *Trypanosoma evansi* in India: an overview. *Infectious diseases of domestic animals and zoonosis in India.* 75:109-120.
- [52]. Fisher M., Say R., (1989): *Manual of Tropical Veterinary Parasitology.* UK: CAB International publication. Pp.100-278.
- [53]. Leak S., (1999): Tsetse biology and ecology: The role in the epidemiology and control of
- [54]. trypanosomosis. CAB International. Wallingford (UK), pp. 152-210.
- [55]. Radostits O., Gay C., Blood D., and Hinchelift K., (1996): Disease caused by protozoa – *Trypanosomes.* *Veterinary Medicine: A Text Book of Disease of Cattle, Sheep, Pig, Goat and Horses.* 9th ed. *Harcourt Publisher Ltd. London.* Pp 1531-1541.
- [56]. Leak S., Mulatu W., Authie E., D'Ieteren G., Peregrine AS., (1993): Epidemiology of bovine trypanosomosis in the Gibe valley, Southern Ethiopia. Tsetse challenge and its relationship to trypanosome prevalence in cattle. *Acta Tropica,* 53: 1221-1234.

- [57]. Seifert H., (1996): Trypanosomosis in Tropical Animal Health. Kluwer Academic publisher, Dordrecht/ Boston/ London. Pp. 152-168.
- [58]. OAU (Organization of African Union) (2001): Trypanosomosis, Tsetse and Africa. The year book report (2001).
- [59]. Sharma A, Singla L., Ashuma., Batth B., Kaur P., Javed M., Juyal P., (2013): Molecular prevalence of Babesia bigemina and Trypanosoma evansi in dairy animals from Punjab, India by duplex PCR: A step forward to detection and management of concurrent latent infections. *Biomed. Res. Int.* Article ID 893862, 8 pp.
- [60]. Taylor M., R. Coop and R. wall (2007): Veterinary Parasitology 3rd ed. Black Well publishing Ltd, oxford. Uk, Pp 96-102, 212-214.

11/22/2023